

# Iwasawa's Main Conjecture for elliptic curves over anticyclotomic $\mathbb{Z}_p$ -extensions

By M. BERTOLINI\* and H. DARMON\*\*

## Contents

1.  $p$ -adic  $L$ -functions
    - 1.1. Modular forms on quaternion algebras
    - 1.2.  $p$ -adic Rankin  $L$ -functions
  2. Selmer groups
    - 2.1. Galois representations and cohomology
    - 2.2. Finite/singular structures
    - 2.3. Definition of the Selmer group
  3. Some preliminaries
    - 3.1.  $\Lambda$ -modules
    - 3.2. Controlling the Selmer group
    - 3.3. Rigid pairs
  4. The Euler system argument
    - 4.1. The Euler system
    - 4.2. The argument
  5. Shimura curves
    - 5.1. The moduli definition
    - 5.2. The Cerednik-Drinfeld theorem
    - 5.3. Character groups
    - 5.4. Hecke operators and the Jacquet-Langlands correspondence
    - 5.5. Connected components
    - 5.6. Raising the level and groups of connected components
  6. The theory of complex multiplication
  7. Construction of the Euler system
  8. The first explicit reciprocity law
  9. The second explicit reciprocity law
- References

## Introduction

Let  $E$  be an elliptic curve over  $\mathbb{Q}$ , let  $p$  be an ordinary prime for  $E$ , and let  $K$  be an imaginary quadratic field. Write  $K_\infty/K$  for the anticyclotomic  $\mathbb{Z}_p$ -extension of  $K$  and set  $G_\infty = \text{Gal}(K_\infty/K)$ .

---

\*Partially supported by GNSAGA (INdAM), M.U.R.S.T., and the EC.

\*\*Partially supported by CICMA and by an NSERC research grant.

Following a construction of Section 2 of [BD1] which is recalled in Section 1, one attaches to the data  $(E, K, p)$  an anticyclotomic  $p$ -adic  $L$ -function  $L_p(E, K)$  which belongs to the Iwasawa algebra  $\Lambda := \mathbb{Z}_p[[G_\infty]]$ . This element, whose construction was inspired by a formula proved in [Gr1], is known, thanks to work of Zhang ([Zh, §1.4]), to interpolate special values of the complex  $L$ -function of  $E/K$  twisted by characters of  $G_\infty$ .

Let  $\text{Sel}(K_\infty, E_{p^\infty})^\vee$  be the Pontrjagin dual of the  $p$ -primary Selmer group attached to  $E$  over  $K_\infty$ , equipped with its natural  $\Lambda$ -module structure, as defined in Section 2. It is a compact  $\Lambda$ -module; write  $\mathcal{C}$  for its characteristic power series, which is well-defined up to units in  $\Lambda$ .

Let  $N_0$  denote the conductor of  $E$ , set  $N = pN_0$  if  $E$  has good ordinary reduction at  $p$ , and set  $N = N_0$  if  $E$  has multiplicative reduction at  $p$  so that  $p$  divides  $N_0$  exactly. It will be assumed throughout that the discriminant of  $K$  is prime to  $N$ , so that  $K$  determines a factorisation

$$N = pN^+N^-,$$

where  $N^+$  (resp.  $N^-$ ) is divisible only by primes different from  $p$  which are split (resp. inert) in  $K$ .

The main goal of the present work is to prove (under the mild technical Assumption 6 on  $(E, K, p)$  given at the end of this introduction) Theorem 1 below, a weak form of the Main Conjecture of Iwasawa Theory for elliptic curves in the ordinary and anticyclotomic setting.

**THEOREM 1.** *Assume that  $N^-$  is the square-free product of an odd number of primes. The characteristic power series  $\mathcal{C}$  divides the  $p$ -adic  $L$ -function  $L_p(E, K)$ .*

The hypothesis on  $N^-$  made in Theorem 1 arises naturally in the anticyclotomic setting, and some justification for it is given at the end of the introduction.

Denote by  $L_p(E, K, s)$  the  $p$ -adic Mellin transform of the measure defined by the element  $L_p(E, K)$  of  $\Lambda$ . Let  $r$  be the rank of the Mordell-Weil group  $E(K)$ . The next result follows by combining Theorem 1 with standard techniques of Iwasawa theory.

**COROLLARY 2.**  $\text{ord}_{s=1} L_p(E, K, s) \geq r$ .

A program of study of  $L_p(E, K, s)$  in the spirit of the work of Mazur, Tate and Teitelbaum [MTT] is outlined in [BD1], and partially carried out in [BD2]–[BD5]. In particular, Section 4 of [BD1] formulates a conjecture predicting the exact order of vanishing of  $L_p(E, K, s)$  at  $s = 1$ . More precisely, set  $E^+ = E$  and let  $E^-$  be the elliptic curve over  $\mathbb{Q}$  obtained by twisting  $E$  by  $K$ . Write

$r^\pm$  for the rank of  $E^\pm(\mathbb{Q})$ , so that  $r = r^+ + r^-$ . Set  $\tilde{r}^\pm = r^\pm + \delta^\pm$ , where

$$\delta^\pm = \begin{cases} 1 & \text{if } E^\pm \text{ has split multiplicative reduction at } p, \\ 0 & \text{otherwise.} \end{cases}$$

Finally set  $\rho := \max(\tilde{r}^+, \tilde{r}^-)$  and  $\tilde{r} := \tilde{r}^+ + \tilde{r}^-$ . Conjecture 4.2 of [BD1] predicts that

$$(1) \quad \text{ord}_{s=1} L_p(E, K, s) = 2\rho = \tilde{r} + |\tilde{r}^+ - \tilde{r}^-|.$$

This conjecture indicates that  $L_p(E, K, s)$  vanishes to order strictly greater than  $r$ , if either  $\tilde{r} > r$  or if  $\tilde{r}^+ \neq \tilde{r}^-$ . The first source of extra vanishing is accounted for by the phenomenon of exceptional zeroes arising when  $p$  is a prime of split multiplicative reduction for  $E$  over  $K$ , which was discovered by Mazur, Tate and Teitelbaum in the cyclotomic setting [MTT]. The second source of extra vanishing is specific to the anticyclotomic setting, and may be accounted for by certain predictable degeneracies in the anticyclotomic  $p$ -adic height, related to the fact that  $\text{Sel}(K_\infty, E_{p^\infty})^\vee$  fails to be semisimple as a module over  $\Lambda$  when  $r^+ \neq r^-$ . (Cf. for example [BD $\frac{1}{2}$ ].)

A more careful study of the  $\Lambda$ -module structure of  $\text{Sel}(K_\infty, E_{p^\infty})$ , which in the good ordinary reduction case is carried out in [BD0] and [BD $\frac{1}{2}$ ], yields the following refinement of Corollary 2 which is consistent with the conjectured equality (1).

**COROLLARY 3.** *If  $p$  is a prime of good ordinary reduction for  $E$ , then*

$$\text{ord}_{s=1} L_p(E, K, s) \geq 2\rho.$$

Let  $\mathcal{O}$  be a finite extension of  $\mathbb{Z}_p$ , and let  $\chi : G_\infty \rightarrow \mathcal{O}^\times$  be a finite order character, extended by  $\mathbb{Z}_p$ -linearity to a homomorphism of  $\Lambda$  to  $\mathcal{O}$ . If  $M$  is any  $\Lambda$ -module, write

$$M^\chi = M \otimes_\chi \mathcal{O},$$

where the tensor product is taken over  $\Lambda$  via the map  $\chi$ .

Let  $\text{III}_p(E/K_\infty)$  denote the  $p$ -primary part of the Shafarevich-Tate group of  $E$  over  $K_\infty$ . A result of Zhang ([Zh, §1.4]) generalising a formula of Gross established in [Gr1] in the special case where  $N$  is prime and  $\chi$  is unramified, relates  $\chi(L_p(E, K))$  to a nonzero multiple of the classical  $L$ -value  $L(E/K, \chi, 1)$  (where one views  $\chi$  as a complex-valued character by choosing an embedding of  $\mathcal{O}$  into  $\mathbb{C}$ ). Theorem 1 combined with Zhang's formula leads to the following corollary, a result which lends some new evidence for the classical Birch and Swinnerton-Dyer conjecture.

**COROLLARY 4.** *If  $L(E/K, \chi, 1) \neq 0$ , then  $E(K_\infty)^\chi$  and  $\text{III}_p(E/K_\infty)^\chi$  are finite.*

*Remarks.* 1. The restriction that  $\chi$  be of  $p$ -power conductor is not essential for the method that is used in this work, so that it should be possible, with little extra effort, to establish Corollary 4 for arbitrary anticyclotomic  $\chi$ , and for the  $\chi$ -part of the full Shafarevich-Tate group and not just its  $p$ -primary part, by the techniques in the proof of Theorem 1.

2. Corollary 4 was also proved in [BD2] by a different, more restrictive method which requires the assumption that  $p$  is a prime of *multiplicative reduction* for  $E/K$  which is inert in  $K$ . Hence, in contrast to the previous remark, the method of [BD2] cannot be used to obtain the finiteness of the full Shafarevich-Tate group of  $E$ , but only of its  $p$ -primary part for a finite set of primes  $p$ .

3. The nonvanishing of  $L(E/K, \chi, 1)$  seems to occur fairly often. For example, Vatsal has shown ([Va1, Th. 1.4]) that  $L(E/K, \chi, 1)$  is nonzero for almost all  $\chi$  when  $\chi$  varies over the anticyclotomic characters of  $p$ -power conductor for a fixed  $p$ .

Another immediate consequence of Theorem 1 is that  $\text{Sel}(K_\infty, E_{p^\infty})$  is a cotorsion  $\Lambda$ -module whenever  $L_p(E, K)$  is not identically 0, so that in particular one has

**COROLLARY 5.** *If  $L_p(E, K)$  is nonzero, then the Mordell-Weil group  $E(K_\infty)$  is finitely generated.*

*Remark.* The nonvanishing of  $L_p(E, K)$  has been established by Vatsal. See for example Theorem 1.1 of [Va2] which even gives a precise formula for the associated  $\mu$ -invariant.

*Assumptions.* Let  $E_p$  be the mod  $p$  representation of  $G_{\mathbb{Q}}$  attached to  $E$ . For simplicity, it is assumed throughout the paper that  $(E, K, p)$  satisfies the following conditions.

**ASSUMPTION 6.** (1) *The prime  $p$  is  $\geq 5$ .*

(2) *The Galois representation attached to  $E_p$  has image isomorphic to  $\mathbf{GL}_2(\mathbb{F}_p)$ .*

(3) *The prime  $p$  does not divide the minimal degree of a modular parametrisation  $X_0(N_0) \rightarrow E$ .*

(4) *For all primes  $\ell$  such that  $\ell^2$  divides  $N$ , and  $p$  divides  $\ell + 1$ , the module  $E_p$  is an irreducible  $I_\ell$ -module.*

*Remarks.* 1. Note that these assumptions are satisfied by all but finitely many primes once  $E$  is fixed, provided that  $E$  has no complex multiplications. They are imposed to simplify the argument and could probably be relaxed.

This is unlike the condition in Theorem 1 which – although it may appear less natural to the uninitiated – is an essential feature of the situation being studied. Indeed, for square-free  $N^-$ , the restriction on the parity of the number of primes appearing in its factorisation is equivalent to requiring that the sign in the functional equation of  $L(E, K, \chi, s)$ , for  $\chi$  a ramified character of  $G_\infty$ , be equal to 1. Without this condition, the  $p$ -adic  $L$ -function  $L_p(E, K, s)$  would vanish identically. See [BD1] for a discussion of this case where it becomes necessary to interpolate the first derivatives  $L'(E, K, \chi, 1)$ .

2. The analogue of Theorem 1 for the cyclotomic  $\mathbb{Z}_p$ -extension has been proved by Kato. Both the proof of Theorem 1 and Kato's proof of the cyclotomic counterpart are based on Kolyvagin's theory of Euler systems.

3. The original ‘‘Euler system’’ argument of Kolyvagin relies on the presence of a systematic supply of algebraic points on  $E$  — the so-called *Heegner points* defined over  $K$  and over abelian extensions of  $K$ . As can be seen from Corollaries 4 and 5, the situation in which we have placed ourselves precludes the existence of a nontrivial norm-compatible system of points in  $E(K_\infty)$ . One circumvents this difficulty by resorting to the theory of congruences between modular forms and the Cerednik-Drinfeld interchange of invariants, which, for each  $n \geq 1$ , realises the Galois representation  $E_{p^n}$  in the  $p^n$ -torsion of the Jacobian of certain Shimura curves for which the Heegner point construction becomes available. By varying the Shimura curves, we produce a compatible collection of cohomology classes in  $H^1(K_\infty, E_{p^n})$ , a collection which can be related to special values of  $L$ -functions and is sufficient to control the Selmer group  $\text{Sel}(K_\infty, E_{p^\infty})$ . It should be noted that this geometric approach to the theory of Euler systems produces ramified cohomology classes in  $H^1(K_\infty, E_{p^n})$  directly without resorting to classes defined over auxiliary ring class field extensions of  $K_\infty$ ; in particular, Kolyvagin's derivative operators make no appearance in the argument. In the terminology of [MR], the strategy of this article produces a ‘‘Kolyvagin system’’ without passing through an Euler system in the sense of [Ru]. This lends some support for the suggestion made in [MR] that Kolyvagin systems are the more fundamental objects of study.

*Acknowledgements.* It is a pleasure to thank Professor Ihara for some useful information on his work, as well as Kevin Buzzard, Ben Howard and the anonymous referees for many helpful comments which led to some corrections and significant improvements in the exposition.

## 1. $p$ -adic $L$ -functions

1.1. *Modular forms on quaternion algebras.* Let  $N^-$  be an arbitrary square-free integer which is the product of an odd number of primes, and let  $N^+$  be any integer prime to  $N^-$ . Let  $p$  be a prime which does not divide

$N^+N^-$  and write  $N = pN^+N^-$ . Let  $B$  be the definite quaternion algebra ramified at all the primes dividing  $N^-$ , and let  $R$  be an Eichler  $\mathbb{Z}[1/p]$ -order of level  $N^+$  in  $B$ . The algebra  $B$  is unique up to isomorphism, and the Eichler order  $R$  is unique up to conjugation by  $B^\times$ , by strong approximation (cf. [Vi, Ch. III, §4 and §5]).

Denote by  $\mathcal{T}$  the Bruhat-Tits tree of  $B_p^\times/\mathbb{Q}_p^\times$ , where

$$B_p := B \otimes \mathbb{Q}_p \simeq M_2(\mathbb{Q}_p).$$

The set  $\mathcal{V}(\mathcal{T})$  of vertices of  $\mathcal{T}$  is indexed by the maximal  $\mathbb{Z}_p$ -orders in  $B_p$ , two vertices being adjacent if their intersection is an Eichler order of level  $p$ . Let  $\vec{\mathcal{E}}(\mathcal{T})$  denote the set of ordered edges of  $\mathcal{T}$ , i.e., the set of ordered pairs  $(s, t)$  of adjacent vertices of  $\mathcal{T}$ . If  $e = (s, t)$ , the vertex  $s$  is called the *source* of  $e$  and the vertex  $t$  is called its *target*; they are denoted by  $s(e)$  and  $t(e)$  respectively.

The tree  $\mathcal{T}$  is endowed with a natural left action of  $B_p^\times/\mathbb{Q}_p^\times$  by isometries corresponding to conjugation of maximal orders by elements of  $B_p^\times$ . This action is transitive on both  $\mathcal{V}(\mathcal{T})$  and  $\vec{\mathcal{E}}(\mathcal{T})$ . Let  $R^\times$  denote the group of invertible elements of  $R$ . The group  $\Gamma := R^\times/\mathbb{Z}[1/p]^\times$  – a discrete subgroup of  $B_p^\times/\mathbb{Q}_p^\times$  in the  $p$ -adic topology – acts naturally on  $\mathcal{T}$  and the quotient  $\mathcal{T}/\Gamma$  is a finite graph.

*Definition 1.1.* A modular form (of weight two) on  $\mathcal{T}/\Gamma$  is a  $\mathbb{Z}_p$ -valued function  $f$  on  $\vec{\mathcal{E}}(\mathcal{T})$  satisfying

$$f(\gamma e) = f(e), \quad \text{for all } \gamma \in \Gamma.$$

Denote by  $S_2(\mathcal{T}/\Gamma)$  the space of such modular forms. It is a free  $\mathbb{Z}_p$ -module of finite rank. More generally, if  $Z$  is any ring, denote by  $S_2(\mathcal{T}/\Gamma, Z)$  the space of  $\Gamma$ -invariant functions on  $\vec{\mathcal{E}}(\mathcal{T})$  with values in  $Z$ .

*Duality.* Let  $e_1, \dots, e_s$  be a set of representatives for the orbits of  $\Gamma$  acting on  $\vec{\mathcal{E}}(\mathcal{T})$ , and let  $w_j$  be the cardinality of the finite group  $\text{Stab}_\Gamma(e_j)$ . The space  $S_2(\mathcal{T}/\Gamma)$  is endowed with a  $\mathbb{Z}_p$ -bilinear pairing defined by

$$(2) \quad \langle f_1, f_2 \rangle = \sum_{i=1}^s w_i f_1(e_i) f_2(e_i).$$

This pairing is nondegenerate so that it identifies  $S_2(\mathcal{T}/\Gamma) \otimes \mathbb{Q}_p$  with its  $\mathbb{Q}_p$ -dual.

*Hecke operators.* Let  $\ell \neq p$  be a prime which does not divide  $p$ . Choose an element  $M_\ell$  of reduced norm  $\ell$  in the  $\mathbb{Z}[1/p]$ -order  $R$  that was used to define  $\Gamma$ . The double coset  $\Gamma M_\ell \Gamma$  decomposes as a disjoint union of left cosets:

$$(3) \quad \Gamma M_\ell \Gamma = \gamma_1 \Gamma \cup \dots \cup \gamma_t \Gamma.$$

Here  $t = \ell + 1$  (resp.  $\ell, 1$ ) if  $\ell$  does not divide  $N^+N^-$  (resp. divides  $N^+, N^-$ ). The function  $f_{|\ell}$  defined on  $\vec{\mathcal{E}}(\mathcal{T})$  by the rule

$$(4) \quad f_{|\ell}(e) = \sum_{i=1}^t f(\gamma_i^{-1}e)$$

is independent of the choice of  $M_\ell$  or of the representatives  $\gamma_1, \dots, \gamma_t$ , and the assignment  $f \mapsto f_{|\ell}$  is a linear endomorphism of  $S_2(\mathcal{T}/\Gamma)$ , called the  $\ell^{\text{th}}$  Hecke operator at  $\ell$  and denoted  $T_\ell$  if  $\ell$  does not divide  $N$ , and  $U_\ell$  if  $\ell$  divides  $N^+N^-$ .

Associated to the prime  $p$  there is a Hecke operator denoted  $U_p$  and defined by the rule

$$(5) \quad (U_p f)(e) = \sum_{s(e')=t(e)} f(e'),$$

where the sum is taken over the  $p$  edges  $e'$  with source equal to the target of  $e$ , not including the edge obtained from  $e$  by reversing the orientation. The Hecke operators  $T_\ell$  (with  $\ell \nmid N$ ) are called the *good* Hecke operators. They are self-adjoint for the pairing on  $S_2(\mathcal{T}/\Gamma)$  defined in (2):

$$(6) \quad \langle T_\ell f_1, f_2 \rangle = \langle f_1, T_\ell f_2 \rangle.$$

*Oldforms and Newforms.* Let  $S_2(\mathcal{V}/\Gamma, Z)$  denote the space of  $\Gamma$ -invariant  $Z$ -valued functions on  $\mathcal{V}(\mathcal{T})$ , equipped with a  $Z$ -valued bilinear pairing as in (2) with edges replaced by vertices. There are two natural “degeneracy maps”  $s^*, t^* : S_2(\mathcal{V}/\Gamma) \rightarrow S_2(\mathcal{T}/\Gamma)$  defined by

$$s^*(f)(e) = f(s(e)), \quad t^*(f)(e) = f(t(e)).$$

A form  $f \in S_2(\mathcal{T}/\Gamma, Z)$  is said to be  $p$ -old if there exist  $\Gamma$ -invariant functions  $f_1$  and  $f_2$  on  $\mathcal{V}(\mathcal{T})$  such that

$$(7) \quad f = s^*(f_1) + t^*(f_2).$$

A form which is orthogonal to the oldforms (i.e., is orthogonal to the image of  $s^*$  and  $t^*$ ) is said to be  $p$ -new. The form  $f$  is  $p$ -new if and only if  $f$  is *harmonic* in the sense that it satisfies

$$(8) \quad s_*(f)(v) := \sum_{s(e)=v} f(e) = 0, \quad t_*(f)(v) := \sum_{t(e)=v} f(e) = 0, \quad \forall v \in \mathcal{V}(\mathcal{T}).$$

This can be seen by noting that  $s_*$  and  $t_*$  are the adjoints of the maps  $s^*$  and  $t^*$  respectively.

*$p$ -isolated forms.* Let  $\mathbb{T}$  be the Hecke algebra acting on the space  $S_2(\mathcal{T}/\Gamma)$ . A form  $f$  in this space is called an *eigenform* if it is a simultaneous eigenvector for all the Hecke operators, i.e.,

$$\begin{aligned} T_\ell(f) &= a_\ell(f)f, & \text{for all } \ell \nmid N, \\ U_\ell(f) &= \alpha_\ell(f)f, & \text{for all } \ell \mid N, \end{aligned}$$

where the eigenvalues  $a_\ell(f)$  and  $\alpha_\ell(f)$  belong to  $\mathbb{Z}_p$ . Such an eigenform determines a maximal ideal  $\mathfrak{m}_f$  of  $\mathbb{T}$  by the rule

$$\mathfrak{m}_f := \langle p, T_\ell - a_\ell(f), U_\ell - \alpha_\ell(f) \rangle.$$

*Definition 1.2.* The eigenform  $f$  is said to be  $p$ -isolated if the completion of  $S_2(\mathcal{T}/\Gamma)$  at  $\mathfrak{m}_f$  is a free  $\mathbb{Z}_p$ -module of rank one.

In other words,  $f$  is  $p$ -isolated if there are no nontrivial congruences between  $f$  and other modular forms in  $S_2(\mathcal{T}/\Gamma)$ . Note that this is really a property of the mod  $p$  eigenform in  $S_2(\mathcal{T}/\Gamma, \mathbb{F}_p)$  associated to  $f$ , or of the maximal ideal  $\mathfrak{m}_f$ , so that it makes sense to say that  $\mathfrak{m}_f$  is  $p$ -isolated if it is attached to (the reduction of) a  $p$ -isolated eigenform.

*The Jacquet-Langlands correspondence.* The complex vector space  $S_2(\mathcal{H}/\Gamma_0(N))$  of classical modular forms of weight 2 on  $\mathcal{H}/\Gamma_0(N)$  is similarly endowed with an action of Hecke operators, which will also be denoted by the symbols  $T_\ell$ ,  $U_\ell$  and  $U_p$  by abuse of notation. Let  $\phi$  be an eigenform on  $\Gamma_0(N)$  which arises from a newform  $\phi_0$  of level  $N_0$ . It is a simultaneous eigenfunction for all the good Hecke operators  $T_\ell$ . Assume that it is also an eigenfunction for the Hecke operator  $U_p$ . Write  $a_\ell$  for the eigenvalue of  $T_\ell$  acting on  $\phi$ , and  $\alpha_p$  for the eigenvalue of  $U_p$  acting on  $\phi$ .

*Remark.* If  $p$  does not divide  $N_0$ , so that  $\phi$  is not new at  $p$ , then the eigenvalue  $\alpha_p$  is a root of the polynomial  $x^2 - a_p x + p$ , where  $a_p$  is the eigenvalue of  $T_p$  acting on  $\phi_0$ . If  $p$  divides  $N_0$ , then  $\phi = \phi_0$  and the eigenvalue  $\alpha_p$  is equal to 1 (resp.  $-1$ ) if the abelian variety attached to  $\phi$  by the Eichler-Shimura construction has split (resp. nonsplit) multiplicative reduction at  $p$ .

**PROPOSITION 1.3.** *Let  $\phi$  be as above. Then there exists an eigenform  $f$  in  $S_2(\mathcal{T}/\Gamma, \mathbb{C})$  satisfying*

$$(9) \quad \begin{aligned} T_\ell f &= a_\ell(\phi) f \text{ for all } \ell \nmid N, \\ U_\ell f &= \alpha_\ell(\phi) f \text{ for all } \ell \mid N^+, \quad U_p f = \alpha_p(\phi) f. \end{aligned}$$

*The form  $f$  with these properties is unique up to multiplication by a nonzero complex number. Conversely, given an eigenform  $f \in S_2(\mathcal{T}/\Gamma, \mathbb{C})$ , there exists an eigenform  $\phi \in S_2(\mathcal{H}/\Gamma_0(N))$  satisfying (9).*

*Proof.* Suppose first that  $p$  divides  $N_0$ , so that  $\phi$  is a newform on  $\Gamma_0(N)$ . Let  $R_0$  be an Eichler  $\mathbb{Z}$ -order of level  $pN^+$  in the definite quaternion algebra of discriminant  $N^-$ . Write  $\hat{R}_0 = R_0 \otimes \hat{\mathbb{Z}} = \prod_\ell R_0 \otimes \mathbb{Z}_\ell$ , and  $\hat{B} := \hat{R}_0 \otimes \mathbb{Q}$ . The Jacquet-Langlands correspondence (which, in this case, can be established by use of the Eichler trace formula as in [Ei]; see also [JL] and the discussion in

Chapter 5 of [DT]) implies the existence of a unique function

$$(10) \quad f : B^\times \backslash \hat{B}^\times / \hat{R}_0^\times \longrightarrow \mathbb{C}$$

satisfying  $T_\ell f = a_\ell f$  for all  $\ell \nmid N$ , and  $U_p f = \alpha_p f$  (where the operators  $T_\ell$  and  $U_p$  are the general Hecke operators defined in terms of double cosets as in [Sh]). Strong approximation identifies the double coset space appearing in (10) with the space  $R^\times \backslash B_p^\times / (R_0)_p^\times$ . The transitive action of  $B_p^\times$  on the set of maximal orders in  $B_p$  by conjugation yields an action of  $B_p^\times$  on  $\mathcal{T}$  by isometries, for which the subgroup  $(R_0)_p^\times$  is equal to the stabiliser of a certain oriented edge. In this way  $B_p^\times / (R_0)_p^\times$  is identified with  $\vec{\mathcal{E}}(\mathcal{T})$ , and  $f$  can thus be viewed as an element of  $S_2(\mathcal{T}/\Gamma, \mathbb{C})$ .

If  $p$  does not divide  $N_0$ , let  $a_p$  denote the eigenvalue of  $T_p$  acting on  $\phi_0$ , and let  $R_0$  denote now the Eichler order of level  $N^+$  in the quaternion algebra  $B$ . As before, to the form  $\phi_0$  is associated a unique function

$$(11) \quad f_0 : B^\times \backslash \hat{B}^\times / \hat{R}_0^\times \longrightarrow \mathbb{C}$$

satisfying  $T_\ell f = a_\ell f$  for all  $\ell \nmid N_0$ . As before, strong approximation makes it possible to identify  $f_0$  with a  $\Gamma$ -invariant function on  $\mathcal{V}(\mathcal{T})$ . In this description, the action of  $T_p$  on  $f_0$  is given by the formula

$$T_p(f_0(v)) = \sum_w f_0(w),$$

where the sum is taken over the  $p+1$  vertices  $w$  of  $\mathcal{T}$  which are adjacent to  $v$ . Define functions  $f_s, f_t : \vec{\mathcal{E}}(\mathcal{T}) \longrightarrow \mathbb{C}$  by the rules:

$$f_s(e) = f_0(s(e)), \quad f_t(e) = f_0(t(e)).$$

The forms  $f_s$  and  $f_t$  both satisfy  $T_\ell(g) = a_\ell g$  for all  $\ell \nmid N$ , and span the two-dimensional eigenspace of forms with this property. A direct calculation reveals that

$$U_p f_s = p f_t, \quad U_p f_t = -f_s + a_p f_t.$$

The function  $f = f_s - \alpha_p f_t$  satisfies  $U_p f = \alpha_p f$ , and is, up to scaling, the unique eigenform in  $S_2(\mathcal{T}/\Gamma, \mathbb{C})$  with this property.

The converse is proved by essentially reversing the argument above: to an eigenform  $f \in S_2(\mathcal{T}/\Gamma, \mathbb{C})$  is associated a function on the adelic coset space attached to  $B^\times$  as in (10); the Jacquet-Langlands correspondence (applied now in the reverse direction) produces the desired  $\phi \in S_2(\mathcal{H}/\Gamma_0(N))$ .

*The Shimura-Taniyama conjecture.* Let  $E$  be an elliptic curve as in the introduction. For each prime  $\ell$  which does not divide  $N$ , set

$$a_\ell = \ell + 1 - \#E(\mathbb{F}_\ell).$$

If  $E$  has good ordinary reduction at  $p$ , let  $\alpha_p \in \mathbb{Z}_p$  be the unique root of the polynomial  $x^2 - a_p x + p$  which is a  $p$ -adic unit. Set  $\alpha_p = 1$  (resp.  $-1$ ) if  $E$  has

split (resp. nonsplit) multiplicative reduction at  $p$ . The following theorem is a consequence of the Shimura-Taniyama conjecture in view of Proposition 1.3.

PROPOSITION 1.4. *There exists an eigenform  $f$  in  $S_2(\mathcal{T}/\Gamma)$  satisfying*

$$T_\ell f = a_\ell f, \text{ for all } \ell \nmid N, \quad U_p f = \alpha_p f,$$

$$f \notin pS_2(\mathcal{T}/\Gamma).$$

*The form  $f$  with these properties is unique up to multiplication by a scalar in  $\mathbb{Z}_p^\times$ .*

*Proof.* Proposition 1.3 shows that there exists a form  $f \in S_2(\mathcal{T}/\Gamma, \mathbb{C})$  satisfying the conclusion of Proposition 1.4. The eigenvalues  $a_\ell$  belong to  $\mathbb{Z}$ , and, since  $E$  is ordinary at  $p$ , the eigenvalue  $\alpha_p$  belongs to the ring of integers  $\mathcal{O}$  of a quadratic extension of  $\mathbb{Q}$  in which  $p$  splits completely. Hence, the form  $f$  can be chosen to lie in  $S_2(\mathcal{T}/\Gamma, \mathcal{O})$ . After applying the unique embedding of  $\mathcal{O}$  into  $\mathbb{Z}_p$  which sends  $\alpha_p$  to a  $p$ -adic unit, and rescaling  $f$  appropriately, one obtains a form in  $S_2(\mathcal{T}/\Gamma)$  satisfying the conclusion of Proposition 1.4.

1.2.  *$p$ -adic Rankin  $L$ -functions.* An eigenform  $f$  in  $S_2(\mathcal{T}/\Gamma)$  is said to be *ordinary* if the eigenvalue  $\alpha_p$  of  $U_p$  acting on  $f$  is a  $p$ -adic unit. This section recalls the definition of the  $p$ -adic Rankin  $L$ -function attached to an ordinary form on  $\mathcal{T}/\Gamma$  and a quadratic algebra  $K \subset B$ .

If  $A$  is any  $\mathbb{Z}$ -algebra, let

$$(12) \quad A_\ell = A \otimes \mathbb{Z}_\ell, \quad \hat{A} = A \otimes \hat{\mathbb{Z}} \subset \prod_\ell A_\ell.$$

Let  $K$  be a quadratic algebra of discriminant prime to  $N$  which embeds in  $B$ . Since  $B$  is definite of discriminant  $N^-$ , the algebra  $K$  is an imaginary quadratic field in which all prime divisors of  $N^-$  are inert. Let  $\mathcal{O}_K$  denote the ring of integers of  $K$  and let  $\mathcal{O} = \mathcal{O}_K[1/p]$  be the maximal  $\mathbb{Z}[1/p]$ -order in  $K$ .

Let  $\tilde{G}_\infty$  denote the group

$$(13) \quad \tilde{G}_\infty = \hat{K}^\times / (\hat{\mathbb{Q}}^\times \prod_{\ell \neq p} \mathcal{O}_\ell^\times K^\times).$$

Fix an embedding

$$(14) \quad \Psi : K \longrightarrow B \quad \text{satisfying} \quad \Psi(K) \cap R = \Psi(\mathcal{O}).$$

Such a  $\Psi$  exists if and only if all the primes dividing  $N^+$  are split in  $K$ . By passing to the adélisation the embedding  $\Psi$  induces a map

$$(15) \quad \hat{\Psi} : \tilde{G}_\infty \longrightarrow B^\times \backslash \hat{B}^\times / \left( \hat{\mathbb{Q}}^\times \prod_{\ell \neq p} R_\ell^\times \right).$$

By strong approximation ([Vi, Ch. III, §4]), the double coset space appearing on the right has a fundamental region contained in  $B_p^\times \subset \hat{B}^\times$ . In fact, strong approximation yields a canonical identification

$$(16) \quad \eta : B^\times \backslash \hat{B}^\times / \left( \hat{\mathbb{Q}}^\times \prod_{\ell \neq p} R_\ell^\times \right) \longrightarrow \Gamma \backslash B_p^\times / \mathbb{Q}_p^\times.$$

The modular form  $f \in S_2(\mathcal{T}/\Gamma)$  determines a pairing between  $\tilde{G}_\infty$  and  $\vec{\mathcal{E}}(\mathcal{T})$  by the rule

$$(17) \quad [\sigma, e]_f := f(\eta \hat{\Psi}(\sigma)e) \in \mathbb{Z}_p.$$

The embedding  $\Psi$  induces an embedding of  $K_p^\times$  into  $B_p^\times$  and hence yields an action of  $K_p^\times/\mathbb{Q}_p^\times$  on  $\mathcal{T}$ . This action fixes a single vertex if  $p$  is inert in  $K$ , and no vertex if  $p$  is split in  $K$ . Let

$$(18) \quad U_n := (1 + p^n \mathcal{O}_K \otimes \mathbb{Z}_p)^\times / (1 + p^n \mathbb{Z}_p)^\times$$

denote the standard compact subgroup of  $K_p^\times/\mathbb{Q}_p^\times$  of level  $n$ . Choose a sequence  $e_1, e_2, \dots, e_n, \dots$  of consecutive edges on  $\mathcal{T}$  satisfying

$$(19) \quad \text{Stab}_{K_p^\times/\mathbb{Q}_p^\times}(e_j) = U_j, \quad j = 1, \dots, n, \dots$$

Since  $\alpha_p$  is a  $p$ -adic unit, the pairing defined by equation (17) can be used to define a  $\mathbb{Z}_p$ -valued distribution  $\tilde{\nu}_f$  on  $\tilde{G}_\infty$  by the rule

$$(20) \quad \tilde{\nu}_f(\sigma U_j) := \alpha_p^{-j} [\sigma, e_j]_f,$$

for all compact open subsets of  $\tilde{G}_\infty$  of the form  $\sigma U_j$  with  $\sigma \in \tilde{G}_\infty$ . The distribution relation for  $\tilde{\nu}_f$  is ensured by the fact that  $f$  is an eigenform for the  $U_p$  operator with eigenvalue  $\alpha_p$ . The distribution  $\tilde{\nu}_f$  gives rise to an element  $\tilde{\mathcal{L}}_f$  in the completed integral group ring  $\mathbb{Z}_p[[\tilde{G}_\infty]]$  by the rule

$$(\tilde{\mathcal{L}}_f)_n := \sum_{g \in \tilde{G}_n} \tilde{\nu}_f(g U_n) \cdot g,$$

where  $\tilde{G}_n := \tilde{G}_\infty / U_n$  so that  $\tilde{G}_\infty$  is the inverse limit of the finite groups  $\tilde{G}_n$ .

Let  $\Delta$  denote the torsion subgroup of  $\tilde{G}_\infty$ , and let

$$(21) \quad G_\infty = \tilde{G}_\infty / \Delta \simeq \mathbb{Z}_p.$$

Write  $\mathcal{L}_f$  for the natural image of  $\tilde{\mathcal{L}}_f$  in the Iwasawa algebra

$$\Lambda = \mathbb{Z}_p[[G_\infty]] \simeq \mathbb{Z}_p[[T]],$$

and denote by  $\nu_f$  the associated measure on  $G_\infty$ . Note that a different choice of edges  $e_j$  satisfying (19) has the effect of multiplying  $\mathcal{L}_f$  by an element of  $G_\infty$ , so that  $\mathcal{L}_f$  is only well-defined up to multiplication by such elements.

*Functional equations.* The Iwasawa algebra is equipped with the involution  $\theta \mapsto \theta^*$  sending any  $\sigma \in G_\infty$  to  $\sigma^{-1}$ . Let  $\varepsilon = \pm 1$  be the sign in the functional equation of the classical  $L$ -function  $L(E/\mathbb{Q}, s)$  attached to  $E/\mathbb{Q}$ . Conjecturally, the value of  $\varepsilon$  determines the parity of the rank of  $E/\mathbb{Q}$ . More precisely, this rank should be even if  $\varepsilon = 1$ , and odd if  $\varepsilon = -1$ . Set  $\varepsilon_p = \varepsilon$  if  $E$  does not have split multiplicative reduction over  $\mathbb{Q}_p$ , and set  $\varepsilon_p = -\varepsilon$  otherwise. The sign  $\varepsilon_p$  is interpreted in [MTT] as the sign in the functional equation for the Mazur-Swinnerton-Dyer  $p$ -adic  $L$ -function attached to  $E/\mathbb{Q}$ . While this  $L$ -function differs markedly from the  $p$ -adic Rankin  $L$ -function considered in this article, one still has

LEMMA 1.5. *The equality*

$$\mathcal{L}_f^* = \varepsilon_p \mathcal{L}_f$$

holds in  $\Lambda$ , up to multiplication by an element of  $G_\infty$ .

*Proof.* See Proposition 2.13 and equation (11) of [BD1].

*Definition 1.6.* The *anticyclotomic Rankin  $L$ -function* attached to  $f$  and  $K$  is the element  $L_p(f, K)$  of  $\Lambda$  defined by

$$L_p(f, K) = \mathcal{L}_f \mathcal{L}_f^*.$$

*Remarks.* 1. Note that  $L_p(f, K)$  is a well-defined element of  $\Lambda$ , since multiplying  $\mathcal{L}_f$  by  $\sigma \in G_\infty$  has the effect of multiplying  $\mathcal{L}_f^*$  by  $\sigma^{-1}$ . Thus the ambiguity in the definition of  $\mathcal{L}_f$  arising from the choice of end in  $\mathcal{T}$  satisfying (19) is cancelled out.

2. Definition 1.6 extends naturally, *mutatis mutandis*, to any eigenform  $g$  in  $S_2(\mathcal{T}/\Gamma, Z)$ , where  $Z$  is any ring in which the eigenvalue of  $U_p$  acting on  $g$  is invertible. In this case the anticyclotomic Rankin  $L$ -function  $L_p(g, K)$  is simply an element of the completed group ring  $Z[[G_\infty]]$ .

Let  $\mu_{f,K}$  be the  $\mathbb{Z}_p$ -valued measure on  $G_\infty$  associated to  $L_p(f, K)$ . The function  $L_p(f, K, s)$  is defined to be the  $p$ -adic Mellin transform of  $\mu_{f,K}$ :

$$L_p(f, K, s) := \int_{G_\infty} g^{s-1} d\mu_{f,K}(g)$$

where  $g^{s-1} := \exp((s-1)\log(g))$ , and  $\log : G_\infty \rightarrow \mathbb{Q}_p$  is a choice of  $p$ -adic logarithm.

*Interpolation properties.* Let  $\phi$  be the normalised eigenform on  $\Gamma_0(N)$  attached to  $f$  via the Jacquet-Langlands correspondence of Proposition 1.3, and let  $\Omega_f = \langle \phi, \phi \rangle$  denote the Peterson scalar product of  $\phi$  with itself. It is known (cf. [Zh, §1.4]) that the measure  $\mu_{f,K}$  on  $G_\infty$  satisfies the following

$p$ -adic interpolation property:

$$\left| \int_{\tilde{G}_\infty} \chi(g) d\mu_{f,K}(g) \right|^2 \doteq L(f, K, \chi, 1) / (\sqrt{\text{Disc}(K)} \Omega_f),$$

for all ramified finite order characters  $\chi$  of  $\tilde{G}_\infty$ . Here the values of  $\chi$  and  $\mu_{f,K}$  are viewed as complex numbers by fixing an embedding of  $\bar{\mathbb{Q}}_p$  in  $\mathbb{C}$ , and the absolute value taken on the left-hand side is the complex one. The symbol  $\doteq$  indicates an equality up to a simple algebraic fudge factor expressed as a product of terms comparatively less important than the quantities explicitly described in the formulas. Note in particular that dividing  $L(f, K, \chi, 1)$  by the complex period  $\Omega_f$  yields an algebraic number.

*Elliptic curves.* If  $E$  is an elliptic curve as in the introduction, let  $f_E$  be the modular form in  $S_2(\mathcal{T}/\Gamma)$  attached to it by Proposition 1.4. The  $p$ -adic  $L$ -function attached to  $E$  and  $K$  is defined by:

$$(22) \quad L_p(E, K) := L_p(f_E, K), \quad L_p(E, K, s) := L_p(f_E, K, s).$$

*Remark.* Note that  $L_p(E, K)$  is only well-defined up to multiplication by a unit in  $\mathbb{Z}_p^\times$ , since the same is true of the form  $f_E$  attached to it by Proposition 1.4.

## 2. Selmer groups

2.1. *Galois representations and cohomology.* Let  $f$  be an ordinary eigenform in  $S_2(\mathcal{T}/\Gamma)$  with coefficients in  $\mathbb{Z}_p$ , and let  $K$  be a quadratic imaginary field in which all primes dividing  $N^-$  (resp.  $N^+$ ) are inert (resp. split). To these two objects a  $p$ -adic  $L$ -function  $L_p(f, K)$  was attached in Section 1. This section introduces an invariant of a more arithmetic nature — the so-called *Selmer group* attached to  $f$  and  $K$ .

*Galois representations.* To  $f$  is attached a continuous representation of the Galois group  $G_{\mathbb{Q}}$ :

$$V_f \simeq \mathbb{Q}_p^2,$$

with determinant the  $p$ -adic cyclotomic character and satisfying

$$(23) \quad \text{trace}((\text{Frob}_\ell)|_{V_f}) = a_\ell(f), \quad \text{for all } \ell \nmid N.$$

This representation is constructed by invoking Proposition 1.3 to associate to  $f$  a classical eigenform  $\phi \in S_2(\mathcal{H}/\Gamma_0(N))$  with the same Hecke eigenvalues as  $f$  at the good primes. The representation  $V_f$  arises in the Jacobian of  $J_0(N)$  by the well-known construction of Eichler and Shimura ([DDT, §3.1]).

The action of the compact group  $G_{\mathbb{Q}}$  is continuous for the  $p$ -adic topology on  $V_f$  and hence preserves a  $\mathbb{Z}_p$ -lattice  $T_f$ . Let

$$(24) \quad A_f = V_f/T_f \simeq (\mathbb{Q}_p/\mathbb{Z}_p)^2$$

be the divisible  $G_{\mathbb{Q}}$ -module attached to  $f$ , and let  $A_{f,n} := A_f[p^n]$  denote the  $p^n$ -torsion submodule of  $A_f$ . It will be convenient occasionally to denote  $A_f$  by  $A_{f,\infty}$ . Likewise write  $T_{f,n} = T_f/p^n T_f$  and set  $T_{f,\infty} := T_f$ . Note that for  $n < \infty$ , the modules  $A_{f,n}$  and  $T_{f,n}$  are isomorphic as  $G_{\mathbb{Q}}$ -modules, but the  $A_{f,n}$  fit naturally into an inductive system while the  $T_{f,n}$  are part of a projective system. It is therefore useful to maintain the notational distinction between the two.

The fact that  $f$  is ordinary at  $p$  implies that  $A_{f,n}$  is ordinary, in the sense that it has a quotient  $A_{f,n}^{(1)}$  which is free of rank one over  $\mathbb{Z}/p^n\mathbb{Z}$  and on which the inertia group  $I_p$  at  $p$  acts trivially. The kernel of the natural projection  $A_{f,n} \rightarrow A_{f,n}^{(1)}$  is a free module of rank one over  $\mathbb{Z}/p^n\mathbb{Z}$ , denoted  $A_{f,n}^{(p)}$ , on which  $I_p$  acts via the the  $p$ -adic cyclotomic character

$$\varepsilon : G_{\mathbb{Q}} \rightarrow \mathbb{Z}_p^{\times}$$

describing the action of  $G_{\mathbb{Q}}$  on the  $p$ -power roots of unity.

In our treatment of the Selmer group attached to  $f$  and  $K$ , it is convenient to make the following technical assumption on  $f$ :

**ASSUMPTION 2.1.** *The Galois representation  $A_{f,1}$  is surjective. Furthermore, for all  $\ell$  dividing  $N_0$  exactly, the Galois representation  $A_{f,1}$  has a unique one-dimensional subspace  $A_{f,1}^{(\ell)}$  on which  $\text{Gal}(\bar{\mathbb{Q}}_{\ell}/\mathbb{Q}_{\ell})$  acts via  $\varepsilon$  or  $-\varepsilon$ .*

*Remarks.* 1. Note that Assumption 2.1 is automatically satisfied for  $\ell$  if  $A_{f,1}$  is ramified at  $\ell$ , because  $A_{f,n}$  arises from the Tate module of an abelian variety which acquires purely toric reduction over the quadratic unramified extension of  $\mathbb{Q}_{\ell}$ . If  $A_{f,1}$  is unramified at  $\ell$ , then the Frobenius element at  $\ell$  acts on  $A_{f,1}$  with eigenvalues  $\pm 1$  and  $\pm \ell$ , and the condition in Assumption 2.1 stipulates that  $p$  should not divide  $\ell^2 - 1$ .

2. For the same reason as explained in Remark 1, the maximal submodule  $A_{f,n}^{(\ell)}$  on which  $G_{\mathbb{Q}_{\ell}}$  acts via  $\pm \varepsilon$  is free of rank one over  $\mathbb{Z}/p^n\mathbb{Z}$ .

**LEMMA 2.2.** *Suppose that  $E$  satisfies Assumption 6 of the introduction. Then Assumption 2.1 is satisfied by the modular form  $f$  attached to  $E$ .*

*Proof.* Note that in this case  $T_f$  is simply isomorphic to the Tate module of  $E$ . The assumption that  $p$  does not divide the degree of the modular parametrisation of  $E$  implies that the newform on  $\Gamma_0(N_0)$  attached to  $E$  is  $p$ -isolated. By Ribet's level-lowering theorem [Ri2], it follows that the Galois

representation attached to  $A_{f,1}$  is ramified at all primes dividing  $N_0$ , and hence Lemma 2.2 follows from Remark 1 after the statement of Assumption 2.1.

*$\mathbb{Z}_p$ -extensions.* Class field theory identifies the group  $\tilde{G}_\infty$  of (13) with the Galois group of the maximal abelian extension  $\tilde{K}_\infty$  of  $K$  which is unramified outside of  $p$  and which is of “dihedral type” over  $\mathbb{Q}$ . The subfield  $K_\infty := \tilde{K}_\infty^\Delta$  is called the *anticyclotomic  $\mathbb{Z}_p$ -extension* of  $K$ . Its Galois group over  $K$  is identified with the group  $G_\infty \simeq \mathbb{Z}_p$  of equation (21). Let  $K_m$  be the  $m^{\text{th}}$  layer of  $K_\infty/K$ , so that  $\text{Gal}(K_m/K) \simeq \mathbb{Z}/p^m\mathbb{Z}$ .

*Galois cohomology.* For each  $m \in \mathbb{N}$  and  $n \in \mathbb{N} \cup \{\infty\}$ , let  $H^1(K_m, A_{f,n})$  and  $H^1(K_m, T_{f,n})$  denote the usual continuous Galois cohomology groups of  $\text{Gal}(\tilde{K}_m/K_m)$  with values in these modules. (Note that

$$H^1(K_m, A_f) := \varinjlim_n H^1(K_m, A_{f,n}), \quad H^1(K_m, T_f) := \varinjlim_n H^1(K_m, T_{f,n}).)$$

To study the behaviour of these groups as  $K_m$  varies over the finite layers of the anticyclotomic  $\mathbb{Z}_p$ -extension, it is convenient to introduce the groups

$$H^1(K_\infty, A_{f,n}) := \varinjlim_m H^1(K_m, A_{f,n}), \quad \hat{H}^1(K_\infty, T_{f,n}) = \varinjlim_m H^1(K_m, T_{f,n}),$$

where the direct limit is taken with respect to the natural restriction maps, and the inverse limit is taken with respect to the norm (corestriction) maps. The compatible actions of the group rings  $\mathbb{Z}_p[G_m]$  on the groups  $H^1(K_m, A_{f,n})$  and  $H^1(K_m, T_{f,n})$  yield an action of the Iwasawa algebra  $\Lambda = \mathbb{Z}_p[[G_\infty]]$  on both of the groups  $H^1(K_\infty, A_{f,n})$  and  $\hat{H}^1(K_\infty, T_{f,n})$ .

*Local cohomology groups.* For each rational prime  $\ell$ , set

$$K_{m,\ell} := K_m \otimes \mathbb{Q}_\ell = \bigoplus_{\lambda|\ell} K_{m,\lambda},$$

where the direct sum is taken over all primes  $\lambda$  of  $K_m$  dividing  $\ell$ , and write for any  $G_{K_m}$ -module  $X$ :

$$H^1(K_{m,\ell}, X) := \bigoplus_{\lambda|\ell} H^1(K_{m,\lambda}, X).$$

Set

$$H^1(K_{\infty,\ell}, A_{f,n}) = \varinjlim_m H^1(K_{m,\ell}, A_{f,n}), \quad \hat{H}^1(K_{\infty,\ell}, T_{f,n}) = \varinjlim_m H^1(K_{m,\ell}, T_{f,n})$$

for the local counterparts of  $H^1(K_\infty, A_{f,n})$  and  $\hat{H}^1(K_\infty, T_{f,n})$ . The Iwasawa algebra  $\Lambda$  acts naturally on these modules in a manner which is compatible with the restriction maps. For each rational prime  $\ell$ , write

$$H^1(I_{m,\ell}, A_{f,n}) := \bigoplus_{\lambda|\ell} H^1(I_{m,\lambda}, A_{f,n}),$$

where  $I_{m,\lambda}$  denotes the inertia group at  $\lambda$ .

*Tate duality.* Let  $\ell$  be a rational prime, and let  $n \in \mathbb{N} \cup \{\infty\}$ . The finite Galois modules  $T_{f,n} = A_{f,n}$  are isomorphic to their own Kummer duals: the Weil pairing gives rise to a canonical  $G_{\mathbb{Q}}$ -equivariant pairing

$$T_{f,n} \times A_{f,n} \longrightarrow \mathbb{Z}/p^n\mathbb{Z}(1) = \mu_{p^n}.$$

Combining this with the cup product pairing in cohomology gives rise to the collection of local Tate pairings at the primes above  $\ell$  over the finite layers  $K_m$  in  $K_{\infty}$ :

$$(25) \quad \langle \cdot, \cdot \rangle_{m,\ell} : H^1(K_{m,\ell}, T_{f,n}) \times H^1(K_{m,\ell}, A_{f,n}) \longrightarrow \mathbb{Q}_p/\mathbb{Z}_p,$$

which gives rise, after passing to the limit with  $m$ , to a perfect pairing

$$\langle \cdot, \cdot \rangle_{\ell} : \hat{H}^1(K_{\infty,\ell}, T_{f,n}) \times H^1(K_{\infty,\ell}, A_{f,n}) \longrightarrow \mathbb{Q}_p/\mathbb{Z}_p.$$

These pairings satisfy the rule

$$\langle \lambda \kappa, s \rangle_{\ell} = \langle \kappa, \lambda^* s \rangle_{\ell},$$

for all  $\lambda \in \Lambda$ , and hence give an isomorphism of  $\Lambda$ -modules

$$\hat{H}^1(K_{\infty,\ell}, T_{f,n}) \longrightarrow H^1(K_{\infty,\ell}, A_{f,n})^{\vee},$$

where the Pontrjagin dual  $X^{\vee}$  of a  $\Lambda$ -module  $X$  is itself endowed with a  $\Lambda$ -module structure by the rule

$$\lambda f(x) := f(\lambda^* x), \quad \text{for all } \lambda \in \Lambda, f \in X^{\vee}, x \in X.$$

*2.2. Finite/singular structures.* Let  $\ell \nmid N$  be a rational prime. The *singular part* of  $H^1(K_{m,\ell}, A_{f,n})$  is the group

$$H_{\text{sing}}^1(K_{m,\ell}, A_{f,n}) := H^1(I_{m,\ell}, A_{f,n})^{G_{\kappa\ell}}.$$

There is a natural map arising from restriction — the so-called *residue map* —

$$\partial_{\ell} : H^1(K_{m,\ell}, A_{f,n}) \longrightarrow H_{\text{sing}}^1(K_{m,\ell}, A_{f,n}).$$

Let  $H_{\text{fin}}^1(K_{m,\ell}, A_{f,n})$  denote the kernel of  $\partial_{\ell}$ . The classes in  $H_{\text{fin}}^1(K_{m,\ell}, A_{f,n})$  are sometimes called the *finite* or *unramified classes*.

Of course, identical definitions can be made in which  $A_{f,n}$  is replaced by  $T_{f,n}$ . By passing to the limit as  $m \rightarrow \infty$  (with either a direct or an inverse limit) the definition of the residue map  $\partial_{\ell}$  extends both to  $H^1(K_{\infty,\ell}, A_{f,n})$  and to  $\hat{H}^1(K_{\infty,\ell}, T_{f,n})$  and the groups

$$\begin{aligned} H_{\text{fin}}^1(K_{\infty,\ell}, A_{f,n}), & \quad \hat{H}_{\text{fin}}^1(K_{\infty,\ell}, T_{f,n}), \\ H_{\text{sing}}^1(K_{\infty,\ell}, A_{f,n}), & \quad \hat{H}_{\text{sing}}^1(K_{\infty,\ell}, T_{f,n}) \end{aligned}$$

are defined in the natural way.

Let  $\ell$  be a prime dividing  $N_0$  exactly. Recall in this case (in view of Assumption 2.1) the distinguished line  $A_{f,n}^{(\ell)}$  consisting of elements on which

$G_{\mathbb{Q}_\ell}$  acts via  $\pm\varepsilon$ . The *ordinary part* of  $H^1(K_{\infty,\ell}, A_{f,n})$  is defined to be the group

$$H_{\text{ord}}^1(K_{\infty,\ell}, A_{f,n}) := H^1(K_{\infty,\ell}, A_{f,n}^{(\ell)}).$$

Finally, at the prime  $p$ , set

$$H_{\text{ord}}^1(K_{\infty,p}, A_{f,n}) := \text{res}_p^{-1} \left( H^1(I_{\infty,p}, A_{f,n}^{(p)}) \right),$$

where  $\text{res}_p : H^1(K_{\infty,p}, A_{f,n}) \rightarrow H^1(I_{\infty,p}, A_{f,n})$  is induced from the restriction maps at the (finitely many) primes of  $K_\infty/K$  above  $p$ .

**PROPOSITION 2.3.** *If  $\ell$  is a prime not dividing  $N$ , the groups  $H_{\text{fin}}^1(K_{\infty,\ell}, A_{f,n})$  and  $\hat{H}_{\text{fin}}^1(K_{\infty,\ell}, T_{f,n})$  are annihilators of each other under the local Tate pairing  $\langle \cdot, \cdot \rangle_\ell$ . The same is true of  $H_{\text{ord}}^1(K_{\infty,\ell}, A_{f,n})$  and  $\hat{H}_{\text{ord}}^1(K_{\infty,\ell}, T_{f,n})$  for  $\ell \nmid N$ . In particular,  $H_{\text{fin}}^1(K_{\infty,\ell}, A_{f,n})$  and  $\hat{H}_{\text{sing}}^1(K_{\infty,\ell}, T_{f,n})$  are the Pontryagin duals of each other.*

*Proof.* The result over the finite layers  $K_m$  follows from standard properties of the local Tate pairing (cf. [DDT, §2.3]), and is then deduced over  $K_\infty$  by passage to the limit.

Proposition 2.3 yields a perfect pairing of  $\Lambda$ -modules (denoted by the same symbols  $\langle \cdot, \cdot \rangle_\ell$  by abuse of notation)

$$(26) \quad \langle \cdot, \cdot \rangle_\ell : \hat{H}_{\text{sing}}^1(K_{\infty,\ell}, T_{f,n}) \times H_{\text{fin}}^1(K_{\infty,\ell}, A_{f,n}) \rightarrow \mathbb{Q}_p/\mathbb{Z}_p.$$

The following lemma makes explicit the structure of the local cohomology groups  $\hat{H}^1(K_{\infty,\ell}, T_{f,n})$  and  $H^1(K_{\infty,\ell}, A_{f,n})$ .

**LEMMA 2.4.** *Suppose that  $\ell$  is a rational prime which does not divide  $N$ . If  $\ell$  is split in  $K/\mathbb{Q}$ , then*

$$\hat{H}_{\text{sing}}^1(K_{\infty,\ell}, T_{f,n}) = 0, \quad H_{\text{fin}}^1(K_{\infty,\ell}, A_{f,n}) = 0.$$

*Proof.* Because  $(\ell) = \lambda_1 \lambda_2$  is split in  $K/\mathbb{Q}$ , the Frobenius element attached to  $\lambda_1$  topologically generates a subgroup of finite index in  $G_\infty$ . Hence  $K_{\infty,\ell}$  is isomorphic to a direct sum of a finite number of copies of the unramified  $\mathbb{Z}_p$ -extension of  $\mathbb{Q}_\ell$ . Since  $A_{f,n}$  is of exponent  $p^n$ , any unramified cohomology class in  $H^1(K_{m,\ell}, A_{f,n})$  becomes trivial after restriction to  $H^1(K_{m',\ell}, A_{f,n})$  for  $m'$  sufficiently large. This implies the second assertion; the first follows from the nondegeneracy of the local Tate pairing displayed in (26).

The primes  $\ell$  which are inert in  $K/\mathbb{Q}$  exhibit a markedly different behaviour, because they split completely in the anticyclotomic tower. It is the presence of such primes which accounts for some of the essential differences between the anticyclotomic theory and the more familiar Iwasawa theory of the cyclotomic  $\mathbb{Z}_p$ -extension.

LEMMA 2.5. *If  $\ell$  does not divide  $N$  and is inert in  $K/\mathbb{Q}$ , then*

$$\hat{H}_{\text{sing}}^1(K_{\infty,\ell}, T_{f,n}) \simeq H_{\text{sing}}^1(K_{\ell}, T_{f,n}) \otimes \Lambda,$$

and

$$H_{\text{fin}}^1(K_{\infty,\ell}, A_{f,n}) \simeq \text{Hom}(H_{\text{sing}}^1(K_{\ell}, T_{f,n}) \otimes \Lambda, \mathbb{Q}_p/\mathbb{Z}_p).$$

*Proof.* Since  $\ell$  is inert in  $K$  and  $K_{\infty}/\mathbb{Q}$  is an extension of dihedral type, the Frobenius element at  $\ell$  in  $\text{Gal}(K_{\infty}/\mathbb{Q})$  is of order two and hence  $\ell$  splits completely in  $K_{\infty}/K$ . The choice of a prime  $\lambda_m$  of  $K_m$  above  $\ell$  thus determines an isomorphism  $H^1(K_{m,\ell}, T_{f,n}) \rightarrow H^1(K_{\ell}, T_{f,n}) \otimes \mathbb{Z}_p[G_m]$ . Choosing a compatible sequence of primes  $\lambda_m$  of  $K_m$  which lie above each other, one obtains an isomorphism

$$\hat{H}^1(K_{\infty,\ell}, T_{f,n}) \simeq H^1(K_{\ell}, T_{f,n}) \otimes \Lambda,$$

from the definition of the completed group ring  $\Lambda$ . The first isomorphism of the lemma now follows by passing to the singular parts of the cohomology, while the second is a consequence of Proposition 2.3.

*Admissible primes.* A rational prime  $\ell$  is said to be *n-admissible* relative to  $f$  if it satisfies the following conditions:

- (1)  $\ell$  does not divide  $N = pN^+N^-$ ;
- (2)  $\ell$  is inert in  $K/\mathbb{Q}$ ;
- (3)  $p$  does not divide  $\ell^2 - 1$ ;
- (4)  $p^n$  divides  $\ell + 1 - a_{\ell}$  or  $\ell + 1 + a_{\ell}$ .

A 1-admissible prime will simply be called *admissible* (so that in particular any *n-admissible* prime is admissible).

Note that if  $\ell$  is *n-admissible*, the module  $T_{f,n}$  is unramified at  $\ell$  and the Frobenius element over  $\mathbb{Q}$  at  $\ell$  acts semisimply on this module with eigenvalues  $\pm\ell$  and  $\pm 1$  which are distinct modulo  $p$ . From this a direct calculation shows that:

LEMMA 2.6. *The local cohomology groups  $H_{\text{sing}}^1(K_{\ell}, T_{f,n})$  and  $H_{\text{fin}}^1(K_{\ell}, T_{f,n})$  are both isomorphic to  $\mathbb{Z}/p^n\mathbb{Z}$ .*

*Proof.* The group  $H_{\text{sing}}^1(K_{\ell}, T_{f,n})$  is identified with  $H^1(I_{\ell}, T_{f,n})^{G_{K_{\ell}}}$ . Since  $T_{f,n}$  is unramified at  $\ell$ , this first cohomology group is identified with a group of homomorphisms, which necessarily factor through the tame inertia group at  $\ell$ . The Frobenius element over  $K$  at  $(\ell)$  acts on this tame inertia group as multiplication by  $\ell^2$ , while it acts on  $T_{f,n}$  with eigenvalues  $\ell^2$  and 1. The result follows from this, in light of the fact that  $p$  does not divide  $\ell^2 - 1$ . Similarly, the group  $H_{\text{fin}}^1(K_{\ell}, A_{f,n})$  is identified with the  $G_{K_{\ell}}$ -coinvariants of  $A_{f,n}$  which are also isomorphic to  $\mathbb{Z}/p^n\mathbb{Z}$ .

LEMMA 2.7. *The local groups  $\hat{H}_{\text{sing}}^1(K_{\infty,\ell}, T_{f,n})$  and  $\hat{H}_{\text{fin}}^1(K_{\infty,\ell}, T_{f,n})$  are each free of rank one over  $\Lambda/p^n\Lambda$ .*

*Proof.* Since  $\ell$  is inert in  $K/\mathbb{Q}$ , Lemma 2.5 implies that  $\hat{H}^1(K_{\ell,\infty}, T_{f,n})$  is isomorphic to  $H^1(K_{\ell}, T_{f,n}) \otimes \Lambda$ . The result now follows from Lemma 2.6.

*Remarks.* 1. Note that the  $n$ -admissible primes are not the primes appearing in Kolyvagin's study of the Selmer groups of elliptic curves, where the condition that  $p^n$  divides  $\ell + 1$  was imposed.

2. The notion of admissible prime introduced here is similar to the one introduced in [BD0, Def. 2.20], the main difference arising from the fact that the local cohomology groups  $H_{\text{fin}}^1(K_{\ell}, A_{f,n})$  and  $H_{\text{sing}}^1(K_{\ell}, T_{f,n})$  are both free of rank one (and not two) over  $\mathbb{Z}/p^n\mathbb{Z}$ .

2.3. *Definition of the Selmer group.* Let  $\ell$  be a prime not dividing  $N$ . Composing restriction from  $K_{\infty}$  to  $K_{\infty,\ell}$  with  $\partial_{\ell}$  yields residue maps on the global cohomology groups, still denoted  $\partial_{\ell}$  by an abuse of notation,

$$(27) \quad \partial_{\ell} : H^1(K_{\infty}, A_{f,n}) \rightarrow H_{\text{sing}}^1(K_{\infty,\ell}, A_{f,n}),$$

$$(28) \quad \partial_{\ell} : \hat{H}^1(K_{\infty}, T_{f,n}) \rightarrow \hat{H}_{\text{sing}}^1(K_{\infty,\ell}, T_{f,n}).$$

Note that if  $\ell$  is split in  $K/\mathbb{Q}$ , the residue map of (28) is 0 by Lemma 2.4.

If  $\partial_{\ell}(\kappa) = 0$  for  $\kappa \in H^1(K_{\infty}, A_{f,n})$  (resp.  $\hat{H}^1(K_{\infty}, T_{f,n})$ ), let

$$v_{\ell}(\kappa) \in H_{\text{fin}}^1(K_{\infty,\ell}, A_{f,n}) \quad (\text{resp. } \hat{H}_{\text{fin}}^1(K_{\infty,\ell}, T_{f,n}))$$

denote the natural image of  $\kappa$  under the restriction map at  $\ell$ .

*Definition 2.8.* The *Selmer group*  $\text{Sel}_{f,n}$  attached to  $f$ ,  $n$  and  $K_{\infty}$  is the group of elements  $s$  in  $H^1(K_{\infty}, A_{f,n})$  satisfying

- (1)  $\partial_{\ell}(s) = 0$  for all rational primes  $\ell$  not dividing  $N$ .
- (2) The class  $s$  is *ordinary* at the primes  $\ell|N^-p$ .
- (3) The class  $s$  is *trivial* at the primes  $\ell|N^+$ .

*Caveat.* Note that the group  $\text{Sel}_{f,n}$  depends on the value of  $N$ , hence on the modular form  $f$  itself, and not just on the Galois representation  $A_{f,n}$  attached to it. The same remark holds for the *compactified Selmer group*  $\hat{H}_S^1(K_{\infty}, T_{f,n})$  defined below:

*Definition 2.9.* Let  $S$  be a square-free integer which is relatively prime to  $N$ . The *compactified Selmer group*  $\hat{H}_S^1(K_{\infty}, T_{f,n})$  attached to  $f$ ,  $S$  and  $K_{\infty}$  is the group of elements  $\kappa$  in  $\hat{H}^1(K_{\infty}, T_{f,n})$  satisfying

- (1)  $\partial_{\ell}(\kappa) = 0$  for all rational primes  $\ell$  not dividing  $SN$ ;

- (2) The class  $\kappa$  is *ordinary* at the primes  $\ell | N^- p$ .
- (3) The class  $\kappa$  is *arbitrary* at the primes  $\ell$  dividing  $N^+$ , and at the primes dividing  $S$ .

*Global reciprocity.* Let  $\kappa \in \hat{H}^1(K_\infty, T_{f,n})$  and let  $s \in H^1(K_\infty, A_{f,n})$  be global cohomology classes. For each rational prime  $q$ , let  $\kappa_q$  and  $s_q$  denote the restrictions of these cohomology classes to the (semi-)local cohomology group attached to the prime  $q$ . The global reciprocity law of class field theory implies that

$$(29) \quad \sum_q \langle \kappa_q, s_q \rangle_q = 0,$$

where the sum is taken over all the rational primes. In particular, if  $\kappa$  belongs to  $\hat{H}_S^1(K_\infty, T_{f,n})$  and  $s$  belongs to  $\text{Sel}_{f,n}$ , then since the local conditions defining these two groups are orthogonal at the primes not dividing  $S$ , and since  $s$  has trivial residue at the primes dividing  $S$ , formula (29) becomes:

$$\sum_{q|S} \langle \partial_q(\kappa), v_q(s) \rangle_q = 0.$$

Of particular interest is the following special case:

PROPOSITION 2.10. *Suppose that  $\kappa$  belongs to  $\hat{H}_\ell^1(K_\infty, A_{f,n})$ . Then*

$$\langle \partial_\ell(\kappa), v_\ell(s) \rangle_\ell = 0,$$

for all  $s \in \text{Sel}_{f,n}$ .

The strategy of the proof of Theorem 1 is to produce, for sufficiently many primes  $\ell$  that are inert in  $K$ , cohomology classes  $\kappa(\ell) \in \hat{H}_\ell^1(K_\infty, A_{f,n})$  whose residue  $\partial_\ell(\kappa(\ell))$  can be related to the  $p$ -adic  $L$ -function  $L_p(f, K)$  constructed in Section 1. Thanks to Proposition 2.10, the elements  $\partial_\ell(\kappa(\ell))$  yield relations in a presentation for  $\text{Sel}_{f,n}^\vee$ .

### 3. Some preliminaries

3.1.  $\Lambda$ -modules. If  $X$  is any module over a ring  $R$ , let  $\text{Fitt}_R(X)$  denote the Fitting ideal of  $X$  over  $R$ . If  $R = \Lambda$  and  $X$  is finitely generated, let  $\text{Char}(X)$  denote the characteristic ideal attached to  $X$ .

PROPOSITION 3.1. *Let  $X$  be a finitely generated  $\Lambda$ -module and let  $\mathcal{L}$  be an element of  $\Lambda$ . Suppose that  $\varphi(\mathcal{L})$  belongs to  $\text{Fitt}_{\mathcal{O}}(X \otimes_\varphi \mathcal{O})$ , for all homomorphisms  $\varphi : \Lambda \rightarrow \mathcal{O}$ , where  $\mathcal{O}$  is a discrete valuation ring. Then  $\mathcal{L}$  belongs to  $\text{Char}(X)$ .*

*Proof.* If  $X$  is not  $\Lambda$ -torsion, then  $\text{Fitt}_\Lambda(X) = 0$ . Since

$$\text{Fitt}_{\mathcal{O}}(X \otimes_{\varphi} \mathcal{O}) = \varphi(\text{Fitt}_\Lambda(X)),$$

it follows that  $\varphi(\mathcal{L}) = 0$  for all  $\varphi$ . This implies (by the Weierstrass preparation theorem, for example) that  $\mathcal{L} = 0$ . Hence one may assume without loss of generality that  $X$  is a  $\Lambda$ -torsion module. In that case the structure theory of  $\Lambda$ -modules ensures the existence of an exact sequence of  $\Lambda$ -modules:

$$(30) \quad X \xrightarrow{j} \bigoplus_i \Lambda/(g_i) \longrightarrow C \longrightarrow 0,$$

where  $C$  and  $\ker j$  are finite  $\Lambda$ -modules and the  $g_i$  are nonzero distinguished polynomials or powers of  $p$ . By definition,  $g := \prod_i g_i$  is a generator of  $\text{Char}(X)$ . Since  $C$  is finite, its  $\Lambda$ -Fitting ideal can be generated by two elements  $\iota_1$  and  $\iota_2$  having no common irreducible factors. By tensoring the exact sequence (30) with  $\mathcal{O}$  one finds that

$$\varphi(\iota_i)\text{Fitt}_{\mathcal{O}}(X \otimes_{\varphi} \mathcal{O}) \subset (\varphi(g)), \quad \text{for } i = 1, 2.$$

It follows by assumption that  $\varphi(g)$  divides  $\varphi(\iota_i \mathcal{L})$  for all  $\varphi$ . Hence (as can be seen by using the Weierstrass preparation theorem)  $g$  divides  $\iota_i \mathcal{L}$  for  $i = 1, 2$ , and therefore  $g$  divides  $\mathcal{L}$ .

**3.2. Controlling the Selmer group.** Suppose now that  $A_{f,1}$  satisfies the irreducibility condition 2 of Assumption 6.

**THEOREM 3.2.** *Let  $s$  be a nonzero element of  $H^1(K, A_{f,1})$ . There exist infinitely many  $n$ -admissible primes  $\ell$  relative to  $f$  such that  $\partial_{\ell}(s) = 0$  and  $v_{\ell}(s) \neq 0$ .*

*Proof.* Let  $\mathbb{Q}(A_{f,n})$  be the extension of  $\mathbb{Q}$  fixed by the kernel of the Galois representation  $A_{f,n}$ . It is unramified at the primes not dividing  $N$ . Since the discriminant of  $K$  is assumed to be prime to  $N$ , the fields  $\mathbb{Q}(A_{f,n})$  and  $K$  are linearly disjoint. Letting  $M$  denote the compositum of these fields, there is therefore a natural inclusion

$$\text{Gal}(M/\mathbb{Q}) = \text{Gal}(K/\mathbb{Q}) \times \text{Gal}(\mathbb{Q}(A_{f,n})/\mathbb{Q}) \subset \{1, \tau\} \times \text{Aut}_{\mathbb{Z}/p^n\mathbb{Z}}(A_{f,n}),$$

so that elements of  $\text{Gal}(M/\mathbb{Q})$  can be labelled by certain pairs  $(\tau^j, T)$  with  $j \in \{0, 1\}$  and  $T \in \text{Aut}_{\mathbb{Z}/p^n\mathbb{Z}}(A_{f,n})$ . Let  $M_s$  be the extension of  $M$  cut out by the image  $\bar{s}$  of  $s$  under restriction to  $H^1(M, A_{f,1}) = \text{Hom}(\text{Gal}(\bar{M}/M), A_{f,1})$ . Assume without loss of generality that  $s$  belongs to a specific eigenspace for the action of  $\tau$ , so that

$$\tau s = \delta s, \quad \text{for some } \delta \in \{1, -1\}.$$

Under this assumption, the extension  $M_s$  is Galois over  $\mathbb{Q}$ , not merely over  $K$ . In fact, by the assumption that  $A_{f,1}$  is an irreducible  $G_{\mathbb{Q}}$ -module,  $\text{Gal}(M_s/\mathbb{Q})$

is identified with the semi-direct product

$$(31) \quad \mathrm{Gal}(M_s/\mathbb{Q}) = A_{f,1} \rtimes \mathrm{Gal}(M/\mathbb{Q}),$$

where the quotient  $\mathrm{Gal}(M/\mathbb{Q})$  acts on the abelian normal subgroup  $A_{f,1}$  of  $\mathrm{Gal}(M_s/\mathbb{Q})$  by the rule

$$(32) \quad (\tau^j, T)(v) = \delta^j \bar{T}v.$$

Here  $\bar{T}$  denotes the natural image of  $T$  in  $\mathrm{Aut}_{\mathbb{F}_p}(A_{f,1})$ . By part 2 of Assumption 6 on the Galois representation  $A_{f,1}$ , the group  $\mathrm{Gal}(M_s/\mathbb{Q})$  contains an element of the form  $(v, \tau, T)$ , where

1. The automorphism  $T$  has eigenvalues  $\delta$  and  $\lambda$ , where  $\lambda \in (\mathbb{Z}/p^n\mathbb{Z})^\times$  is not equal to  $\pm 1 \pmod{p}$  and has order prime to  $p$ . (Note that here the assumption that  $p > 3$  is needed.)

2. The vector  $v \in A_{f,1}$  is nonzero and belongs to the  $\delta$ -eigenspace for  $\bar{T}$ .

Let  $\ell \nmid N$  be a rational prime which is unramified in  $M_s/\mathbb{Q}$  and satisfies

$$(33) \quad \mathrm{Frob}_\ell(M_s/\mathbb{Q}) = (v, \tau, T).$$

By the Chebotarev density theorem, there exist infinitely many such primes. In fact, the set of such primes has positive density. The fact, immediate from (33), that  $\mathrm{Frob}_\ell(M/\mathbb{Q}) = (\tau, T)$  implies that  $\ell$  is an admissible prime. To see that  $v_\ell(s) \neq 0$ , choose a prime  $\lambda$  of  $M$  above  $\ell$ , and let  $d$  be the (necessarily even) degree of the corresponding residue field extension. Then

$$\mathrm{Frob}_\lambda(M_s/M) = (v, \tau, T)^d = v + \delta \bar{T}v + \bar{T}^2v + \cdots + \delta \bar{T}^{d-1}v = dv.$$

Let  $\bar{s}$  denote the image of  $s$  in

$$H^1(M, A_{f,1}) = \mathrm{Hom}(\mathrm{Gal}(\bar{M}/M), A_{f,1})$$

under restriction. Since  $d$  is prime to  $p$  by property 1 of  $T$ , it follows that  $\bar{s}(\mathrm{Frob}_\lambda(M_s/M)) = d\bar{s}(v) \neq 0$ , so that the restriction at  $\lambda$  of  $\bar{s}$  is nonzero. Hence, so is  $v_\ell(s)$ , *a fortiori*.

*Global cohomology groups.* By [BD0, Def. 2.22], a finite set  $S$  of primes is said to be *n-admissible* relative to  $f$  if

1. All  $\ell \in S$  are *n-admissible* relative to  $f$ .
2. The map  $\mathrm{Sel}(K, A_{f,n}) \longrightarrow \bigoplus_{\ell \in S} H_{\mathrm{fin}}^1(K_\ell, A_{f,n})$  is injective.

A direct argument based on Theorem 3.2 shows that *n-admissible* sets exist. (See also the proof of Lemma 2.23 of [BD0].) In fact, any finite collection of *n-admissible* primes can be enlarged to an *n-admissible* set.

**PROPOSITION 3.3.** *If  $S$  is an *n-admissible* set, then the group  $\hat{H}_S^1(K_\infty, T_{f,n})$  is free of rank  $\#S$  over  $\Lambda/p^n\Lambda$ .*

*Proof.* The fact that  $H_S^1(K_m, T_{f,n})$  is free over  $\mathbb{Z}/p^n\mathbb{Z}[G_m]$  is essentially Theorem 3.2 of [BD0], whose proof carries over, *mutatis mutandis*, to the present context with its slightly modified notion of admissible prime. Proposition 3.3 follows by passing to the limit as  $m \rightarrow \infty$ .

**THEOREM 3.4.** *Let  $\mathfrak{m}_\Lambda$  denote the maximal ideal of  $\Lambda$ . Then*

- (1) *The natural map from  $H^1(K, A_{f,1}) \rightarrow H^1(K_\infty, A_{f,n})[\mathfrak{m}_\Lambda]$  induced by restriction is an isomorphism.*
- (2) *If  $S$  is an  $n$ -admissible set, the natural map from  $\hat{H}_S^1(K_\infty, T_{f,n})/\mathfrak{m}_\Lambda$  to  $H^1(K, T_{f,1})$  induced by corestriction is injective.*

*Proof.* Let  $I_\Lambda$  denote the augmentation ideal of  $\Lambda$ . The inflation-restriction sequence from  $K$  to  $K_m$  gives the exact sequence

$$\begin{aligned} H^1(K_m/K, A_{f,n}^{G_{K^m}}) &\longrightarrow H^1(K, A_{f,n}) \xrightarrow{j} H^1(K_m, A_{f,n})[I_\Lambda] \longrightarrow \\ &\longrightarrow H^2(K_m/K, A_{f,n}^{G_{K^m}}). \end{aligned}$$

By part 2 of Assumption 6 in the introduction, the module  $A_{f,n}^{G_{K^m}}$  is trivial. (Otherwise, the Galois representation attached to  $A_{f,n}$  would have solvable image, contradicting the hypotheses that were made in the introduction.) Hence the map  $j$  is an isomorphism. Taking the  $G_K$ -cohomology of the exact sequence

$$0 \longrightarrow A_{f,1} \longrightarrow A_{f,n} \xrightarrow{p} A_{f,n-1} \longrightarrow 0$$

and using the fact that  $A_{f,1}^{G_K} = 0$  once again, we see that the natural map

$$(34) \quad H^1(K, A_{f,1}) \longrightarrow H^1(K, A_{f,n})[p]$$

is an isomorphism. It follows that the natural map

$$H^1(K, A_{f,1}) \longrightarrow H^1(K_m, A_{f,n})[\mathfrak{m}_\Lambda]$$

is an isomorphism as well. Part 1 of Theorem 3.4 follows by taking the direct limit as  $m \rightarrow \infty$ .

Part 2 of Theorem 3.4 follows directly from Proposition 3.3.

**3.3. Rigid pairs.** Let  $W_f := \text{ad}_0(A_{f,1}) = \text{Hom}_0(A_{f,1}, A_{f,1})$  be the adjoint representation attached to  $A_{f,1}$ , i.e., the vector space of trace zero endomorphisms of  $A_{f,1}$ . It is a three-dimensional  $\mathbb{F}_p$ -vector space endowed with a natural action of  $G_\mathbb{Q}$  arising from the conjugation of endomorphisms. Write  $W_f^* := \text{Hom}(W_f, \mu_p)$  for the Kummer dual of  $W_f$ .

Recall that  $A_{f,1}$  is ordinary at  $p$ , so that there is an exact sequence of  $I_p$ -modules

$$0 \longrightarrow A_{f,1}^{(p)} \longrightarrow A_{f,1} \longrightarrow A_{f,1}^{(1)} \longrightarrow 0$$

where  $A_{f,1}^{(p)}$  represents the subspace on which  $I_p$  acts via the cyclotomic character  $\varepsilon$ , and  $A_{f,1}^{(1)}$  represents the  $I_p$ -coinvariants of  $A_{f,1}$ . Let

$$W_f^{(p)} := \text{Hom}(A_{f,1}^{(1)}, A_{f,1}^{(p)}).$$

It is an  $I_p$ -submodule of  $W_f$ ; let  $W_f^{(1)} := W_f/W_f^{(p)}$ . The classes in  $H^1(\mathbb{Q}_p, W_f)$  whose restriction at  $p$  belong to  $H^1(I_p, W_f^{(p)})$  are called *ordinary* at  $p$ .

Likewise, if  $\ell$  is a prime which divides  $N$  exactly, recall the submodules  $A_{f,1}^{(\ell)}$  and  $A_{f,1}^{(1)}$  on which  $G_{\mathbb{Q}_\ell}$  acts by  $\pm\varepsilon$  and  $\pm 1$  respectively. (These submodules are well-defined, by virtue of Assumption 2.1.) Set

$$W_f^{(\ell)} := \text{Hom}(A_{f,1}^{(1)}, A_{f,1}^{(\ell)}).$$

The classes in  $H^1(\mathbb{Q}, W_f)$  whose restriction at  $\ell$  belongs to  $H^1(\mathbb{Q}_\ell, W_f^{(\ell)})$  are called *ordinary* at  $\ell$ .

If  $\ell$  is an admissible prime for  $f$ , the eigenvalues of  $\text{Frob}_\ell$  acting on the Galois representation  $A_{f,1}$  are  $\pm 1$  and  $\pm\ell$ . Recall also that  $\ell^2 \neq 1$  belongs to  $\mathbb{F}_p^\times$ . Therefore, the eigenvalues of  $\text{Frob}_\ell$  acting on  $W_f$  (resp.  $W_f^*$ ) are the distinct elements  $1, \ell$ , and  $\ell^{-1}$  (resp.  $\ell, 1$  and  $\ell^2$ ) of  $\mathbb{F}_p^\times$ . Let  $W_f^{(\ell)}$  and  $W_f^{*(\ell)}$  denote the one-dimensional  $\mathbb{F}_p$ -subspace on which  $\text{Frob}_\ell$  acts with eigenvalue  $\ell$ . The classes in  $H^1(\mathbb{Q}, W_f)$  whose restrictions at  $\ell$  belong to  $H^1(\mathbb{Q}_\ell, W_f^{(\ell)})$  are called *ordinary* at  $\ell$ . (In [Ram, §3], these classes are referred to as *null cocycles*.) Note that  $H^1(\mathbb{Q}_\ell, W_f)$  decomposes as a direct sum of two one-dimensional  $\mathbb{F}_p$ -subspaces,

$$H^1(\mathbb{Q}_\ell, W_f) = H_{\text{fin}}^1(\mathbb{Q}_\ell, W_f) \oplus H_{\text{ord}}^1(\mathbb{Q}_\ell, W_f^{(\ell)}),$$

where

$$H_{\text{fin}}^1(\mathbb{Q}_\ell, W_f) := H^1(\mathbb{Q}_\ell^{\text{nr}}/\mathbb{Q}_\ell, W_f)$$

is the space of unramified cocycles. A similar remark holds for  $W_f^*$ .

Let  $S$  be a square-free product of admissible primes for  $f$ .

*Definition 3.5.* The  $S$ -Selmer group attached to  $W_f$ , denoted  $\text{Sel}_S(\mathbb{Q}, W_f)$ , is the subspace of cohomology classes  $\xi \in H^1(\mathbb{Q}, W_f)$  satisfying

- (1) For all  $\ell$  which do not divide  $NS$ , the image of  $\xi$  in  $H^1(\mathbb{Q}_\ell, W_f)$  belongs to  $H_{\text{fin}}^1(\mathbb{Q}_\ell, W_f)$ .
- (2) The class  $\xi$  is *ordinary* at the primes  $\ell$  dividing  $NS$  exactly.
- (3) The class  $\xi$  belongs to the kernel of the restriction to  $H^1(I_\ell, W_f)$  if  $\ell$  is a prime such that  $\ell^2$  divides  $N^+$ .

Similar definitions can be made for  $\text{Sel}_S(\mathbb{Q}, W_f^*)$ . Note that  $H^1(\mathbb{Q}_\ell, W_f^{(\ell)})$  and  $H^1(\mathbb{Q}_\ell, W_f^{*(\ell)})$  are orthogonal to each other under the local Tate pairing.

PROPOSITION 3.6. *The modular form  $f$  is  $p$ -isolated if and only if  $\text{Sel}_1(\mathbb{Q}, W_f)$  is trivial.*

*Proof.* Let  $R$  denote the universal ring attached to deformations  $\rho$  of the Galois representation  $A_{f,1}$ , satisfying

- (1) The determinant of  $\rho$  is the cyclotomic character describing the action of  $G_{\mathbb{Q}}$  on the  $p$ -power roots of unity.
- (2)  $\rho$  is unramified outside  $NS$ .
- (3)  $\rho$  is ordinary at  $p$ , i.e., the restriction of  $\rho$  to  $I_p$  is of the form  $\begin{pmatrix} \varepsilon & * \\ 0 & 1 \end{pmatrix}$ .
- (4) For all  $\ell$  dividing  $N^+N^-S$  exactly, the restriction of  $\rho$  to a decomposition group at  $\ell$  is ordinary, i.e., is of the form  $\begin{pmatrix} \varepsilon & * \\ 0 & 1 \end{pmatrix}$ .

The ring  $R$  is a complete local Noetherian  $\mathbb{Z}_p$ -algebra with residue field  $\mathbb{F}_p$ . Let  $\mathfrak{m}$  denote the maximal ideal of  $R$ . Standard results of deformation theory (cf. Lemma 2.39 and Sections 2.6 and 2.7 of [DDT]) identify  $\mathfrak{m}/(p, \mathfrak{m}^2)$  with the Pontryagin dual of  $\text{Sel}_S(\mathbb{Q}, W_f)$ . It follows that  $R = \mathbb{Z}_p$  if and only if  $\text{Sel}_S(\mathbb{Q}, W_f)$  is trivial. When  $S = 1$ , a calculation as in [W, §3] shows that the ring  $R$  surjects onto the ring  $\mathbb{T}_f$  of Hecke operators acting on the space  $S_2(\mathcal{T}/\Gamma)$ , completed at the maximal ideal attached to  $f$ . A deep result of Wiles ([W], [DDT, Th. 3.42]) asserts that this surjection is an isomorphism. Hence the fact that  $R = \mathbb{Z}_p$  is equivalent to the fact that  $\mathbb{T}_f = \mathbb{Z}_p$ , which in turn is equivalent to the fact that the modular form  $f$  is  $p$ -isolated.

If  $S$  is a square-free product of admissible primes for  $A_{f,1}$ , let  $\text{Sel}_{(S)}(\mathbb{Q}, W_f)$  denote the Selmer group defined in the same way as  $\text{Sel}_S(\mathbb{Q}, W_f)$  above, but with no condition imposed at the primes of  $S$ . Let  $\text{Sel}_{[S]}(\mathbb{Q}, W_f)$  denote the subgroup of  $\text{Sel}_S(\mathbb{Q}, W_f)$  consisting of classes that are trivial at the primes in  $S$ . This notation can be combined: thus, if  $S_1, S_2, S_3$  are pairwise coprime square-free products of admissible primes, the group  $\text{Sel}_{S_1(S_2)[S_3]}(\mathbb{Q}, W_f)$  is given the obvious meaning. Similar definitions can be made with  $W_f$  replaced by  $W_f^*$ . Note that the Selmer groups  $\text{Sel}_{(S)}(\mathbb{Q}, W_f)$  and  $\text{Sel}_{[S]}(\mathbb{Q}, W_f^*)$  are dual Selmer groups in the sense of [DDT, §2.3], and the same is true of  $\text{Sel}_S(\mathbb{Q}, W_f)$  and  $\text{Sel}_S(\mathbb{Q}, W_f^*)$ .

PROPOSITION 3.7. *If  $f$  is  $p$ -isolated, and  $\ell$  is an admissible prime for  $f$ , then  $\text{Sel}_{(\ell)}(\mathbb{Q}, W_f)$  and  $\text{Sel}_{(\ell)}(\mathbb{Q}, W_f^*)$  are one-dimensional  $\mathbb{F}_p$ -vector spaces.*

*Proof.* It follows from a direct calculation of orders of local cohomology groups, combined with Theorem 2.18 of [DDT], that the groups  $\text{Sel}_1(\mathbb{Q}, W_f)$  and  $\text{Sel}_1(\mathbb{Q}, W_f^*)$  have the same cardinality. By Proposition 3.6, both these

groups are trivial. Applying Theorem 2.18 of [DDT] once more, and using the fact that  $p$  divides  $(\ell + 1)^2 - a_\ell^2$ , one finds that

$$\#\mathrm{Sel}_{(\ell)}(\mathbb{Q}, W_f) / \#\mathrm{Sel}_{[\ell]}(\mathbb{Q}, W_f^*) = p.$$

Hence  $\mathrm{Sel}_{(\ell)}(\mathbb{Q}, W_f)$  is one-dimensional over  $\mathbb{F}_p$ . The same argument, with  $W_f$  and  $W_f^*$  interchanged, shows that  $\mathrm{Sel}_{(\ell)}(\mathbb{Q}, W_f^*)$  is one-dimensional as well.

Suppose that  $f$  is  $p$ -isolated so that the conclusion of Proposition 3.7 holds. If  $\ell \neq \ell_1$  is any admissible prime, write

$$v_\ell : \mathrm{Sel}_{(\ell_1)}(\mathbb{Q}, W_f) \longrightarrow H_{\mathrm{fin}}^1(\mathbb{Q}_\ell, W_f), \quad v_\ell^* : \mathrm{Sel}_{(\ell_1)}(\mathbb{Q}, W_f^*) \longrightarrow H_{\mathrm{fin}}^1(\mathbb{Q}_\ell, W_f^*)$$

for the natural maps induced from restriction at  $\ell$ .

**PROPOSITION 3.8.** (1) *If  $\mathrm{Sel}_{\ell_1}(\mathbb{Q}, W_f) \neq 0$ , and  $v_{\ell_2}$  and  $v_{\ell_2}^*$  are both nonzero, then  $\mathrm{Sel}_{\ell_1 \ell_2}(\mathbb{Q}, W_f) = 0$ .*

(2) *If  $\mathrm{Sel}_{\ell_1}(\mathbb{Q}, W_f) = 0$ , and  $v_{\ell_2}$  is 0, then either  $\mathrm{Sel}_{\ell_1 \ell_2}(\mathbb{Q}, W_f) = 0$  or  $\mathrm{Sel}_{\ell_2}(\mathbb{Q}, W_f)$  is one-dimensional.*

*Proof.* 1. Since  $\ell_1$  is admissible, we may invoke Proposition 3.7 and let  $\xi_1$  and  $\xi_1^*$  be generators of  $\mathrm{Sel}_{(\ell_1)}(\mathbb{Q}, W_f)$  and  $\mathrm{Sel}_{(\ell_1)}(\mathbb{Q}, W_f^*)$  respectively. Note that  $\xi_1$  belongs to  $\mathrm{Sel}_{\ell_1}(\mathbb{Q}, W_f)$  (i.e., the restriction of  $\xi_1$  at  $\ell_1$  is ordinary) since  $\mathrm{Sel}_{\ell_1}(\mathbb{Q}, W_f) \neq 0$ . A calculation using Theorem 2.18 of [DDT] shows that  $\mathrm{Sel}_{\ell_1}(\mathbb{Q}, W_f)$  and  $\mathrm{Sel}_{\ell_1}(\mathbb{Q}, W_f^*)$  have the same dimension, and hence  $\xi_1^*$  also belongs to  $H_{\ell_1}^1(\mathbb{Q}, W_f^*)$ .

By Theorem 2.18 of [DDT],

$$\#\mathrm{Sel}_{\ell_1(\ell_2)}(\mathbb{Q}, W_f) / \#\mathrm{Sel}_{\ell_1[\ell_2]}(\mathbb{Q}, W_f^*) = p.$$

The assumption that  $v_{\ell_2}(\xi_1^*) \neq 0$  implies that  $\#\mathrm{Sel}_{\ell_1[\ell_2]}(\mathbb{Q}, W_f^*) = 0$ . Hence  $\mathrm{Sel}_{\ell_1(\ell_2)}(\mathbb{Q}, W_f) = \mathrm{Sel}_{\ell_1}(\mathbb{Q}, W_f)$  is one-dimensional and spanned by  $\xi_1$ . Now the assumption that  $v_{\ell_2}(\xi_1) \neq 0$  shows that  $\mathrm{Sel}_{\ell_1 \ell_2}(\mathbb{Q}, W_f) = 0$ .

2. It is convenient to distinguish two cases:

a) If  $\mathrm{Sel}_{\ell_2}(\mathbb{Q}, W_f) = 0$ , then the space  $\mathrm{Sel}_{(\ell_1)\ell_2}$  is one-dimensional by Theorem 2.18 of [DDT]. The assumption that  $v_{\ell_2}(\xi_1) = 0$  shows that in fact  $\xi_1$  generates  $\mathrm{Sel}_{(\ell_1)\ell_2}(\mathbb{Q}, W_f)$ . But  $\xi_1$  does not belong to  $\mathrm{Sel}_{\ell_1(\ell_2)}(\mathbb{Q}, W_f)$  since if it did it would also belong to  $\mathrm{Sel}_{\ell_1}(\mathbb{Q}, W_f)$  which is trivial by assumption. Hence

$$\mathrm{Sel}_{\ell_1 \ell_2}(\mathbb{Q}, W_f) = \mathrm{Sel}_{(\ell_1)\ell_2}(\mathbb{Q}, W_f) \cap \mathrm{Sel}_{\ell_1(\ell_2)}(\mathbb{Q}, W_f) = 0.$$

b) If  $\mathrm{Sel}_{\ell_2}(\mathbb{Q}, W_f) \neq 0$ , it is necessarily one-dimensional since it lies in the one-dimensional space  $\mathrm{Sel}_{(\ell_2)}(\mathbb{Q}, W_f)$ .

*Definition 3.9.* A pair  $(\ell_1, \ell_2)$  of admissible primes is said to be a *rigid pair* if the Selmer group  $\text{Sel}_{\ell_1 \ell_2}(\mathbb{Q}, W_f)$  is trivial.

In addition to Theorem 3.2 guaranteeing the existence of a plentiful supply of  $n$ -admissible primes sufficient to control the Selmer group  $\text{Sel}_{f,n}$ , there arises the need for the somewhat more technical Theorems 3.10 and 3.11 below guaranteeing the existence of a large supply of rigid pairs of  $n$ -admissible primes in certain favorable circumstances.

**THEOREM 3.10.** *Suppose that  $f$  is a  $p$ -isolated eigenform in  $S_2(\mathcal{T}/\Gamma)$ , and let  $\ell_1$  be an admissible prime for  $f$ . Let  $s$  be a nonzero class in  $H^1(K, A_{f,1})$ . For any  $n$ , there exist infinitely many  $n$ -admissible primes  $\ell_2$  such that*

- (1)  $\partial_{\ell_2}(s) = 0$  and  $v_{\ell_2}(s) \neq 0$ .
- (2) *Either  $(\ell_1, \ell_2)$  is a rigid pair, or  $\text{Sel}_{\ell_2}(\mathbb{Q}, W_f)$  is one-dimensional.*

*Proof.* Assume as in the proof of Theorem 3.2 that  $s$  belongs to a fixed eigenspace for complex conjugation, so that  $\tau s = \delta s$  for some  $\delta \in \{1, -1\}$ . Write  $M = K(A_{f,n})$  and let  $M_s$  be the Galois extension of  $M$  cut out by  $s$  as in the proof of Theorem 3.2.

The fact that  $f$  is  $p$ -isolated implies, by Proposition 3.7, that the Selmer groups  $\text{Sel}_{(\ell_1)}(\mathbb{Q}, W_f)$  and  $\text{Sel}_{(\ell_1)}(\mathbb{Q}, W_f^*)$  are one-dimensional over  $\mathbb{F}_p$ . Let  $\xi$  and  $\xi^*$  denote as before generators of these spaces. The image  $\bar{\xi}$ ,  $\bar{\xi}^*$  of  $\xi$ ,  $\xi^*$  in  $H^1(M, W_f) = \text{Hom}(G_M, W_f)$  and  $H^1(M, W_f^*)$  cuts out extensions  $M_{\bar{\xi}}$  and  $M_{\bar{\xi}^*}$  of  $M$  whose Galois groups are identified, via  $\bar{\xi}$  and  $\bar{\xi}^*$ , with  $W_f$  and  $W_f^*$  respectively.

Let  $M_{s, \xi, \xi^*}$  denote the compositum of  $M_s$ ,  $M_{\bar{\xi}}$ , and  $M_{\bar{\xi}^*}$  over  $M$ . Since the Galois representations  $A_{f,1}$ ,  $W_f$  and  $W_f^*$  are absolutely irreducible and pairwise nonisomorphic, the Galois group of  $M_{s, \xi, \xi^*}$  over  $\mathbb{Q}$  is isomorphic to the semi-direct product

$$\text{Gal}(M_{s, \xi, \xi^*}/\mathbb{Q}) = (A_{f,1} \times W_f \times W_f^*) \rtimes \text{Gal}(M/\mathbb{Q}),$$

where the action of the quotient  $\text{Gal}(M/\mathbb{Q})$  on the abelian normal subgroup  $(A_{f,1} \times W_f \times W_f^*)$  is given by

$$(\tau^j, T)(v, w, w^*) = (\delta^j \bar{T}v, \bar{T}w \bar{T}^{-1}, \bar{T}w^* \bar{T}^{-1} \det(T)).$$

*Case 1.* Suppose that  $\xi$  belongs to  $\text{Sel}_{\ell_1}(\mathbb{Q}, W_f)$ , so that  $\xi^*$  also belongs to  $\text{Sel}_{\ell_1}(\mathbb{Q}, W_f^*)$ . The group  $\text{Gal}(M_{s, \xi, \xi^*}/\mathbb{Q})$  contains an element of the form  $(v, w, w^*, \tau, T)$ , where

1. The transformation  $T$  acting on  $A_{f,n}$  has eigenvalues  $\delta$  and  $\lambda$ , where  $\lambda$  is an element of  $(\mathbb{Z}/p^n\mathbb{Z})^\times$  of order prime to  $p$  which is  $\neq \pm 1$ .

2. The vector  $v$  belongs to the unique line in  $A_{f,1}$  on which  $T$  acts by  $\delta$ .
3. The vector  $w$  (resp.  $w^*$ ) belongs to the unique line in  $W_f$  (resp.  $W_f^*$ ) which is fixed by  $T$ .

Let  $\ell_2$  be a rational prime satisfying

$$(35) \quad \text{Frob}_{\ell_2}(M_{s,\xi,\xi^*}/\mathbb{Q}) = (v, w, w^*, \tau, T).$$

There are infinitely many such primes  $\ell_2$ , by the Chebotarev density theorem. Now, note that

1. By the same reasoning as in the proof of Theorem 3.2, the prime  $\ell_2$  is  $n$ -admissible and  $v_{\ell_2}(s) \neq 0$ .
2. A similar argument shows that  $v_{\ell_2}(\xi) \neq 0$  and  $v_{\ell_2}(\xi^*) \neq 0$ . From this it follows, by part 1 of Proposition 3.8, that  $\text{Sel}_{\ell_1\ell_2}(\mathbb{Q}, W_f) = 0$ . Hence  $(\ell_1, \ell_2)$  is a rigid pair, and Theorem 3.10 follows.

*Case 2.* Suppose that  $\xi$  does not belong to  $\text{Sel}_{\ell_1}(\mathbb{Q}, W_f)$ , so that  $\xi^*$  also does not belong to  $\text{Sel}_{\ell_1}(\mathbb{Q}, W_f^*)$ . Keeping the notations of Case 1, let  $\ell_2$  be a rational prime satisfying

$$(36) \quad \text{Frob}_{\ell_2}(M_{s,\xi,\xi^*}/\mathbb{Q}) = (v, 0, 0, \tau, T).$$

There are infinitely many such primes  $\ell_2$ , by the Chebotarev density theorem. Note that the prime  $\ell_2$  is  $n$ -admissible and  $v_{\ell_2}(s) \neq 0$ , and that  $v_{\ell_2}(\xi)$  and  $v_{\ell_2}(\xi^*)$  both vanish. It follows from part 2 of Proposition 3.8 that either  $\text{Sel}_{\ell_1\ell_2}(\mathbb{Q}, W_f)$  is trivial – i.e.,  $(\ell_1, \ell_2)$  is a rigid pair – or that  $\text{Sel}_{\ell_2}(\mathbb{Q}, W_f)$  is one-dimensional.

**THEOREM 3.11.** *Suppose that  $f$  is a  $p$ -isolated eigenform in  $S_2(\mathcal{T}/\Gamma)$ , and let  $\ell_1$  be an admissible prime for  $f$ . Let  $s$  be a nonzero class in  $H^1(K, A_{f,1})$ . Suppose further that  $\text{Sel}_{\ell_1}(\mathbb{Q}, W_f)$  is one-dimensional over  $\mathbb{F}_p$ . Then there exist infinitely many  $n$ -admissible primes  $\ell_2$  such that*

- (1)  $\partial_{\ell_2}(s) = 0$  and  $v_{\ell_2}(s) \neq 0$ .
- (2)  $(\ell_1, \ell_2)$  is a rigid pair.

*Proof.* This follows directly from the analysis of Case 1 in the proof of Theorem 3.10.

*Congruences between modular forms.* Let  $\ell_1, \ell_2$  be distinct  $n$ -admissible primes relative to  $f$ , such that  $p^n$  divides  $\ell_1 + 1 - \varepsilon_1 a_{\ell_1}(f)$  and  $\ell_2 + 1 - \varepsilon_2 a_{\ell_2}(f)$ , for  $\varepsilon_1$  and  $\varepsilon_2$  equal to  $\pm 1$ . Let  $B'$  be the definite quaternion algebra of discriminant  $\text{Disc}(B)_{\ell_1\ell_2}$ , let  $R'$  be an Eichler  $\mathbb{Z}[1/p]$ -order of level  $N^+$  in  $B'$  and let  $\Gamma' := (R')^\times / \mathbb{Z}[1/p]^\times$ . The theory of congruences between modular forms yields the following proposition:

PROPOSITION 3.12. *There exists an eigenform  $g \in S_2(\mathcal{T}/\Gamma', \mathbb{Z}/p^n\mathbb{Z})$  such that the equalities modulo  $p^n$  hold:*

$$(37) \quad T_q g \equiv a_q(f)g \quad (q \nmid N\ell_1\ell_2), \quad U_q g \equiv a_q(f)g \quad (q|N), \\ U_{\ell_1} g \equiv \varepsilon_1 g, \quad U_{\ell_2} g \equiv \varepsilon_2 g.$$

*If furthermore the pair  $(\ell_1, \ell_2)$  is a rigid pair, then  $g$  can be lifted to an eigenform with coefficients in  $\mathbb{Z}_p$  satisfying (37) above. This form is  $p$ -isolated.*

*Proof.* The existence of the mod  $p^n$  eigenform  $g$ , which relies on the concepts and notation introduced in Sections 5 and 9, is given in Theorem 9.3. This  $g$  corresponds to a surjective algebra homomorphism  $f_{\ell_2, \ell_1} : \mathbb{T}_{\ell_2, \ell_1} \longrightarrow \mathbb{Z}/p^n\mathbb{Z}$ , where  $\mathbb{T}_{\ell_2, \ell_1}$  is the Hecke algebra defined in the proof of Theorem 9.3. If  $(\ell_1, \ell_2)$  is a rigid pair, this algebra is isomorphic to  $\mathbb{Z}_p$  and therefore  $f_{\ell_2, \ell_1}$  lifts to characteristic 0 so that  $g$  can be lifted (uniquely) to a form in  $S_2(\mathcal{T}/\Gamma')$ .

#### 4. The Euler system argument

4.1. *The Euler system.* Section 7 describes the construction of certain global cohomology classes

$$\kappa(\ell) \in \hat{H}_\ell^1(K_\infty, T_{f,n}),$$

indexed by the  $n$ -admissible primes  $\ell$  attached to  $f$ . The proof of Theorem 1 relies crucially on the existence of these classes and on their behaviour under localisation described in Theorems 4.1 and 4.2 below. Both theorems are instances of explicit reciprocity laws relating these explicit cohomology classes to special values of  $L$ -functions, and form the technical heart of the proof of Theorem 1.

Observe that when  $\ell$  is an  $n$ -admissible prime, the local cohomology group  $\hat{H}^1(K_{\infty, \ell}, T_{f,n})$  decomposes as a direct sum

$$\begin{aligned} \hat{H}^1(K_{\infty, \ell}, T_{f,n}) &= \hat{H}_{\text{fin}}^1(K_{\infty, \ell}, T_{f,n}) \oplus \hat{H}_{\text{ord}}^1(K_{\infty, \ell}, T_{f,n}) \\ &= \hat{H}_{\text{fin}}^1(K_{\infty, \ell}, T_{f,n}) \oplus \hat{H}_{\text{sing}}^1(K_{\infty, \ell}, T_{f,n}). \end{aligned}$$

The map  $\partial_\ell$  is simply the projection onto the second factor, while  $v_\ell$  can be extended naturally to a map

$$v_\ell : \hat{H}^1(K_{\infty, \ell}, T_{f,n}) \longrightarrow \hat{H}_{\text{fin}}^1(K_{\infty, \ell}, T_{f,n}) = \hat{H}^1(K_{\infty, \ell}, T_{f,n}) / \hat{H}_{\text{ord}}^1(K_{\infty, \ell}, T_{f,n})$$

defined as the projection onto the first factor.

THEOREM 4.1. *If  $\ell$  is an  $n$ -admissible prime, then  $v_\ell(\kappa(\ell)) = 0$ . The equality*

$$\partial_\ell(\kappa(\ell)) = \mathcal{L}_f \pmod{p^n}$$

*holds in  $\hat{H}_{\text{sing}}^1(K_{\infty, \ell}, T_{f,n}) \simeq \Lambda/p^n\Lambda$ , up to multiplication by elements of  $\mathbb{Z}_p^\times$  and  $G_\infty$ .*

Note that the ambiguity in the statement of Theorem 4.1 is unavoidable, since the identification of  $H_{\text{sing}}^1(K_{\infty,\ell}, T_{f,n})$  with  $\Lambda/p^n\Lambda$ , and the element  $\mathcal{L}_f$ , are both only defined up to multiplication by elements in  $\mathbb{Z}_p^\times$  and  $G_\infty$ .

Theorem 4.1 is proved in Section 8.

The second theorem describes the localisation of  $\kappa(\ell_1)$  at an  $n$ -admissible prime  $\ell_2$  which is different from  $\ell_1$ . Recall the discrete subgroup  $\Gamma'$  of  $\mathbf{PSL}_2(\mathbb{Q}_p)$  and the  $\mathbb{Z}/p^n\mathbb{Z}$ -valued eigenform  $g$  in  $S_2(\mathcal{T}/\Gamma', \mathbb{Z}/p^n\mathbb{Z})$  attached to  $f$  and  $(\ell_1, \ell_2)$  in Proposition 3.12.

**THEOREM 4.2.** *The equality*

$$v_{\ell_2}(\kappa(\ell_1)) = \mathcal{L}_g$$

holds in  $\hat{H}_{\text{fin}}^1(K_{\infty,\ell_2}, T_{f,n}) \simeq \Lambda/p^n\Lambda$ , up to multiplication by elements of  $\mathbb{Z}_p^\times$  and  $G_\infty$ .

Theorem 4.2 is proved in Section 9.

Since the definition of  $g$  is symmetric in  $\ell_1$  and  $\ell_2$ , one obtains the following reciprocity formula for the classes  $\kappa(\ell)$ :

**COROLLARY 4.3.** *For all pairs of  $n$ -admissible primes  $\ell_1, \ell_2$  attached to  $f$ , the equality*

$$v_{\ell_1}(\kappa(\ell_2)) = v_{\ell_2}(\kappa(\ell_1))$$

holds in  $\Lambda/p^n\Lambda$ , up to multiplication by elements of  $\mathbb{Z}_p^\times$  and  $G_\infty$ .

**4.2. The argument.** To an ordinary eigenform  $f \in S_2(\mathcal{T}/\Gamma)$  with coefficients in  $\mathbb{Z}_p$  one has associated two invariants: the  $p$ -adic  $L$ -function  $L_p(f, K) \in \Lambda$  (Section 1) and the Selmer group  $\text{Sel}_{f,n}$  (Section 2). This section explains the proof of Theorem 1. In our approach based on congruences between modular forms, it is indispensable to prove the following generalisation which is stronger insofar as it applies to all  $p$ -isolated modular eigenforms in  $S_2(\mathcal{T}/\Gamma)$  with coefficients in  $\mathbb{Z}_p$  satisfying Assumption 2.1.

**THEOREM 4.4.** *Let  $f$  be an ordinary eigenform in  $S_2(\mathcal{T}/\Gamma)$  with coefficients in  $\mathbb{Z}_p$  which is  $p$ -isolated, and satisfies Assumption 2.1. The characteristic power series of  $\text{Sel}_{f,\infty}^\vee$  divides the  $p$ -adic  $L$ -function  $L_p(f, K)$ .*

*Proof.* By Proposition 3.1, it suffices to show that

$$(38) \quad \varphi(\mathcal{L}_f)^2 \text{ belongs to } \text{Fitt}_{\mathcal{O}}(\text{Sel}_{f,\infty}^\vee \otimes_{\varphi} \mathcal{O}),$$

for all homomorphisms  $\varphi$  of  $\Lambda$  into a discrete valuation ring  $\mathcal{O}$ . For this it is enough to show that

$$(39) \quad \varphi(\mathcal{L}_f)^2 \text{ belongs to } \text{Fitt}_{\mathcal{O}}(\text{Sel}_{f,n}^\vee \otimes_{\varphi} \mathcal{O}), \quad \text{for all } n \geq 1.$$

Fix  $\mathcal{O}$ ,  $\varphi$ , and  $n$ . Write  $\pi$  for a uniformiser of  $\mathcal{O}$ , and let  $e := \text{ord}_\pi(p)$  be the ramification degree of  $\mathcal{O}$  over  $\mathbb{Z}_p$ . Write

$$t_f := \text{ord}_\pi(\varphi(\mathcal{L}_f)).$$

Assume without loss of generality that

1.  $t_f < \infty$ . (Otherwise,  $\varphi(\mathcal{L}_f) = 0$  and (39) is trivially verified.)
2. The group  $\text{Sel}_{f,n}^\vee \otimes \mathcal{O}$  is nontrivial. (Otherwise, its Fitting ideal is equal to  $\mathcal{O}$  and (39) is trivially verified.)

Theorem 4.4 (or rather, equation (39)) is proved by induction on  $t_f$ . We begin by describing the construction of certain cohomology classes attached to an admissible prime  $\ell$ . Let  $\ell$  be any  $(n + t_f)$ -admissible prime, and enlarge  $\{\ell\}$  to an  $(n + t_f)$ -admissible set  $S$ . Let

$$\kappa(\ell) \in \hat{H}_\ell^1(K_\infty, T_{f,n+t_f}) \subset \hat{H}_S^1(K_\infty, T_{f,n+t_f})$$

be the cohomology class attached to  $\ell$  as in Section 4.1, and denote by  $\kappa_\varphi(\ell)$  the natural image of this class in

$$\mathcal{M} := \hat{H}_S^1(K_\infty, T_{f,n+t_f}) \otimes_\varphi \mathcal{O}.$$

Note that this module is free over  $\mathcal{O}/p^{(n+t_f)}$ , by Proposition 3.3. By Theorem 4.1,

$$\text{ord}_\pi(\kappa_\varphi(\ell)) \leq \text{ord}_\pi(\partial_\ell \kappa_\varphi(\ell)) = \text{ord}_\pi(\varphi(\mathcal{L}_f)) = t_f,$$

so that  $t := \text{ord}_\pi(\kappa_\varphi(\ell)) \leq t_f$ . Choose an element  $\tilde{\kappa}_\varphi(\ell) \in \mathcal{M}$  satisfying

$$\pi^t \tilde{\kappa}_\varphi(\ell) = \kappa_\varphi(\ell).$$

Now,  $\tilde{\kappa}_\varphi(\ell)$  is well defined modulo the  $\pi^t$ -torsion subgroup of  $\mathcal{M}$ , which is contained in the kernel of the natural homomorphism

$$\hat{H}_S^1(K_\infty, T_{f,n+t_f}) \otimes_\varphi \mathcal{O} \longrightarrow \hat{H}_S^1(K_\infty, T_{f,n}) \otimes_\varphi \mathcal{O}.$$

To remove this ambiguity, let  $\kappa'_\varphi(\ell)$  be the natural image of the class  $\tilde{\kappa}_\varphi(\ell)$  in  $\hat{H}_S^1(K_\infty, T_{f,n}) \otimes_\varphi \mathcal{O}$ . The key properties of the class  $\kappa'_\varphi(\ell)$  are summarised in Lemmas 4.5 and 4.6 below.

LEMMA 4.5. *The class  $\kappa'_\varphi(\ell)$  enjoys the following properties:*

- (1)  $\text{ord}_\pi(\kappa'_\varphi(\ell)) = 0$ .
- (2)  $\partial_q \kappa'_\varphi(\ell) = 0$ , for all  $q \nmid \ell N^-$ .
- (3)  $v_\ell(\kappa'_\varphi(\ell)) = 0$ .
- (4)  $\text{ord}_\pi(\partial_\ell \kappa'_\varphi(\ell)) = t_f - t$ .

*Proof.* The first property follows from the fact that  $\text{ord}_\pi(\kappa_\varphi(\ell)) = t$ . The second is a direct consequence of the fact that  $\kappa(\ell)$  belongs to  $\hat{H}_\ell^1(K_\infty, T_{f,n+t_f})$ , while the third and fourth follow from Theorem 4.1.

LEMMA 4.6. *The element  $\partial_\ell(\kappa'_\varphi(\ell))$  belongs to the kernel of the natural homomorphism*

$$\eta_\ell : \hat{H}_{\text{sing}}^1(K_{\infty,\ell}, T_{f,n}) \otimes_\varphi \mathcal{O} \longrightarrow \text{Sel}_{f,n}^\vee \otimes_\varphi \mathcal{O}.$$

*Proof.* Let  $I_\varphi$  denote the kernel of  $\varphi$ . By the global reciprocity law of class field theory, the class  $\tilde{\kappa}_\varphi(\ell)$  satisfies

$$(40) \quad \sum_{q|S} \langle \partial_q(\tilde{\kappa}_\varphi(\ell)), s_q \rangle_q = 0,$$

for all  $s \in \text{Sel}_{f,n+t_f}[I_\varphi]$ . (Here,  $s_q$  simply denotes the natural image of  $s$  in  $H_{\text{fin}}^1(K_{q,\infty}, A_{f,n+t_f})$ .) On the other hand,  $\pi^t \tilde{\kappa}_\varphi(\ell) = \kappa_\varphi(\ell)$  has trivial residue at all the primes  $q \neq \ell$ . Hence, for such primes, the element  $\partial_q(\tilde{\kappa}_\varphi(\ell))$  annihilates

$$\pi^t H_{\text{fin}}^1(K_{\infty,q}, A_{f,n+t_f})[I_\varphi] \supset H_{\text{fin}}^1(K_{\infty,q}, A_{f,n})[I_\varphi].$$

Hence, if  $s$  belongs to  $\text{Sel}_{f,n}[I_\varphi]$ , the terms in the sum (40) corresponding to the primes  $q \neq \ell$  are zero. Hence so is the term corresponding to  $\ell$ . It follows that  $\partial_\ell(\kappa'_\varphi(\ell))$  annihilates the image of  $\text{Sel}_{f,n}[I_\varphi]$  in  $H_{\text{fin}}^1(K_{\infty,\ell}, A_{f,n})$ , as was to be shown.

We now turn to the proof of (39), in the case where  $t_f = 0$ , which provides the basis for the induction argument.

PROPOSITION 4.7. *If  $t_f = 0$  (i.e.,  $\mathcal{L}_f$  is a unit), then  $\text{Sel}_{f,n}^\vee$  is trivial.*

*Proof.* Since  $\mathcal{L}_f$  is a unit, Theorem 4.1 implies that  $\partial_\ell(\kappa_\varphi(\ell))$  generates  $\hat{H}_{\text{sing}}^1(K_{\infty,\ell}, T_{f,n}) \otimes_\varphi \mathcal{O}$ , for all  $n$ -admissible primes  $\ell$ . Hence the map  $\eta_\ell$  of Lemma 4.6 is trivial for all such primes. This is enough to conclude that  $\text{Sel}_{f,n}^\vee$  is trivial. For otherwise, Nakayama's lemma implies that the group

$$\text{Sel}_{f,n}^\vee / \mathfrak{m}_\Lambda = (\text{Sel}_{f,n}[\mathfrak{m}_\Lambda])^\vee$$

is nonzero. Let  $s$  be a nontrivial element of  $\text{Sel}_{f,n}[\mathfrak{m}_\Lambda]$ . By part 1 of Theorem 3.4,  $s$  can be viewed as an element of  $H^1(K, A_{f,1})$ . Invoking Theorem 3.2, choose an  $n$ -admissible prime  $\ell$  such that  $v_\ell(s) \neq 0$ . The nondegeneracy of the local Tate pairing implies that  $\eta_\ell$  is nonzero, a contradiction.

Turning now to the general case of equation (39), let  $\Pi$  be the set of rational primes  $\ell$  satisfying the following conditions:

- (1)  $\ell$  is  $(n + t_f)$ -admissible.
- (2) The quantity  $t = \text{ord}_\pi(\kappa_\varphi(\ell))$  is minimal, among all primes satisfying condition 1.

Note that the set  $\Pi$  is nonempty, by Theorem 3.2. Let  $t$  be the common value of  $\text{ord}_\pi(\kappa_\varphi(\ell))$  for  $\ell \in \Pi$ .

LEMMA 4.8.  $t < t_f$ .

*Proof.* Suppose not. Then  $\text{ord}_\pi(\kappa_\varphi(\ell)) = t_f$ , for all  $(n + t_f)$ -admissible primes  $\ell$ . Let  $s$  be a nonzero element of  $H^1(K, A_{f,1}) \cap \text{Sel}_{f,n}$ , which exists by Theorem 3.4. Invoking Theorem 3.2, choose an  $(n + t_f)$ -admissible prime  $\ell$  such that  $v_\ell(s) \neq 0$ . By Lemma 4.5, the natural image of  $\partial_\ell(\kappa'_\varphi(\ell))$  in  $H^1(K_\ell, T_{f,1}) \otimes_\varphi \mathcal{O}$  is nonzero. By Lemma 4.6, it is also orthogonal to  $v_\ell(s)$  with respect to the local Tate pairing, contradicting the fact that the vectors  $\partial_\ell(\kappa'_\varphi(\ell))$  and  $v_\ell(s)$  are both supposed to be nonzero and that the Tate pairing is a perfect duality between these two one-dimensional vector spaces over  $\mathcal{O}/\pi$ .

LEMMA 4.9. *There exist primes  $\ell_1, \ell_2 \in \Pi$  such that  $(\ell_1, \ell_2)$  is a rigid pair.*

*Proof.* Choose any  $\ell_1$  in  $\Pi$ , and let  $s$  denote the natural image of  $\kappa'_\varphi(\ell_1)$  in

$$\begin{aligned} \hat{H}_S^1(K_\infty, T_{f,n}) \otimes_\varphi \mathcal{O}/(\pi) &= \left( \hat{H}_S^1(K_\infty, T_{f,n})/\mathfrak{m}_\Lambda \right) \otimes (\mathcal{O}/\pi) \\ &\subset H^1(K, T_{f,1}) \otimes (\mathcal{O}/\pi), \end{aligned}$$

where the last inclusion (cf. part 2 of Theorem 3.4) is induced from the corestriction map and the natural projection  $T_{f,n} \rightarrow T_{f,1}$ . Observe that the class  $s$  is a nonzero element of  $H^1(K, T_{f,1}) \otimes (\mathcal{O}/\pi)$  which satisfies  $\partial_q(s) = 0$  for all  $q \nmid \ell_1 N$ . Invoking Theorem 3.10, choose an  $n + t_f$ -admissible prime  $\ell_2$  such that

1.  $v_{\ell_2}(s) \neq 0$ , and
2. either  $(\ell_1, \ell_2)$  is a rigid pair, or  $\text{Sel}_{\ell_2}(\mathbb{Q}, W_f)$  is one-dimensional.

The reader will note that:

$$(41) \quad t = \text{ord}_\pi(\kappa_\varphi(\ell_1)) \leq \text{ord}_\pi(\kappa_\varphi(\ell_2)) \leq \text{ord}_\pi(v_{\ell_1}(\kappa_\varphi(\ell_2))).$$

The first inequality holds by the minimality assumption made in the choice of the prime  $\ell_1$ . The second inequality is a consequence of the fact that  $v_{\ell_1}$  is a homomorphism. By the reciprocity law of Corollary 4.3, and the choice of  $\ell_2$ ,

$$(42) \quad \text{ord}_\pi(v_{\ell_1}(\kappa_\varphi(\ell_2))) = \text{ord}_\pi(v_{\ell_2}(\kappa_\varphi(\ell_1))) = \text{ord}_\pi(\kappa_\varphi(\ell_1)).$$

(To see the second equality, note that the inequality

$$\text{ord}_\pi(v_{\ell_2}(\kappa_\varphi(\ell_1))) \geq \text{ord}_\pi(\kappa_\varphi(\ell_1))$$

is clear, and that strict inequality holds precisely when  $v_{\ell_2}(s) = 0$ .) Combining (41) and (42), it follows that the inequalities must be equalities throughout, so that

$$t = \text{ord}_\pi(\kappa_\varphi(\ell_1)) = \text{ord}_\pi(\kappa_\varphi(\ell_2)).$$

Hence  $\ell_2$  belongs to  $\Pi$ . If  $(\ell_1, \ell_2)$  is a rigid pair, we are done. Otherwise, the group  $\text{Sel}_{\ell_2}(\mathbb{Q}, W_f)$  is one-dimensional. In that case one can repeat the argument above, with  $\ell_1$  replaced by  $\ell_2$ , invoking this time Theorem 3.11 instead of 3.10 to obtain a pair  $(\ell_2, \ell_3)$  satisfying the conclusion of Lemma 4.9. This completes the proof of the lemma.

Let  $(\ell_1, \ell_2)$  be a rigid pair of  $(n + t_f)$ -admissible primes in  $\Pi$ , whose existence is guaranteed by Lemma 4.9. By Theorem 4.2, note that  $t = t_g = \text{ord}_\pi(\varphi(\mathcal{L}_g))$ , where  $g$  is the  $p$ -isolated eigenform in  $S_2(\mathcal{T}/\Gamma')$  attached to  $f$  and  $(\ell_1, \ell_2)$  by Proposition 3.12.

Let  $\text{Sel}_{[\ell_1, \ell_2]}$  denote the subgroup of  $\text{Sel}_{f, n}$  consisting of classes which are locally trivial at the primes dividing  $\ell_1$  and  $\ell_2$ . By definition, there is a natural exact sequence of  $\Lambda$ -modules

$$(43) \quad 0 \longrightarrow S_{\ell_1 \ell_2}^f \longrightarrow \text{Sel}_{f, n}^\vee \longrightarrow \text{Sel}_{[\ell_1, \ell_2]}^\vee \longrightarrow 0,$$

where  $S_{\ell_1 \ell_2}^f$  denotes the kernel of the natural surjection of duals of Selmer groups. Note the natural surjection given by local Tate duality:

$$\eta_f : (\hat{H}_{\text{sing}}^1(K_{\infty, \ell_1}, A_{f, n}) \oplus \hat{H}_{\text{sing}}^1(K_{\infty, \ell_2}, A_{f, n})) \longrightarrow S_{\ell_1 \ell_2}^f$$

induced from the inclusion

$$(S_{\ell_1 \ell_2}^f)^\vee \subset H_{\text{fin}}^1(K_{\infty, \ell_1}, A_{f, n}) \oplus H_{\text{fin}}^1(K_{\infty, \ell_2}, A_{f, n}).$$

The domain of  $\eta_f$  is isomorphic to  $(\Lambda/p^n \Lambda)^2$ , by Lemma 2.7. Let  $\eta_f^\varphi$  denote the map induced from  $\eta_f$  after tensoring by  $\mathcal{O}$  via  $\varphi$ . The domain of  $\eta_f^\varphi$  is isomorphic to  $(\mathcal{O}/p^n \mathcal{O})^2$ . By Lemma 4.6, the kernel of  $\eta_f^\varphi$  contains the vectors  $(\partial_{\ell_1} \kappa'_\varphi(\ell_1), 0)$  and  $(0, \partial_{\ell_2} \kappa'_\varphi(\ell_2))$  in

$$\left( \hat{H}_{\text{sing}}^1(K_{\infty, \ell_1}, A_{f, n}) \oplus \hat{H}_{\text{sing}}^1(K_{\infty, \ell_2}, A_{f, n}) \right) \otimes_\varphi \mathcal{O} \simeq (\mathcal{O}/p^n \mathcal{O})^2.$$

By part 3 of Lemma 4.5,

$$t_f - t_g = \text{ord}_\pi(\partial_{\ell_1} \kappa'_\varphi(\ell_1)) = \text{ord}_\pi(\partial_{\ell_2} \kappa'_\varphi(\ell_2)).$$

Hence

$$(44) \quad \pi^{2(t_f - t_g)} \text{ belongs to the Fitting ideal of } S_{\ell_1 \ell_2}^f \otimes_\varphi \mathcal{O}.$$

One may repeat the above argument with the modular form  $g$ . Thus we have an exact sequence similar to (43) but involving  $g$  instead of  $f$ :

$$(45) \quad 0 \longrightarrow S_{\ell_1 \ell_2}^g \longrightarrow \text{Sel}_{g, n}^\vee \longrightarrow \text{Sel}_{[\ell_1, \ell_2]}^\vee \longrightarrow 0,$$

as well as a surjection given by local Tate duality:

$$\eta_g : (\hat{H}_{\text{fin}}^1(K_{\infty, \ell_1}, A_{f, n}) \oplus \hat{H}_{\text{fin}}^1(K_{\infty, \ell_2}, A_{f, n})) \longrightarrow S_{\ell_1 \ell_2}^g.$$

By global reciprocity, the kernel of the map  $\eta_g^\varphi$  obtained from  $\eta_g$  after tensoring by  $\mathcal{O}$  via  $\varphi$  contains the elements

$$(v_{\ell_1} \kappa'_\varphi(\ell_1), v_{\ell_2} \kappa'_\varphi(\ell_1)) = (0, v_{\ell_2} \kappa'_\varphi(\ell_1))$$

as well as  $(v_{\ell_1} \kappa'_\varphi(\ell_2), 0)$ . But

$$\text{ord}_\pi(v_{\ell_2} \kappa'_\varphi(\ell_1)) = \text{ord}_\pi(v_{\ell_1} \kappa'_\varphi(\ell_2)) = t_g - t = 0.$$

It follows from this that the module  $S_g \otimes_\varphi \mathcal{O}$  is trivial, and the natural surjection

$$(46) \quad \text{Sel}_{g, n}^\vee \otimes_\varphi \mathcal{O} \longrightarrow \text{Sel}_{[\ell_1 \ell_2]}^\vee \otimes_\varphi \mathcal{O} \text{ is an isomorphism.}$$

Recall that, by Lemma 4.8,

$$t_g < t_f,$$

and that the eigenform  $g$  satisfies all the hypotheses of Theorem 4.4, including Assumption 2.1, in light of the fact that  $\ell_1$  and  $\ell_2$  are admissible. One is thus in a position to invoke the induction hypothesis to conclude that

$$(47) \quad \varphi(\mathcal{L}_g)^2 \text{ belongs to the Fitting ideal of } \text{Sel}_{g, n}^\vee \otimes_\varphi \mathcal{O}.$$

The theory of Fitting ideals implies that

$$\begin{aligned} \pi^{2t_f} &= \pi^{2(t_f - t_g)} \pi^{2t_g} \\ &\in \text{Fitt}_{\mathcal{O}}(S_{\ell_1 \ell_2}^f \otimes \mathcal{O}) \text{Fitt}_{\mathcal{O}}(\text{Sel}_{g, n}^\vee \otimes \mathcal{O}), \quad \text{by (44) and (47)} \\ &= \text{Fitt}_{\mathcal{O}}(S_{\ell_1 \ell_2}^f \otimes \mathcal{O}) \text{Fitt}_{\mathcal{O}}(\text{Sel}_{[\ell_1 \ell_2]}^\vee \otimes \mathcal{O}), \quad \text{by (46)} \\ &\subset \text{Fitt}_{\mathcal{O}}(\text{Sel}_{f, n}^\vee \otimes \mathcal{O}), \quad \text{by (43)}. \end{aligned}$$

Hence (39) is proved:  $\varphi(\mathcal{L}_f)^2$  belongs to the Fitting ideal of  $\text{Sel}_{f, n}^\vee \otimes_\varphi \mathcal{O}$ . Theorem 4.4 follows.

## 5. Shimura curves

The construction and the properties of the Euler system used in the argument of Section 4 are based on the geometry over  $\mathbb{Z}$  of certain Shimura curves, which are reviewed in this section.

Let  $M$  be a positive integer, and let  $M = M^+ M^-$  be an integer decomposition of  $M$  such that  $M^-$  is a squarefree product (possibly empty) of an even number of prime factors. Following [Ro] and [BD1], one may attach to such a decomposition a Shimura curve  $X_{M^+, M^-}$ . If  $M^-$  is equal to 1,  $X_{M^+, M^-}$  is the classical modular curve  $X_0(M)$  of level  $M$ . In the general case,  $X_{M^+, M^-}$  is the Shimura curve with level  $M^+$ -structure associated to the indefinite quaternion algebra of discriminant  $M^-$ .

5.1. *The moduli definition.* The curve  $X = X_{M^+, M^-}$  has the following moduli interpretations.

*Models over  $\mathbb{Z}[\frac{1}{M}]$ .* Let  $\mathcal{B}$  be the indefinite quaternion algebra over  $\mathbb{Q}$  of discriminant  $M^-$ . Fix a maximal order  $\mathcal{R}_{\max}$  in  $\mathcal{B}$ , and an Eichler order  $\mathcal{R}$  of level  $M^+$  contained in  $\mathcal{R}_{\max}$ . Write  $\mathcal{F} = \mathcal{F}_{\mathbb{Z}[\frac{1}{M}]}$  for the functor from the category of schemes over  $\mathbb{Z}[\frac{1}{M}]$  to the category of sets which associates to a scheme  $S$  the set of isomorphism classes of triples  $(A, \iota, C)$ , where:

- (1)  $A$  is an abelian scheme over  $S$  of relative dimension 2,
- (2)  $\iota : \mathcal{R}_{\max} \rightarrow \text{End}(A)$  is an action of  $\mathcal{R}_{\max}$  on  $A$ ,
- (3)  $C$  is a level  $M^+$ -structure on  $A$ , that is, a subgroup scheme of  $A$  of order  $(M^+)^2$  which is stable and cyclic for the action of  $\mathcal{R}_{\max}$ .

If  $M^-$  is strictly greater than 1, the functor  $\mathcal{F}$  is coarsely representable by a smooth projective scheme  $X_{\mathbb{Z}[\frac{1}{M}]}$  over  $\mathbb{Z}[\frac{1}{M}]$ , with smooth fibers. Let  $\mathcal{H}_{\infty}$  be the complex upper half-plane, and let  $\mathcal{R}_1^{\times}$  be the group of norm 1 elements in  $\mathcal{R}$ . Fix an embedding of  $\mathcal{B}$  in  $M_2(\mathbb{R})$ , and write  $\Gamma_{\infty, 1}$  for the natural image of  $\mathcal{R}_1^{\times}$  in  $\mathbf{PGL}_2(\mathbb{R})$ . The group  $\Gamma_{\infty, 1}$  acts discontinuously (on the right) on  $\mathcal{H}_{\infty}$ , and the complex points  $X(\mathbb{C})$  of the generic fiber  $X = X_{\mathbb{Q}}$  are identified with the Riemann surface  $\mathcal{H}_{\infty}/\Gamma_{\infty, 1}$ . For more information, see [BC, Ch. III] and [Bu].

If  $M^-$  is equal to 1, then  $\mathcal{B}$  is isomorphic to the split quaternion algebra  $M_2(\mathbb{Q})$ , and one may assume that  $\mathcal{R}_{\max}$  corresponds to the standard maximal order  $M_2(\mathbb{Z})$ . A triple  $(A, i, C)$  as above is then of the form  $(E \times E, \iota, C_E \times C_E)$  where  $E$  is an elliptic curve,  $C_E$  is a level  $M$ -structure on  $E$ , and the action  $\iota$  is the natural matrix action of  $M_2(\mathbb{Z})$  on  $E \times E$ . Thus, the functor  $\mathcal{F}$  is coarsely representable by the affine modular curve  $Y_0(M)_{\mathbb{Z}[\frac{1}{M}]}$ . The projective completion  $X_0(M)_{\mathbb{Z}[\frac{1}{M}]}$  of  $Y_0(M)_{\mathbb{Z}[\frac{1}{M}]}$ , obtained by adding a finite set of cusps, is a moduli space for generalized elliptic curves with level  $M$ -structure. See [DR] and [Bu].

*Models over  $\mathbb{Z}_{\ell}$  for  $\ell \mid M^+$ .* Assume that  $\ell$  is a prime dividing exactly  $M^+$ . (This is the only case which is relevant to the arguments of this paper.)

The reader is referred to [Ed, §§3 and 4], [DR], [KM], and [Bu] for the definition of the variant of the moduli functor  $\mathcal{F}$  which can be used in this case. The resulting canonical model  $X_{\mathbb{Z}_{\ell}}$  is a *nodal model* of  $X$ , in the sense of [Ed]. That is:

- (1)  $X_{\mathbb{Z}_{\ell}}$  is proper and flat over  $\mathbb{Z}_{\ell}$ , and its generic fiber is  $X$  (viewed as a curve over  $\mathbb{Q}_{\ell}$ ),
- (2) the irreducible components of the special fiber  $X_{\mathbb{F}_{\ell}}$  of  $X_{\mathbb{Z}_{\ell}}$  are smooth, and the only singularities of  $X_{\mathbb{F}_{\ell}}$  are ordinary double points.

More precisely, the special fiber  $X_{\mathbb{F}_\ell}$  consists of two copies of the irreducible curve  $(X_{M^+/\ell, M^-})_{\mathbb{F}_\ell}$  intersecting transversally at the supersingular points, which are identified via the Frobenius morphism at  $\ell$ . (Note that  $X_{M^+/\ell, M^-}$  has good reduction at  $\ell$ , and a description of  $(X_{M^+/\ell, M^-})_{\mathbb{F}_\ell}$  follows from the model  $(X_{M^+/\ell, M^-})_{\mathbb{Z}[\frac{1}{M/\ell}]}$  given above.)

*Models over  $\mathbb{Z}_\ell$  for  $\ell \mid M^-$ .* Assume that  $M^-$  is strictly greater than 1. Fix a prime  $\ell$  dividing  $M^-$ . As before, one may define a model  $X_{\mathbb{Z}_\ell}$  of  $X$  over  $\mathbb{Z}_\ell$  via moduli. The new moduli functor  $\mathcal{F}_{\mathbb{Z}_\ell}$  on schemes over  $\mathbb{Z}_\ell$  is defined similarly to  $\mathcal{F}_{\mathbb{Z}[\frac{1}{M}]}$ , except for the requirement that the action  $\iota$  be *special* in the sense of [BC, §III.3]. The functor  $\mathcal{F}_{\mathbb{Z}_\ell}$  is coarsely representable by a scheme  $X_{\mathbb{Z}_\ell}$ , which is a nodal model of  $X$ . A more precise description of  $X_{\mathbb{Z}_\ell}$  is given in Section 5.2 where, in particular, it is explained that the irreducible components of  $X_{\mathbb{F}_\ell}$  are rational curves.

5.2. *The Cerednik-Drinfeld theorem.* (References: see [JoLi1] and [BC] for details, and [BD3, §4] for an exposition.) Assume that  $M^-$  is strictly greater than 1, and let  $\ell$  be a prime dividing  $M^-$ . Let  $B$  be the definite quaternion algebra over  $\mathbb{Q}$  of discriminant  $M^-/\ell$ , and let  $R$  be an Eichler order in  $B$  of level  $M^+\ell$ . Set  $B_\ell = B \otimes \mathbb{Z}_\ell$  and  $R_\ell = R \otimes \mathbb{Z}_\ell$ , and fix an isomorphism  $\kappa_\ell : B_\ell \xrightarrow{\sim} M_2(\mathbb{Q}_\ell)$  mapping  $R_\ell$  onto the standard Eichler order of level  $\ell$  in  $M_2(\mathbb{Z}_\ell)$ , consisting of matrices which are upper triangular modulo  $\ell$ . Write  $\Gamma_\ell$ , respectively,  $\Gamma_{\ell,1}$  for the image of  $R[\frac{1}{\ell}]^\times$ , respectively,  $R[\frac{1}{\ell}]_1^\times$  in  $\mathbf{PGL}_2(\mathbb{Q}_\ell)$ , where  $R[\frac{1}{\ell}]_1^\times$  is the subgroup of norm 1 elements in  $R[\frac{1}{\ell}]^\times$ .

Let  $\mathbb{C}_\ell$  be a completion of an algebraic closure of  $\mathbb{Q}_\ell$ , and let  $\hat{\mathcal{H}}_\ell$  be the  $\ell$ -adic upper half-plane, viewed as a formal scheme over  $\mathbb{Z}_\ell$ . The generic fiber of  $\hat{\mathcal{H}}_\ell$  is identified with a rigid analytic space  $\mathcal{H}_\ell$  over  $\mathbb{Q}_\ell$  whose  $\mathbb{C}_\ell$ -valued points are

$$\mathcal{H}_\ell(\mathbb{C}_\ell) = \mathbb{P}^1(\mathbb{C}_\ell) - \mathbb{P}^1(\mathbb{Q}_\ell) = \mathbb{C}_\ell - \mathbb{Q}_\ell.$$

The special fiber of  $\hat{\mathcal{H}}_\ell$  consists of an infinite sequence of projective lines, intersecting at ordinary double points. The  $\ell$ -adic group  $\Gamma_{\ell,1}$  acts discontinuously on  $\hat{\mathcal{H}}_\ell$  and  $\mathcal{H}_\ell$  on the right. The action on  $\mathcal{H}_\ell(\mathbb{C}_\ell)$  is given by the rule

$$z\gamma = \frac{az + b}{cz + d}, \quad \text{where } \gamma^{-1} \text{ is represented by } \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

The quotients  $\hat{\mathcal{H}}_\ell/\Gamma_{\ell,1}$  and  $\mathcal{H}_\ell/\Gamma_{\ell,1}$  exist in the category of formal schemes and of rigid analytic spaces, respectively. Since the irreducible components of the special fiber of  $\hat{\mathcal{H}}_\ell$  have finite stabilizer in  $\Gamma_{\ell,1}$ , it follows that the irreducible components of the special fiber of  $\mathcal{H}_\ell/\Gamma_{\ell,1}$  are rational curves.

Write  $\hat{X}_{\mathbb{Z}_\ell}$  for the formal completion of  $X_{\mathbb{Z}_\ell}$  along its special fiber, where  $X_{\mathbb{Z}_\ell}$  is the model of  $X$  over  $\mathbb{Z}_\ell$  introduced in Section 5.1. Let  $X^{\text{an}}$  be the rigid analytic space over  $\mathbb{Q}_\ell$  associated to  $X$ . Fix a quadratic unramified extension

$\mathbb{Q}_{\ell^2}$  of  $\mathbb{Q}_{\ell}$ , and denote by  $\mathbb{Z}_{\ell^2}$  the ring of integers of  $\mathbb{Q}_{\ell^2}$  and by  $\mathbb{F}_{\ell^2}$  its residue field.

**THEOREM 5.1.** *The formal schemes  $\hat{X}_{\mathbb{Z}_{\ell}}$  and  $\hat{\mathcal{H}}_{\ell}/\Gamma_{\ell,1}$  are naturally isomorphic over  $\mathbb{Z}_{\ell^2}$ . In particular,  $X^{\text{an}}$  is isomorphic to  $\mathcal{H}_{\ell}/\Gamma_{\ell,1}$  over  $\mathbb{Q}_{\ell^2}$ , and  $X_{\mathbb{F}_{\ell}}$  is isomorphic to the special fiber of  $\hat{\mathcal{H}}_{\ell}/\Gamma_{\ell,1}$  over  $\mathbb{F}_{\ell^2}$ .*

Theorem 5.1 can be proved by comparing the moduli definition of  $X_{\mathbb{Z}_{\ell}}$  given in Section 5.1 with Drinfeld's interpretation [Dr] of  $\hat{\mathcal{H}}_{\ell}$ , where the base-change of  $\hat{\mathcal{H}}_{\ell}$  to the maximal unramified extension of  $\mathbb{Z}_{\ell}$  is identified with the classifying space of certain formal groups of dimension 2 and height 4 which are endowed with an action of the local order  $\mathcal{R}_{\text{max}} \otimes \mathbb{Z}_{\ell}$ . To obtain the version of the Cerednik-Drinfeld theorem given here, one must descend the isomorphism obtained from the above comparison to  $\mathbb{Z}_{\ell}$ , using the arguments of Jordan and Livné in [JoLi1]; see also [BC, Ch. III].

**5.3. Character groups.** Let  $\ell$  be a prime dividing  $M$  exactly, and let  $X_{\mathbb{Z}_{\ell^2}}$  be a fixed nodal model of  $X$  over the unramified quadratic extension  $\mathbb{Z}_{\ell^2}$  of  $\mathbb{Z}_{\ell}$ . (It is convenient to extend scalars from  $\mathbb{Z}_{\ell}$  to  $\mathbb{Z}_{\ell^2}$ , also in view of the isomorphism of Theorem 5.1.) The *dual graph*  $\mathcal{G}_{\ell} = \mathcal{G}_{\ell}(X)$  of  $X$  at  $\ell$  is defined to be the finite graph determined by the following properties:

- (1) The set of vertices  $\mathcal{V}(\mathcal{G}_{\ell})$  is the set of irreducible (geometric) components of the special fiber  $X_{\mathbb{F}_{\ell^2}}$ .
- (2) The set of (unoriented) edges  $\mathcal{E}(\mathcal{G}_{\ell})$  is the set of singular points of  $X_{\mathbb{F}_{\ell^2}}$ .
- (3) Two vertices  $v$  and  $v'$  are joined by an edge  $e$  if  $v$  and  $v'$  intersect at the singular point  $e$ .

Let  $xy = \ell^{\nu_e}$  be a local equation for  $X_{\mathbb{Z}_{\ell^2}}$  at the double point  $e$ . The assignment  $e \mapsto \nu_e$  equips the graph  $\mathcal{G}_{\ell}$  with a weight function  $\nu : \mathcal{E}(\mathcal{G}_{\ell}) \rightarrow \mathbb{N}$ .

Fix an orientation on  $\mathcal{G}_{\ell}$ , that is, a pair of maps  $s, t : \mathcal{E}(\mathcal{G}_{\ell}) \rightarrow \mathcal{V}(\mathcal{G}_{\ell})$  such that  $s(e)$  and  $t(e)$  are the ends of the edge  $e$ , called the source and the target of  $e$ , respectively.

Let  $\mathbb{Z}[\mathcal{V}(\mathcal{G}_{\ell})]$  and  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_{\ell})]$  be the module of formal divisors with integer coefficients supported on  $\mathcal{V}(\mathcal{G}_{\ell})$  and  $\mathcal{E}(\mathcal{G}_{\ell})$ , respectively. Let  $\mathbb{Z}[\mathcal{V}(\mathcal{G}_{\ell})]^0$  and  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_{\ell})]^0$  be the submodule of degree zero divisors in  $\mathbb{Z}[\mathcal{V}(\mathcal{G}_{\ell})]$  and  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_{\ell})]$ , respectively. Let

$$\partial_* : \mathbb{Z}[\mathcal{E}(\mathcal{G}_{\ell})] \rightarrow \mathbb{Z}[\mathcal{V}(\mathcal{G}_{\ell})]^0$$

be the  $\mathbb{Z}$ -linear boundary map defined by the rule  $\partial_*(e) = t(e) - s(e)$ .

Let  $J = J(X)$  be the jacobian of  $X$ , and let  $J_{\mathbb{Z}_{\ell^2}}$  be the Néron model of  $J$  over  $\mathbb{Z}_{\ell^2}$ . Write  $J_{\mathbb{F}_{\ell^2}}$  for the special fiber of  $J$ ,  $J_{\mathbb{F}_{\ell^2}}^0$  for the connected component of the origin in  $J_{\mathbb{F}_{\ell^2}}$ , and  $\Phi_{\ell} = \Phi_{\ell}(X)$  for the group of connected

(geometric) components  $J_{\mathbb{F}_{\ell^2}}/J_{\mathbb{F}_{\ell^2}}^0$ . Denote by  $\mathcal{X}_\ell = \mathcal{X}_\ell(X)$  the *character group*  $\text{Hom}(T_{\mathbb{F}_{\ell^2}}, \mathbb{G}_m)$  of  $J$  at  $\ell$ , where  $T_{\mathbb{F}_{\ell^2}}$  is the maximal torus of  $J_{\mathbb{F}_{\ell^2}}^0$ .

PROPOSITION 5.2. *The module  $\mathcal{X}_\ell$  is isomorphic to the kernel of  $\partial_*$ , and hence fits into the exact sequence*

$$0 \rightarrow \mathcal{X}_\ell \rightarrow \mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)] \xrightarrow{\partial_*} \mathbb{Z}[\mathcal{V}(\mathcal{G}_\ell)]^0 \rightarrow 0.$$

*Sketch of proof.* The module  $\mathcal{X}_\ell$  is canonically identified with the integral homology group  $H_1(\mathcal{G}_\ell, \mathbb{Z})$ . Since the kernel of the map  $\partial_*$  (which depends on the choice of orientation on  $\mathcal{G}_\ell$ ) computes  $H_1(\mathcal{G}_\ell, \mathbb{Z})$ , the proposition follows. (For more details, see [Ed, §1].)

*The character group  $\mathcal{X}_\ell$  for  $\ell \mid M^+$ .* Let  $X_{\mathbb{Z}_{\ell^2}}$  be (the base change of) the model of  $X$  at a prime  $\ell$  dividing exactly  $M^+$ , introduced in Section 5.1. Let  $\mathcal{S}_\ell = \mathcal{S}_\ell(X)$  be the set of supersingular points of the special fiber  $X_{\mathbb{F}_{\ell^2}}$ . By the facts recalled in Section 5.1,  $\mathcal{S}_\ell$  is equal to  $\mathcal{E}(\mathcal{G}_\ell)$ ; furthermore,  $\mathcal{V}(\mathcal{G}_\ell)$  contains two elements, say  $v_1$  and  $v_2$ . Fix an orientation on  $\mathcal{G}_\ell$  so that  $s(e) = v_1$  and  $t(e) = v_2$  for all edges  $e$ .

PROPOSITION 5.3. *The character group  $\mathcal{X}_\ell$  is identified with the group  $\mathbb{Z}[\mathcal{S}_\ell]^0$  of degree zero divisors with  $\mathbb{Z}$ -coefficients supported on  $\mathcal{S}_\ell$ .*

*Proof.* In view of the above remarks, Proposition 5.3 follows directly from Proposition 5.2.

*The character group  $\mathcal{X}_\ell$  for  $\ell \mid M^-$ .* Let  $\mathcal{T}_\ell$  be the Bruhat-Tits tree of  $\mathbf{PGL}_2(\mathbb{Q}_\ell)$ .

PROPOSITION 5.4. *The dual graph  $\mathcal{G}_\ell$  is identified with  $\mathcal{T}_\ell/\Gamma_{\ell,1}$ .*

*Proof.* This follows from Theorem 5.1, by the fact that  $\mathcal{T}_\ell$  is identified with the dual graph of the special fiber of the formal scheme  $\hat{\mathcal{H}}_\ell$  (see [BC, Ch. I]).

From now on, fix the following orientation on  $\mathcal{E}(\mathcal{G}_\ell)$ . Let  $v_0$  be the vertex of  $\mathcal{T}_\ell$  corresponding to the local maximal order  $M_2(\mathbb{Z}_\ell)$ . Say that a vertex of  $\mathcal{T}_\ell$  is *even* or *odd* depending on whether its distance from  $v_0$  is even or odd, respectively. Since the elements of  $\Gamma_{\ell,1}$  have determinant 1, they send even vertices of  $\mathcal{T}_\ell$  to even ones, and odd vertices to odd ones. Thus, there is a well defined notion of even and odd vertex for the quotient graph  $\mathcal{G}_\ell$ . Define the source and target maps  $s, t : \mathcal{E}(\mathcal{G}_\ell) \rightarrow \mathcal{V}(\mathcal{G}_\ell)$  so that  $s(e)$  is the even vertex of  $e$  and  $t(e)$  is the odd vertex of  $e$ .

Write  $\delta_*$  for the restriction to  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0$  of the map  $\partial_*$  (relative to the above choice of orientation).

PROPOSITION 5.5. *The module  $\mathcal{X}_\ell$  fits into the exact sequence*

$$0 \rightarrow \mathcal{X}_\ell \rightarrow \mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0 \xrightarrow{\delta_*} \mathbb{Z}[\mathcal{V}(\mathcal{G}_\ell)]^0.$$

*Proof.* By the choice of orientation made above, the elements of  $H_1(\mathcal{G}_\ell, \mathbb{Z})$  belong to  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0$ . Proposition 5.5 follows directly from Proposition 5.2.

5.4. *Hecke operators and the Jacquet-Langlands correspondence.* (References: see [Ri1], [Ri2] and [JoLi3].) Assume that  $M^-$  is strictly greater than 1, and let  $\ell$  be a prime dividing  $M^-$ . This section is concerned with the study of natural families of Hecke operators acting on the terms of the exact sequence of Proposition 5.5. The natural Hecke algebra acting by Picard functoriality on the jacobian  $J = \text{Pic}^0(X)$  induces an action on the character group  $\mathcal{X}_\ell$ . On the other hand, in order to define an action of Hecke operators on  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0$  and  $\text{Im}(\delta_*)$ , one must use the interpretation of these modules in terms of double coset spaces provided by Lemma 5.6.

*Remark.* A Hecke correspondence  $T$  on  $X$  induces endomorphisms  $T$  and  $\xi$  of  $J$  via Picard (contravariant) and Albanese (covariant) functoriality. The reader is referred to the discussion in [Ri2, pp. 445–6] for details on the definitions. Unless stated otherwise, the Hecke actions considered in this section and in the following ones will be induced from the Hecke action on  $J$  obtained from Picard functoriality. If  $w_{M^+,1}$  denotes the Atkin-Lehner involution defined in [BD1, §1.5], the relation

$$w_{M^+,1} T w_{M^+,1} = \xi$$

holds. (This can be seen by imitating the arguments in Proposition 3.54 of [Sh].) In particular, the Hecke operators corresponding to the primes which do not divide  $M^+$  induce the same endomorphism via Picard and Albanese functoriality. (For the primes dividing  $M^-$ , this can also be checked by observing that the corresponding Hecke operators are involutions; in this case, a general property of curves shows that the two functorialities induce the same endomorphism.) In view of the above remarks, the detailed discussion of Hecke actions contained in Chapters 3 and 4 of [Ri2] extends to the more general setting of this paper.

Let  $B$ ,  $R$  and  $\kappa_\ell$  be as in Section 5.2, and let  $\underline{R}$  be the Eichler order of level  $M^+$  which contains  $R$  and is mapped by  $\kappa_\ell$  to  $M_2(\mathbb{Z}_\ell)$ . Use the notation of formula (12).

LEMMA 5.6. (1) *The set  $\mathcal{E}(\mathcal{G}_\ell)$  is identified with the double coset space  $\hat{R}^\times \backslash \hat{B}^\times / B^\times$ .*

(2) *The set  $\mathcal{V}(\mathcal{G}_\ell)$  is identified with the disjoint union  $(\hat{R}^\times \backslash \hat{B}^\times / B^\times) \times \{0, 1\}$  of the double coset space  $\hat{R}^\times \backslash \hat{B}^\times / B^\times$  with itself.*

*Proof.* Strong approximation (see [Vi, p. 61]) yields the identifications

$$\hat{R}^\times \backslash \hat{B}^\times / B^\times = R_\ell^\times \backslash B_\ell^\times / R[\frac{1}{\ell}]^\times, \quad \underline{\hat{R}}^\times \backslash \hat{B}^\times / B^\times = \underline{R}_\ell^\times \backslash B_\ell^\times / R[\frac{1}{\ell}]^\times.$$

Let  $\tilde{\mathcal{G}}_\ell$  be the graph  $\mathcal{T}_\ell / \Gamma_\ell$ , and let  $\vec{\mathcal{E}}(\tilde{\mathcal{G}}_\ell)$  be the set of oriented edges of  $\tilde{\mathcal{G}}_\ell$ . Using the map  $\kappa_\ell$ , one obtains the identifications

$$\hat{R}^\times \backslash \hat{B}^\times / B^\times = \vec{\mathcal{E}}(\tilde{\mathcal{G}}_\ell), \quad \underline{\hat{R}}^\times \backslash \hat{B}^\times / B^\times = \mathcal{V}(\tilde{\mathcal{G}}_\ell).$$

To conclude, observe that  $\mathcal{E}(\mathcal{G}_\ell)$  is identified with  $\vec{\mathcal{E}}(\tilde{\mathcal{G}}_\ell)$  by mapping the unoriented edge  $\{v, w\} \pmod{\Gamma_{\ell,1}}$  to the oriented edge  $(v, w) \pmod{\Gamma_\ell}$  if  $v$  is even, and to  $(w, v) \pmod{\Gamma_\ell}$  if  $v$  is odd (see [BD4, Lemma 2.2] for more details). Moreover, the disjoint union  $\mathcal{V}(\tilde{\mathcal{G}}_\ell) \times \{0, 1\}$  is identified with  $\mathcal{V}(\mathcal{G}_\ell)$  by mapping  $(\tilde{v}, 0)$  to the even lift of  $\tilde{v}$  and  $(\tilde{v}, 1)$  to the odd lift of  $\tilde{v}$ , under the natural projection  $\mathcal{V}(\tilde{\mathcal{G}}_\ell) \rightarrow \mathcal{V}(\mathcal{G}_\ell)$ .

Let

$$\alpha : \hat{R}^\times \backslash \hat{B}^\times / B^\times \rightarrow \underline{\hat{R}}^\times \backslash \hat{B}^\times / B^\times$$

be the natural projection induced by the inclusion  $R \subset \underline{R}$ . Denote by  $\hat{w}$  the element of  $\hat{B}^\times$  whose local components away from  $\ell$  are equal to 1, and whose local component at  $\ell$  maps by  $\kappa_\ell$  to the diagonal matrix  $\text{diag}(1, \ell)$ . Since the Eichler order  $S = \hat{w} \hat{R}^\times \hat{w}^{-1} \cap B$  of level  $M^+$  contains  $R$ , there is a natural projection from  $\hat{R}^\times \backslash \hat{B}^\times / B^\times$  to  $\hat{S}^\times \backslash \hat{B}^\times / B^\times$ . Call  $\beta$  the composition of this projection with the identification of  $\hat{S}^\times \backslash \hat{B}^\times / B^\times$  with  $\underline{\hat{R}}^\times \backslash \hat{B}^\times / B^\times$  induced by the assignment  $\hat{b} \mapsto \hat{w}^{-1} \hat{b}$ .

Let

$$\mathcal{X}'_\ell = \mathbb{Z}[\hat{R}^\times \backslash \hat{B}^\times / B^\times]^0, \quad \text{respectively,} \quad \mathcal{X}''_\ell = \mathbb{Z}[\underline{\hat{R}}^\times \backslash \hat{B}^\times / B^\times]^0$$

be the module of degree zero formal divisors with  $\mathbb{Z}$ -coefficients supported on  $\hat{R}^\times \backslash \hat{B}^\times / B^\times$ , respectively, on  $\underline{\hat{R}}^\times \backslash \hat{B}^\times / B^\times$ .

Define a degeneracy map

$$d_* : \mathcal{X}'_\ell \rightarrow (\mathcal{X}''_\ell)^2$$

by extending by linearity the rule  $d_*(x) = (\beta(x), 0) - (0, \alpha(x))$  for  $x$  in  $\hat{R}^\times \backslash \hat{B}^\times / B^\times$ .

**PROPOSITION 5.7.** *The modules  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0$  and  $\text{Im}(\delta_*)$  of Proposition 5.5 are identified with  $\mathcal{X}'_\ell$  and  $(\mathcal{X}''_\ell)^2$ , respectively. Moreover, the map  $\delta_*$  corresponds under these identifications to  $d_*$  (up to sign).*

*Proof.* Lemma 5.6 implies directly that  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0$  and  $\text{Im}(\delta_*)$  are identified with  $\mathcal{X}'_\ell$  and with a submodule of  $\mathbb{Z}[\underline{\hat{R}}^\times \backslash \hat{B}^\times / B^\times]^2$ . A more careful study shows that  $d_*$  is surjective, and corresponds to  $\delta_*$  under the above identifications. Proposition 5.7 follows.

Let  $\mathbb{T}$  be the Hecke algebra acting faithfully on  $\mathcal{X}_\ell$ , induced by the Hecke algebra acting on the jacobian  $J$  (by Picard functoriality). The module  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0$  is also endowed with a faithful action of an algebra  $\mathbb{T}'$  of Hecke correspondences, coming from its double coset description given in Lemma 5.6: see [BD1, 1.5] and also Section 1.1. Similarly, the identification of  $\text{Im}(\delta_*)$  with  $(\mathcal{X}_\ell'')^2$  equips  $\text{Im}(\delta_*)$  with the diagonal action of Hecke operators  $T_q''$  for  $q$  not dividing  $M/\ell$ , and  $U_q''$  for  $q$  dividing  $M/\ell$ . Moreover, the quotient  $\text{Im}(\delta_*)$  of  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0$  is stable for the induced action of  $\mathbb{T}'$ : denote by  $\mathbb{T}''$  the algebra quotient of  $\mathbb{T}'$  acting faithfully on it. Write  $\tilde{\mathbb{T}}$  for the polynomial ring with  $\mathbb{Z}$ -coefficients generated by the indeterminates  $\tilde{T}_q$ , for primes  $q$  not dividing  $M$ , and  $\tilde{U}_q$ , for primes  $q$  dividing  $M$ . Note that the Hecke algebras  $\mathbb{T}$ ,  $\mathbb{T}'$  and  $\mathbb{T}''$  are natural quotients of  $\tilde{\mathbb{T}}$ .

PROPOSITION 5.8. (1) *The exact sequence of Proposition 5.5 is equivariant for the natural actions of  $\tilde{\mathbb{T}}$  on  $\mathcal{X}_\ell$ ,  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0$ , and  $\text{Im}(\delta_*)$  defined above.*

(2) *For  $q \neq \ell$ , the  $q^{\text{th}}$  Hecke operator  $T_q' \in \mathbb{T}'$  ( $q \nmid M$ ) or  $U_q' \in \mathbb{T}'$  ( $q|M$ ) acts on  $\text{Im}(\delta_*) \simeq (\mathcal{X}_\ell'')^2$  via the diagonal action induced by the natural  $q^{\text{th}}$  Hecke operator  $T_q''$  ( $q \nmid M$ ) or  $U_q'' \in \mathbb{T}'$  ( $q|M$ ), respectively, acting on the double coset space  $\mathcal{X}_\ell''$ . Moreover, the induced action of  $U_\ell' \in \mathbb{T}'$  on  $\text{Im}(\delta_*)$  is given by the formula  $(x, y) \mapsto (T_\ell''x - y, \ell x)$ , where  $T_\ell''$  is the  $\ell^{\text{th}}$  Hecke operator acting on  $\mathcal{X}_\ell''$ .*

(3) *The Hecke algebra  $\mathbb{T}'$  is isomorphic to the Hecke algebra acting on modular forms of level  $M$  which are new at  $M^-/\ell$ . Its quotient  $\mathbb{T}$ , respectively,  $\mathbb{T}''$  is isomorphic to the Hecke algebra acting on modular forms of level  $M$  which are new at  $M^-$ , respectively, which are new at  $M^-/\ell$  and old at  $\ell$ .*

*Sketch of proof. Step 1.* Recalling that  $M^-$  is divisible by an even number of primes, fix a prime  $m$  such that  $m \neq \ell$  and  $m | M^-$ . Denote by  $X'$ , respectively,  $X''$  the Shimura curve  $X_{M+\ell m, M^-/\ell m}$ , respectively,  $X_{M+m, M^-/\ell m}$ . Write  $\mathcal{X}_m(X')$ , respectively,  $\mathcal{X}_m(X'')$  for the character group of  $X'$ , respectively,  $X''$  at  $m$ . These character groups are described in Proposition 5.3: one has

$$\mathcal{X}_m(X') = \mathbb{Z}[\mathcal{S}_m(X')]^0, \quad \mathcal{X}_m(X'') = \mathbb{Z}[\mathcal{S}_m(X'')]^0.$$

Let  $a$  be the natural projection from  $X'$  to  $X''$ , and let  $b$  be the composition of the Atkin-Lehner involution at  $\ell$  acting on  $X'$  with  $a$ . They induce a surjective degeneracy map

$$b_* - a_* : \mathcal{X}_m(X') \rightarrow \mathcal{X}_m(X'')^2.$$

The module  $\mathcal{X}_m(X')$ , respectively,  $\mathcal{X}_m(X'')$  is equipped with the action of a Hecke algebra induced by the Hecke algebra acting (by Picard functoriality) on

$J' = \text{Pic}^0(X')$ , respectively,  $J'' = \text{Pic}^0(X'')$ . Moreover,  $\mathcal{X}_m(X'')^2$  is endowed with a quotient Hecke action, induced by the map  $b_* - a_*$ .

*Step 2.* The assignment sending a supersingular modulus to its ring of endomorphisms equipped with an orientation sets up a bijection of  $\mathcal{S}_m(X')$ , respectively,  $\mathcal{S}_m(X'')$  onto the set of  $B^\times$ -conjugacy classes of oriented Eichler orders in  $B$  of level  $M^+\ell$ , respectively,  $M^+$  (see [BD2, §2]). These conjugacy classes are classified by the elements of the double coset space  $\hat{R}^\times \backslash \hat{B}^\times / B^\times$ , respectively,  $\hat{R}^\times \backslash \hat{B}^\times / B^\times$  (see [BD3, §1]). One obtains the isomorphisms

$$\mathcal{X}_m(X') \simeq \mathcal{X}'_\ell, \quad \mathcal{X}_m(X'') \simeq \mathcal{X}''_\ell.$$

It can be checked that the above identifications are Hecke equivariant, and that the degeneracy maps  $d_*$  and  $b_* - a_*$  coincide (up to sign) under these identifications.

*Step 3.* The previous steps give a geometric interpretation of the sequence of Proposition 5.5 in terms of the Shimura curves  $X$ ,  $X'$  and  $X''$ . Then, the proof of Proposition 5.8 in the special case where  $M^- = \ell m$  is a product of two primes is contained in Ribet's paper [Ri2]: see Theorems 3.19, 3.20 and 3.21. The details on the general case are provided in [Ri1].

Let us state for future use the exact sequence of character groups considered in the proof of Proposition 5.8.

**PROPOSITION 5.9.** *Let  $m$  be a prime  $\neq \ell$  which divides  $M^-$ . There is a Hecke equivariant exact sequence*

$$0 \rightarrow \mathcal{X}_\ell(X_{M^+, M^-}) \rightarrow \mathcal{X}_m(X_{M^+ \ell m, M^- / \ell m}) \rightarrow \mathcal{X}_m(X_{M^+ m, M^- / \ell m})^2 \rightarrow 0.$$

*Remark.* The identification of the group  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0$  with  $\mathcal{X}_m(X_{M^+ \ell m, M^- / \ell m})$  and of  $\text{Im}(\delta_*)$  with  $\mathcal{X}_m(X_{M^+ m, M^- / \ell m})^2$  can be made entirely canonical. The reader should consult [Ri1] for a thorough discussion of this issue.

**COROLLARY 5.10** (The Jacquet-Langlands correspondence). *The subring of the endomorphism ring of  $J$  generated by the natural Hecke correspondences on  $X$  is identified with the Hecke algebra acting on cusp forms of level  $M$  which are new at  $M^-$ .*

*Proof.* By Theorem 5.1, combined with the theory of  $\ell$ -adic uniformization of Mumford and Tate (see [GvdP, Chs. VI and VIII]),  $J = \text{Pic}^0(X)$  has purely toric reduction at  $\ell$ . Hence, the Hecke algebra  $\mathbb{T}$  acting on  $\mathcal{X}_\ell(X)$  is canonically identified with the Hecke algebra acting on  $J$ . The result follows from the interpretation of  $\mathbb{T}$  given in Proposition 5.8.

5.5. *Connected components.* Let  $\ell$  be a prime dividing  $M^-$ , and let  $\Phi_\ell$  be the group of connected components of  $J$  at  $\ell$ , defined in Section 5.3.

Let us begin by reviewing Grothendieck's description of  $\Phi_\ell$ . Let

$$\langle \cdot, \cdot \rangle : \mathcal{X}_\ell \times \mathcal{X}_\ell \rightarrow \mathbb{Z}$$

be the natural *monodromy pairing* on  $\mathcal{X}_\ell$ . It is defined to be the restriction to  $\mathcal{X}_\ell$ , by the embedding of Proposition 5.2, of the diagonal pairing on  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]$  given by the rule

$$\langle e, e' \rangle = \nu_e \delta_{e,e'},$$

where  $\nu_e$  is the weight of the edge  $e$  defined in Section 5.3. Let

$$j : \mathcal{X}_\ell \rightarrow \mathcal{X}_\ell^\vee$$

be the map induced by the monodromy pairing, where  $\mathcal{X}_\ell^\vee$  denotes the  $\mathbb{Z}$ -dual of  $\mathcal{X}_\ell$ . Using the notation introduced before Proposition 5.8, note that the map  $j$  is  $\mathbb{T}$ -equivariant, provided that the action of  $\mathbb{T}$  on the source of  $j$  is induced by Albanese functoriality and the action of  $\mathbb{T}$  on the target of  $j$  is induced by Picard functoriality. The reader is referred to the remark at the beginning of Section 5.4, where these two actions are compared, and to [Ri2, pp. 448–449] for an extended discussion.

PROPOSITION 5.11. *The group  $\Phi_\ell$  is canonically identified with the cokernel of  $j$ , and hence fits into the Hecke-equivariant exact sequence*

$$0 \rightarrow \mathcal{X}_\ell \xrightarrow{j} \mathcal{X}_\ell^\vee \xrightarrow{\tau_\ell} \Phi_\ell \rightarrow 0.$$

*Proof.* See Theorems 11.5 and 12.5 of [Groth], and Sections 1 and 2 of [Ed].

COROLLARY 5.12. *There is a natural map*

$$\mathbb{Z}[\mathcal{V}(\mathcal{G}_\ell)]^0 \xrightarrow{\omega_\ell} \Phi_\ell.$$

*Proof.* Recall the exact sequence

$$0 \rightarrow \mathcal{X}_\ell \xrightarrow{i} \mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)] \xrightarrow{\partial^*} \mathbb{Z}[\mathcal{V}(\mathcal{G}_\ell)] \rightarrow \mathbb{Z} \rightarrow 0$$

of Proposition 5.2, where the map from  $\mathbb{Z}[\mathcal{V}(\mathcal{G}_\ell)]$  to  $\mathbb{Z}$  is the degree map. The free  $\mathbb{Z}$ -modules  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]$  and  $\mathbb{Z}[\mathcal{V}(\mathcal{G}_\ell)]$  can be identified canonically with their  $\mathbb{Z}$ -duals, by using their distinguished bases. Taking the  $\mathbb{Z}$ -dual of the above sequence yields

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}[\mathcal{V}(\mathcal{G}_\ell)] \xrightarrow{\partial^*} \mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)] \xrightarrow{i^\vee} \mathcal{X}_\ell^\vee \rightarrow 0,$$

where the map from  $\mathbb{Z}$  to  $\mathbb{Z}[\mathcal{V}(\mathcal{G}_\ell)]$  is the diagonal embedding, and  $\partial^*$  sends a vertex  $v$  to  $-\sum_{s(e)=v} e$  if  $v$  is even, and to  $\sum_{t(e)=v} e$  if  $v$  is odd. Let  $j_0$  denote the map from  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]$  to itself induced by the pairing  $\langle \cdot, \cdot \rangle$  defined above. Note that the map  $j$  of Proposition 5.11 is equal to  $i^\vee \circ j_0 \circ i$ . Since  $\Phi_\ell$  is identified

with the cokernel of  $j$ , the assignment sending an element  $x$  of  $\mathbb{Z}[\mathcal{V}(\mathcal{G}_\ell)]^0$  to the element  $(\tau_\ell \circ i^\vee \circ j_0)(y)$  of  $\Phi_\ell$ , where  $y$  is chosen so that  $\partial_*(y) = x$ , defines the sought-for map  $\omega_\ell$ .

By the notation of Proposition 5.8, a  $\tilde{\mathbb{T}}$ -module  $C$  is *Eisenstein* if the relations

$$\tilde{T}_q c = (q+1)c \quad \text{for } q \nmid M$$

hold for all  $c$  in  $C$ .

**PROPOSITION 5.13.** *The restriction of the map  $\omega_\ell$  of Proposition 5.12 to  $\text{Im}(\delta_*)$ , viewed as a submodule of  $\mathbb{Z}[\mathcal{V}(\mathcal{G}_\ell)]^0$  via Proposition 5.5, induces a  $\tilde{\mathbb{T}}$ -equivariant map*

$$\bar{\omega}_\ell : \text{Im}(\delta_*) / ((U'_\ell)^2 - 1) \rightarrow \Phi_\ell.$$

*Furthermore, the kernel and cokernel of  $\bar{\omega}_\ell$  are Eisenstein.*

*Remark.* The proof of Proposition 5.13 follows the argument in the proof of Theorem 4.3 of [Ri2], where the result is proved in the special case where  $M^-$  is the product of two primes. See also the previous work of Jordan and Livné [JoLi2]. For the details on the Hecke compatibility in the statement of Proposition 5.13, see Chapters 3 and 4 of [Ri2], particularly Remark 3.24, and the remark at the beginning of Section 5.4.

*Sketch of proof.* Let  $j'$  be the map from  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0$  to its  $\mathbb{Z}$ -dual  $(\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0)^\vee$  induced by the pairing  $\langle \cdot, \cdot \rangle$  defined before. Similarly, let

$$\langle\langle \cdot, \cdot \rangle\rangle : \mathbb{Z}[\mathcal{V}(\mathcal{G}_\ell)] \times \mathbb{Z}[\mathcal{V}(\mathcal{G}_\ell)] \rightarrow \mathbb{Z}$$

be the pairing defined by the rule  $\langle\langle v, v' \rangle\rangle = \nu_v \delta_{v, v'}$ , where the weight  $\nu_v$  of the vertex  $v$  is defined to be the order of the stabilizer in  $\Gamma_{\ell,1}$  of any lift of  $v$  to  $\mathcal{V}(\mathcal{T}_\ell)$ . (Note that the quantity  $\nu_e$ , defined in a more geometric fashion in Section 5.3, could equivalently be defined to be the order of the stabilizer in  $\Gamma_{\ell,1}$  of any lift of  $e$  to  $\mathcal{E}(\mathcal{T}_\ell)$ .) Let

$$j'' : \text{Im}(\delta_*) \rightarrow \text{Im}(\delta_*)^\vee$$

be the map induced by  $\langle\langle \cdot, \cdot \rangle\rangle$ . Fix a prime divisor  $m \neq \ell$  of  $M^-$ , and let  $\Phi'_m$ , respectively  $\Phi''_m$ , denote the group of connected components of the Néron model over  $\mathbb{Z}_{m^2}$  of the jacobian of  $X_{M+\ell m, M^-/\ell m}$ , respectively, of  $X_{M+m, M^-/\ell m}$ . The groups  $\Phi'_m$  and  $\Phi''_m$  are Eisenstein. This follows from a generalization to Shimura curves of Theorem 3.12 of [Ri2], which can be obtained from the results of [Bu] and [JoLi3]. The analogue of Proposition 5.11 applied to the character groups at  $m$  of  $X_{M+\ell m, M^-/\ell m}$  and  $X_{M+m, M^-/\ell m}$ , respectively (see Theorem 11.5 and 12.5 of [Groth], and Sections 1 and 2 of [Ed]), combined with Proposition 5.9 and the proof of Proposition 5.8, yields the identifications

$$\Phi'_m = (\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0)^\vee / j'(\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0), \quad \Phi''_m \times \Phi''_m = \text{Im}(\delta_*)^\vee / j''(\text{Im}(\delta_*)).$$

Let

$$j'_0 : \text{Im}(\delta_*) \rightarrow \frac{(\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0)^\vee}{j'(i(\mathcal{X}_\ell))}$$

be the composition of the isomorphism of  $\text{Im}(\delta_*)$  with  $\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0/i(\mathcal{X}_\ell)$ , induced by  $\delta_*$ , with the map induced by  $j'$ . Set

$$\sigma : \text{Im}(\delta_*) \rightarrow \text{Im}(\delta_*), \quad (x, y) \mapsto ((\ell + 1)x + T_\ell'' y, T_\ell'' x + (\ell + 1)y).$$

One obtains a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Im}(\delta_*) & \xrightarrow{j''} & \text{Im}(\delta_*)^\vee & \longrightarrow & \Phi_m'' \times \Phi_m'' \longrightarrow 0 \\ & & \downarrow \sigma & & \downarrow \delta_*^\vee & & \downarrow \\ 0 & \longrightarrow & \text{Im}(\delta_*) & \xrightarrow{j'_0} & \frac{(\mathbb{Z}[\mathcal{E}(\mathcal{G}_\ell)]^0)^\vee}{j'(i(\mathcal{X}_\ell))} & \longrightarrow & \Phi_m' \longrightarrow 0, \end{array}$$

where  $\delta_*^\vee$  is induced by the  $\mathbb{Z}$ -dual of  $\delta_*$ . Note that  $\delta_*^\vee$  is injective, and its cokernel is identified with  $\Phi_\ell$ , in view of Proposition 5.11. Moreover, the cokernel of  $\sigma$  is identified with  $\text{Im}(\delta_*)/((U'_\ell)^2 - 1)$ . For, a calculation shows that the composition of the isomorphism

$$(x, y) \mapsto (-x, T_\ell'' x - y)$$

of  $\text{Im}(\delta_*)$  with  $\sigma$  gives the action of  $(U'_\ell)^2 - 1$ . Proposition 5.13 follows from the snake lemma applied to the above diagram.

Let  $X_{\mathbb{Z}_{\ell^2}}$  be (the base change of) the model of  $X$  introduced in Section 5.1, and let  $X_{\mathbb{F}_{\ell^2}}$  be the special fiber of  $X_{\mathbb{Z}_{\ell^2}}$ . Write

$$D = \sum_P n_P P \in \text{Div}^0(X)$$

for a degree zero divisor on  $X$  with integer coefficients, and  $\text{Supp}(D)$  for the support of  $D$ . Assume that  $D$  satisfies the following assumptions:

- (1) Each  $P \in \text{Supp}(D)$  is defined over  $\mathbb{Q}_{\ell^2}$ .
- (2) Each  $P \in \text{Supp}(D)$  reduces modulo  $\ell$  to a point  $r_\ell(P)$  which belongs to a unique irreducible component of  $X_{\mathbb{F}_{\ell^2}}$  (that is,  $r_\ell(P)$  is not a double point in  $X_{\mathbb{F}_{\ell^2}}$ ).

Thus,  $D$  determines an element

$$r_\ell(D) = \sum_P n_P r_\ell(P)$$

in  $\mathbb{Z}[\mathcal{V}(\mathcal{G}_\ell)]^0$ . Consider the specialization map

$$\partial_\ell : J(\mathbb{Q}_{\ell^2}) \rightarrow \Phi_\ell.$$

**PROPOSITION 5.14.** *Let  $D \in \text{Div}^0(X)$  be a divisor satisfying the above assumptions, and let  $[D] \in J(\mathbb{Q}_{\ell^2})$  be the class of  $D$ . Then*

$$\partial_\ell([D]) = \omega_\ell(r_\ell(D)).$$

*Proof.* This follows from Edixhoven's results in [Ed, §2]. (These results are built on Raynaud's description of  $\partial_\ell$  in terms of the minimal model of  $X$  over  $\mathbb{Z}_{\ell^2}$ , given in [Ray].)

5.6. *Raising the level and groups of connected components.* Let  $f : \vec{\mathcal{E}}(\mathcal{T})/\Gamma \rightarrow \mathbb{Z}_p$  be a form on  $\mathcal{T}/\Gamma$ , defined as in Section 1. Thus,  $f$  is attached to an integer factorization  $N = N^+N^-$ , where  $N^-$  is squarefree and divisible by an odd number of primes and  $N^+$  is divisible by  $p$ . (Note that this differs slightly from the notation of Section 1, where  $N$  was written as  $pN^+N^-$  and  $N^+$  was assumed to be prime to  $p$ .)

Let  $m$  be a distinguished prime divisor of  $N^-$ . In what follows,  $\ell$  denotes a prime number which does not divide  $N$ . Write  $\mathbb{T} = \mathbb{T}_{N^+,N^-}$ , respectively  $\mathbb{T}_\ell = \mathbb{T}_{N^+,N^-\ell}$ , for the Hecke algebra acting on cusp forms on  $\Gamma_0(N)$ , respectively on  $\Gamma_0(N\ell)$ , which are new at  $N^-$ , respectively at  $N^-\ell$ . Denote by  $t_q$  and  $u_q$ , respectively,  $T_q$  and  $U_q$  the Hecke operators in  $\mathbb{T}$ , respectively in  $\mathbb{T}_\ell$ .

The form  $f$  yields a surjective homomorphism

$$f : \mathbb{T} \rightarrow \mathbb{Z}/p^n\mathbb{Z}$$

(still denoted  $f$  by an abuse of notation). Write  $\mathcal{I}_f$  for the kernel of  $f$ , and  $\mathfrak{m}_f$  for the unique proper maximal ideal of  $\mathbb{T}$  containing  $\mathcal{I}_f$ .

In view of Corollary 5.10,  $\mathbb{T}$  can be identified with the  $m$ -new quotient of the Hecke algebra  $\mathbb{T}_{N^+,N^-/m}$  acting faithfully on  $X_{N^+,N^-/m}$ . Since the character group  $\mathcal{X}_m(X_{N^+,N^-/m})$  arises from the  $m$ -new part of  $J_{N^+,N^-/m}$ , it follows that  $\mathbb{T}$  acts naturally on it (by Picard functoriality).

**THEOREM 5.15.** *Assume:*

- (1)  $\mathfrak{m}_f$  is residually irreducible,
- (2) the completion  $\mathcal{X}_m(X_{N^+,N^-/m})_{\mathfrak{m}_f}$  of  $\mathcal{X}_m(X_{N^+,N^-/m})$  at  $\mathfrak{m}_f$  is free, of rank 1 over the completed Hecke algebra  $\mathbb{T}_{\mathfrak{m}_f}$ ,
- (3)  $\ell$  is a  $n$ -admissible prime relative to  $f$ .

*Then:*

- (1) *There exists a surjective homomorphism*

$$f_\ell : \mathbb{T}_\ell \rightarrow \mathbb{Z}/p^n\mathbb{Z}$$

*such that  $f_\ell(T_q) = f(t_q)$  for all  $q \nmid N\ell$ ,  $f_\ell(U_q) = f(u_q)$  for all  $q \mid N$ , and  $f_\ell(U_\ell) = \varepsilon$ , where  $\varepsilon = \pm 1$  is such that  $p^n$  divides  $\ell + 1 - \varepsilon f(t_\ell)$ .*

- (2) *Let  $\mathcal{I}_{f_\ell} \subset \mathbb{T}_\ell$  denote the kernel of the homomorphism  $f_\ell$ , and let  $\Phi_\ell$  denote the group of connected components of the Néron model of  $J_{N^+,N^-\ell}$  over  $\mathbb{Z}_{\ell^2}$ , defined in Section 5.3. There is a group isomorphism*

$$\Phi_\ell/\mathcal{I}_{f_\ell} \simeq \mathbb{Z}/p^n\mathbb{Z}.$$

*Remark.* Compare the proof of Theorem 5.15 with Section 7 of [Ri2], in which the special case, where  $M^-$  is a product of two primes and  $\mathcal{I}_f = \mathfrak{m}_f$  is a maximal ideal, is treated.

*Proof.* The second assumption implies the existence of an isomorphism

$$\mathcal{X}_m(X_{N+m, N^-/m})^2/\mathcal{I}_f \simeq (\mathbb{Z}/p^n\mathbb{Z})^2.$$

Write  $T'_q$  and  $U'_q$  for the Hecke operators in  $\mathbb{T}_{N+\ell, N^-}$ . By Proposition 5.9 and Proposition 5.8, there is an action of  $\mathbb{T}_{N+\ell, N^-}$  on  $\mathcal{X}_m(X_{N+m, N^-/m})^2$ , induced by the diagonal action of  $t_q$  for  $q \nmid N\ell$  and  $u_q$  for  $q \mid N$ , and such that the Hecke operator  $U'_\ell$  acts via the formula

$$(x, y) \mapsto (t_\ell x - y, \ell x).$$

Since  $\ell$  is  $n$ -admissible,  $t_\ell$  is equal modulo  $\mathcal{I}_f$  to  $\varepsilon(\ell + 1)$ . By combining this relation with the above formula for the action of  $U'_\ell$  and with the fact that  $p \geq 5$ , one obtains that  $U'_\ell + \varepsilon$  is invertible on  $\mathcal{X}_m(X_{N+m, N^-/m})^2/\mathcal{I}_f$ , and that the isomorphisms

$$\mathcal{X}_m(X_{N+m, N^-/m})^2/\langle \mathcal{I}_f, U'_\ell - \varepsilon \rangle \simeq \mathcal{X}_m(X_{N+m, N^-/m})^2/\langle \mathcal{I}_f, (U'_\ell)^2 - 1 \rangle \simeq \mathbb{Z}/p^n\mathbb{Z}$$

hold. Hence, the action of  $\mathbb{T}_{N+\ell, N^-}$  on the above quotient is via a surjective homomorphism

$$f'_\ell : \mathbb{T}_{N+\ell, N^-} \rightarrow \mathbb{Z}/p^n\mathbb{Z}.$$

Write  $\mathcal{I}_{f'_\ell}$  for the kernel of  $f'_\ell$ . By Proposition 5.13 and the residual irreducibility of  $\mathfrak{m}_f$ , combined with Proposition 5.9 and the remark after it, one finds an isomorphism

$$\Phi_\ell/\mathcal{I}_{f'_\ell} \simeq \mathcal{X}_m(X_{N+m, N^-/m})^2/\langle \mathcal{I}_f, (U'_\ell)^2 - 1 \rangle.$$

It follows that  $f'_\ell$  factors through  $\mathbb{T}_\ell$ , giving the sought for character  $f_\ell$ , and that  $\Phi_\ell/\mathcal{I}_{f'_\ell}$  is isomorphic to  $\mathbb{Z}/p^n\mathbb{Z}$ .

Write  $X^{(\ell)}$  for the Shimura curve  $X_{N^+, N^-/\ell}$ ,  $J^{(\ell)}$  for the jacobian of  $X^{(\ell)}$ , and  $\text{Pic}(X^{(\ell)})$  for the Picard variety of  $X^{(\ell)}$ . Denote by  $\mathcal{X}_\ell$  the character group of  $J^{(\ell)}$  at  $\ell$ , and by  $\text{Ta}_p(J^{(\ell)})$  the  $p$ -adic Tate module of  $J^{(\ell)}$ . Note that the Hecke algebra  $\mathbb{T}_\ell$  acts faithfully on  $J^{(\ell)}$ .

LEMMA 5.16. *Under the assumptions of Theorem 5.15, the exponent of the  $\mathbb{Z}$ -module  $\text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$  is equal to  $p^n$ .*

*Proof.* Since  $\text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$  is naturally a  $\mathbb{T}_\ell/\mathcal{I}_{f_\ell}$ -module, and  $\mathbb{T}_\ell/\mathcal{I}_{f_\ell}$  is isomorphic to  $\mathbb{Z}/p^n\mathbb{Z}$ , it follows that the exponent of  $\text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$  is at most  $p^n$ . On the other hand, the Mumford-Tate theory of  $\ell$ -adic uniformization, combined with the Cerednik-Drinfeld theorem (see Section 5.2 and [GvdP, Chs. VI and VIII]) shows the existence of a symmetric pairing

$$(48) \quad [ \ , \ ] : \mathcal{X}_\ell \times \mathcal{X}_\ell \rightarrow \mathbb{Q}_\ell^\times$$

such that there exists an exact sequence of  $\mathbb{T}_\ell[\text{Gal}(\bar{\mathbb{Q}}_{\ell^2}/\mathbb{Q}_{\ell^2})]$ -modules

$$(49) \quad 0 \rightarrow \mathcal{X}_\ell \xrightarrow{\tilde{j}} \mathcal{X}_\ell^\vee \otimes \bar{\mathbb{Q}}_{\ell^2}^\times \rightarrow J^{(\ell)}(\bar{\mathbb{Q}}_{\ell^2}) \rightarrow 0,$$

where  $\mathcal{X}_\ell^\vee$  denotes the  $\mathbb{Z}$ -dual  $\text{Hom}(\mathcal{X}_\ell, \mathbb{Z})$  of  $\mathcal{X}_\ell$ , and where the map  $\tilde{j}$  is induced by [ , ]. The sequence (49) is Hecke-equivariant, provided that the action of  $\mathbb{T}_\ell$  on the first term is induced by Albanese functoriality, and the action on the second and third term is induced by Picard functoriality. Note also that the need to extend scalars to  $\mathbb{Q}_{\ell^2}$  in (49) arises from the fact that the isomorphism of rigid spaces stated in Theorem 5.1 is only defined over  $\mathbb{Q}_{\ell^2}$  and not over  $\mathbb{Q}_\ell$ . The pairing [ , ] is related to monodromy pairing of Section 5.5 by the rule

$$(50) \quad \text{ord}_\ell \circ [ , ] = \langle , \rangle$$

(see [M, Th. 7.6]). In view of Theorem 5.15, choose an element  $c$  of  $\Phi_\ell/\mathcal{I}_{f_\ell}$  of order  $p^n$ , and lift  $c$  to an element  $\tilde{c}$  of  $\Phi_\ell$  of  $p$ -power order  $p^{n'}$ , with  $n' \geq n$ . In view of Proposition 5.11, fix an element  $b \in \mathcal{X}_\ell^\vee$  such that  $\tau_\ell(b) = \tilde{c}$ , and let  $a$  be the element of  $\mathcal{X}_\ell$  such that  $p^{n'}b = j(a)$ . Formula (50) shows that  $\text{ord}_\ell$  of the period  $\tilde{j}(a) \in \mathcal{X}_\ell^\vee \otimes \bar{\mathbb{Q}}_\ell^\times$  is divisible by  $p^{n'}$ . Thanks to the sequence (49), the choice of a  $p^{n'}$ -root of  $\tilde{j}(a)$  determines an element  $\tilde{t}$  of  $J^{(\ell)}[p^{n'}]$  defined over an unramified extension of  $\mathbb{Q}_\ell$ , whose natural image in  $\Phi_\ell$  is equal to  $\tilde{c}$ . Writing  $t$  for the image of  $\tilde{t}$  in  $J^{(\ell)}[p^{n'}]/\mathcal{I}_{f_\ell}$ , then the natural image of  $t$  in  $\Phi_\ell/\mathcal{I}_{f_\ell}$  is equal to  $c$ . Since

$$\text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell} = J^{(\ell)}[p^{n'}]/\mathcal{I}_{f_\ell},$$

it follows that  $t$  is an element of order  $\geq p^n$  in  $\text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$ . Since the exponent of  $\text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$  is at most  $p^n$ , this proves Lemma 5.16.

**THEOREM 5.17.** *Under the assumptions of Theorem 5.15, the Galois representations  $\text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$  and  $T_{f,n}$  are isomorphic.*

*Proof.* Let  $\mathfrak{m}_{f_\ell}$  be the maximal ideal in  $\mathbb{T}_\ell$  containing  $\mathcal{I}_{f_\ell}$ . (Thus,  $\mathbb{T}_\ell/\mathfrak{m}_{f_\ell}$  is isomorphic to  $\mathbb{Z}/p\mathbb{Z}$ .) The proof is naturally divided into two steps. In the first step, one works modulo  $\mathfrak{m}_{f_\ell}$ , and a known result of [BoLeRi] on the structure of  $\text{Ta}_p(J^{(\ell)})/\mathfrak{m}_{f_\ell}$  is invoked.

*Step 1.* This first step shows that  $\text{Ta}_p(J^{(\ell)})/\mathfrak{m}_{f_\ell}$  is isomorphic to  $T_{f,1}$ . Taking  $p$ -torsion in the sequence (49) yields the exact sequence of  $\mathbb{T}_\ell[\text{Gal}(\bar{\mathbb{Q}}_{\ell^2}/\mathbb{Q}_{\ell^2})]$ -modules

$$(51) \quad 0 \rightarrow \mathcal{X}_\ell^\vee \otimes \mu_p \rightarrow J^{(\ell)}[p] \rightarrow \mathcal{X}_\ell/p \rightarrow 0.$$

After tensoring (51) with  $\mathbb{T}_\ell/\mathfrak{m}_{f_\ell}$ , one finds

$$(52) \quad 0 \rightarrow ((\mathcal{X}_\ell^\vee/\mathfrak{m}_{f_\ell})/\mathcal{Y}) \otimes \mu_p \rightarrow J^{(\ell)}[p]/\mathfrak{m}_{f_\ell} \rightarrow \mathcal{X}_\ell/\mathfrak{m}_{f_\ell} \rightarrow 0,$$

where  $\mathcal{Y}$  is a certain submodule of  $\mathcal{X}_\ell^\vee/\mathfrak{m}_{f_\ell}$ . Taking Galois cohomology over  $\mathbb{Q}_{\ell^2}$  of (52) yields an exact sequence

$$(53) \quad \begin{aligned} \mathcal{X}_\ell/\mathfrak{m}_{f_\ell} &\rightarrow H^1(\mathbb{Q}_{\ell^2}, ((\mathcal{X}_\ell^\vee/\mathfrak{m}_{f_\ell})/\mathcal{Y}) \otimes \mu_p) \\ &\rightarrow H^1(\mathbb{Q}_{\ell^2}, J^{(\ell)}[p]/\mathfrak{m}_{f_\ell}) \rightarrow H^1(\mathbb{Q}_{\ell^2}, \mathcal{X}_\ell/\mathfrak{m}_{f_\ell}). \end{aligned}$$

Note the identifications

$$(54) \quad \begin{aligned} H^1(\mathbb{Q}_{\ell^2}, ((\mathcal{X}_\ell^\vee/\mathfrak{m}_{f_\ell})/\mathcal{Y}) \otimes \mu_p) &= ((\mathcal{X}_\ell^\vee/\mathfrak{m}_{f_\ell})/\mathcal{Y}) \otimes H^1(\mathbb{Q}_{\ell^2}, \mu_p) \\ &= ((\mathcal{X}_\ell^\vee/\mathfrak{m}_{f_\ell})/\mathcal{Y}) \otimes \mathbb{Q}_{\ell^2}^\times/(\mathbb{Q}_{\ell^2}^\times)^p = (\mathcal{X}_\ell^\vee/\mathfrak{m}_{f_\ell})/\mathcal{Y}, \end{aligned}$$

where the last equality follows from the fact that  $\ell$  is an admissible prime and hence  $p \nmid \ell^2 - 1$ . Moreover,  $J^{(\ell)}[p]/\mathfrak{m}_{f_\ell} = \mathrm{Ta}_p(J^{(\ell)})/\mathfrak{m}_{f_\ell}$ , and

$$H^1(\mathbb{Q}_{\ell^2}, \mathcal{X}_\ell/\mathfrak{m}_{f_\ell}) = \mathrm{Hom}_{\mathrm{unr}}(\mathrm{Gal}(\bar{\mathbb{Q}}_{\ell^2}/\mathbb{Q}_{\ell^2}), \mathcal{X}_\ell/\mathfrak{m}_{f_\ell}) = \mathrm{Hom}(\mathbb{Z}/p\mathbb{Z}, \mathcal{X}_\ell/\mathfrak{m}_{f_\ell})$$

by local class field theory, since  $p \nmid \ell^2 - 1$ . Thus, (53) can be re-written as

$$(55) \quad \mathcal{X}_\ell/\mathfrak{m}_{f_\ell} \rightarrow (\mathcal{X}_\ell^\vee/\mathfrak{m}_{f_\ell})/\mathcal{Y} \rightarrow H^1(\mathbb{Q}_{\ell^2}, \mathrm{Ta}_p(J^{(\ell)})/\mathfrak{m}_{f_\ell}) \rightarrow H_{\mathrm{unr}}^1(\mathbb{Q}_{\ell^2}, \mathcal{X}_\ell/\mathfrak{m}_{f_\ell}).$$

The first map in (55) is induced by the monodromy pairing on  $\mathcal{X}_\ell$ : this follows from the definition of the sequence (49) which induces (55); see also the Proposition on page IV–32 of [Se2], which covers a special case. By Proposition 5.11, one obtains the exact sequence

$$(56) \quad 0 \rightarrow \bar{\Phi}_\ell/\mathfrak{m}_{f_\ell} \rightarrow H^1(\mathbb{Q}_{\ell^2}, \mathrm{Ta}_p(J^{(\ell)})/\mathfrak{m}_{f_\ell}) \rightarrow H_{\mathrm{unr}}^1(\mathbb{Q}_{\ell^2}, \mathcal{X}_\ell/\mathfrak{m}_{f_\ell}),$$

where  $\bar{\Phi}_\ell/\mathfrak{m}_{f_\ell}$  is a quotient of  $\Phi_\ell/\mathfrak{m}_{f_\ell}$ . By the main result of [BoLeRi], the module  $\mathrm{Ta}_p(J^{(\ell)})/\mathfrak{m}_{f_\ell}$  is semisimple over  $\mathbb{F}_p[\mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})]$ , and hence is isomorphic to  $k \geq 1$  copies of  $T_{f,1}$  by the Eichler-Shimura relations. Hence,  $H^1(\mathbb{Q}_{\ell^2}, \mathrm{Ta}_p(J^{(\ell)})/\mathfrak{m}_{f_\ell})$  is isomorphic to  $H^1(\mathbb{Q}_{\ell^2}, T_{f,1})^k$ . In view of the results of Section 2.2, the  $\mathbb{F}_p$ -vector space  $H^1(\mathbb{Q}_{\ell^2}, T_{f,1})^k$  is  $2k$ -dimensional; furthermore, it can be decomposed as the direct sum of two  $k$ -dimensional subspaces, one of which is generated by unramified cohomology classes, and the other by ramified cohomology classes. Since  $\Phi_\ell/\mathfrak{m}_{f_\ell}$  is 1-dimensional by Theorem 5.15, it follows from (56) and the proof of Lemma 2.6 that  $\bar{\Phi}_\ell/\mathfrak{m}_{f_\ell}$  is equal to  $\Phi_\ell/\mathfrak{m}_{f_\ell}$ , and that  $k$  is equal to 1. Hence  $\mathrm{Ta}_p(J^{(\ell)})/\mathfrak{m}_{f_\ell}$  is isomorphic to  $T_{f,1}$ .

*Step 2.* It remains to show that  $\mathrm{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$  is isomorphic to  $T_{f,n}$ . There is a natural  $G_{\mathbb{Q}}$ -equivariant projection

$$(57) \quad \mathrm{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell} \xrightarrow{\pi} \mathrm{Ta}_p(J^{(\ell)})/\mathfrak{m}_{f_\ell}.$$

In view of Lemma 5.16, let  $t$  be an element of order  $p^n$  in  $\mathrm{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$ , and let  $\bar{t} = \pi(t)$ . Since  $T_{f,1}$  is irreducible, there is an element  $g \in G_{\mathbb{Q}}$  such that  $\bar{t}$  and  $\bar{t}^g$  are a basis for  $\mathrm{Ta}_p(J^{(\ell)})/\mathfrak{m}_{f_\ell}$ . Nakayama's lemma shows that

$\mathrm{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$  is generated by  $t$  and  $t^g$ . Moreover, since  $g$  acts as an automorphism of  $\mathrm{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$ , the order of  $t^g$  is equal to the order of  $t$ . Finally, using the assumption made in the introduction that the Galois representation  $T_{f,1}$  has image isomorphic to  $\mathbf{GL}_2(\mathbb{F}_p)$ , one checks directly that the submodules generated by  $t$  and  $t^g$  have trivial intersection, so that  $\mathrm{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$  is isomorphic to  $(\mathbb{Z}/p^n\mathbb{Z})^2$ . This implies that  $\mathrm{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$  is isomorphic to  $T_{f,n}$ .

*Remarks.* 1. Repeating the argument in the proof of the first step of Theorem 5.17, with  $p^n$  replacing  $p$  and the ideal  $\mathcal{I}_{f_\ell}$  replacing  $\mathfrak{m}_{f_\ell}$ , yields the exact sequence (analogue of (56))

$$(58) \quad 0 \rightarrow \bar{\Phi}_\ell/\mathcal{I}_{f_\ell} \rightarrow H^1(K_\ell, \mathrm{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}) \rightarrow H_{\mathrm{unr}}^1(K_\ell, \mathcal{X}_\ell/\mathcal{I}_{f_\ell}),$$

where  $\bar{\Phi}_\ell/\mathcal{I}_{f_\ell}$  is a quotient of  $\Phi_\ell/\mathcal{I}_{f_\ell}$ . This exact sequence will be of use in the proof of Corollary 5.18.

2. In view of Remark 1, equations (49) and (54), with  $\mathcal{I}_{f_\ell}$  replacing  $\mathfrak{m}_{f_\ell}$ , show that the natural Kummer map from  $J^{(\ell)}(K_\ell)$  to  $H^1(K_\ell, \mathrm{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell})$  factors through the map of specialization to connected components from  $J^{(\ell)}(K_\ell)$  to  $\bar{\Phi}_\ell/\mathcal{I}_{f_\ell}$ .

Recall the quadratic imaginary field  $K$  introduced in the previous sections. Note that the  $n$ -admissible prime  $\ell$  is inert in  $K$  by definition, so that the completion  $K_\ell$  is isomorphic to  $\mathbb{Q}_{\ell^2}$ . Given  $m \geq 0$ , let  $\mathcal{O}_m = \mathbb{Z} + p^{m+1}\mathcal{O}_K$  be the order of  $K$  of conductor  $p^{m+1}$ . Let  $\tilde{K}_m$  be the ring class field of  $K$  of conductor  $p^{m+1}$ , and let  $\tilde{K}_\infty$  be the union of the  $\tilde{K}_m$ . (The field  $\tilde{K}_\infty$  was introduced in Section 2.1.) The field  $\tilde{K}_m$  can be constructed by adjoining to  $K$  the value  $j(\mathcal{O}_m)$  of the modular function  $j$  (viewed as a function of lattices) on  $\mathcal{O}_m$ . By the theory of complex multiplication,  $\tilde{K}_m$  is an abelian extension of  $K$ , which contains the Hilbert class field  $\tilde{K}$  of  $K$ . The Galois group of  $\tilde{K}_m$  over  $\tilde{K}$  is cyclic, of order  $p^m(p - \varepsilon(p))/u$ , where  $\varepsilon(p)$  is equal to 1 or  $-1$  depending on whether  $p$  is split or inert in  $K$ , respectively, and where  $u$  denotes half the order of the group of units of  $K$ . Write  $\tilde{G}_m$  for the Galois group of  $\tilde{K}_m$  over  $K$ , and  $\tilde{G}_\infty$  for the Galois group of  $\tilde{K}_\infty$  over  $K$ . By class field theory, the Galois group  $\tilde{G}_\infty$  is identified with the group  $\tilde{G}_\infty$  defined in Section 1.2 in terms of the ideles of  $K$ .

Write  $\Phi_{\ell,m}$ , respectively,  $\Phi'_{\ell,m}$  for  $\bigoplus_{\lambda|\ell} \Phi_\lambda$ , where the sum is taken over the primes  $\lambda$  of  $K_m$  and of  $\tilde{K}_m$  dividing  $\ell$ , respectively, and  $\Phi_\lambda$  denotes the group of connected components of  $J^{(\ell)}$  at  $\lambda$ . Define  $\hat{\Phi}_\ell$ , respectively  $\tilde{\Phi}_\ell$ , to be the inverse limit of the groups  $\Phi_{\ell,m}$ , respectively  $\Phi'_{\ell,m}$ , with respect to the norm maps. Since the prime  $\ell$  is inert in  $K$ , it splits completely in  $\tilde{K}_\infty/K$ . Hence, the choice of a prime of  $\tilde{K}_\infty$  above  $\ell$  identifies  $\hat{\Phi}_\ell$  with  $\Phi_\ell \otimes \mathbb{Z}[[G_\infty]]$ , and  $\tilde{\Phi}_\ell$  with  $\Phi_\ell \otimes \mathbb{Z}[[\tilde{G}_\infty]]$ . It follows that the identification of  $\Phi_\ell/\mathcal{I}_{f_\ell}$  with  $\mathbb{Z}/p^n\mathbb{Z}$  of

Theorem 5.15 yields isomorphisms

$$\hat{\Phi}_\ell/\mathcal{I}_{f_\ell} \simeq \Lambda/p^n\Lambda, \quad \tilde{\Phi}_\ell/\mathcal{I}_{f_\ell} \simeq \mathbb{Z}/p^n\llbracket\tilde{G}_\infty\rrbracket.$$

Write  $\hat{J}^{(\ell)}(K_\infty)/\mathcal{I}_{f_\ell}$ , respectively  $\hat{J}^{(\ell)}(\tilde{K}_\infty)/\mathcal{I}_{f_\ell}$ , for the inverse limit of the modules  $J^{(\ell)}(K_{m,\ell})/\mathcal{I}_{f_\ell}$ , respectively  $J^{(\ell)}(\tilde{K}_{m,\ell})/\mathcal{I}_{f_\ell}$ , with respect to the norm maps. The inverse limit of the maps of specialization to connected components yields the maps

$$(59) \quad \hat{\partial}_\ell : \hat{J}^{(\ell)}(K_\infty)/\mathcal{I}_{f_\ell} \rightarrow \hat{\Phi}_\ell/\mathcal{I}_{f_\ell} \simeq \Lambda/p^n,$$

$$(60) \quad \tilde{\partial}_\ell : \hat{J}^{(\ell)}(\tilde{K}_\infty)/\mathcal{I}_{f_\ell} \rightarrow \tilde{\Phi}_\ell/\mathcal{I}_{f_\ell} \simeq \mathbb{Z}/p^n\llbracket\tilde{G}_\infty\rrbracket.$$

COROLLARY 5.18. (1) *There is an isomorphism*

$$\Phi_\ell/\mathcal{I}_{f_\ell} \rightarrow H_{\text{sing}}^1(K_\ell, T_{f,n}),$$

*which is canonical up to the choice of an identification of  $\text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$  with  $T_{f,n}$ .*

(2) *There is an isomorphism*

$$\hat{\Phi}_\ell/\mathcal{I}_{f_\ell} \rightarrow \hat{H}_{\text{sing}}^1(K_{\infty,\ell}, T_{f,n}),$$

*which is canonical up to the choice of an identification of  $\text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$  with  $T_{f,n}$ .*

(3) *There is a commutative diagram*

$$\begin{array}{ccc} \hat{J}^{(\ell)}(K_\infty)/\mathcal{I}_{f_\ell} & \longrightarrow & \hat{H}^1(K_\infty, T_{f,n}) \\ \downarrow \hat{\partial}_\ell & & \downarrow \partial_\ell \\ \hat{\Phi}_\ell/\mathcal{I}_{f_\ell} & \longrightarrow & \hat{H}_{\text{sing}}^1(K_{\infty,\ell}, T_{f,n}), \end{array}$$

*where the top horizontal arrow arises from the natural Kummer map, the lower horizontal arrow is the isomorphism defined above, and  $\partial_\ell$  is the residue map defined in Section 2.2. Furthermore, there is a similar commutative diagram having the group  $\hat{J}^{(\ell)}(\tilde{K}_\infty)/\mathcal{I}_{f_\ell}$  as its source.*

*Proof.* By Theorem 5.17,  $\text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$  is isomorphic to  $T_{f,n}$ . Hence, by the results of Section 2.2,  $H^1(K_\ell, \text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell})$  is free of rank two over  $\mathbb{Z}/p^n\mathbb{Z}$ ; moreover, its submodule of unramified classes has rank one. By Theorem 5.15,  $\Phi_\ell/\mathcal{I}_{f_\ell}$  is isomorphic to  $\mathbb{Z}/p^n\mathbb{Z}$ . The exact sequence (58) shows that all the elements of  $\tilde{\Phi}_\ell/\mathcal{I}_{f_\ell}$  map to ramified classes. It follows that  $\tilde{\Phi}_\ell/\mathcal{I}_{f_\ell}$  is equal to  $\Phi_\ell/\mathcal{I}_{f_\ell}$ . Furthermore, the residue map induces a surjection of  $\Phi_\ell/\mathcal{I}_{f_\ell}$  onto the rank one group  $H_{\text{sing}}^1(K_\ell, T_{f,n})$ . This defines the isomorphism stated in part one of Corollary 5.18. The second part is a formal consequence of the first. The third part follows from the definition of the isomorphism in part two, combined with Remark 2 after the proof of Theorem 5.17.

### 6. The theory of complex multiplication

(Reference: [BD1, §2]). Fix a positive integer  $M$ , a rational prime  $p$ , and an imaginary quadratic field  $K$  of discriminant  $-D$ , such that  $p \mid M$  but  $p^2 \nmid M$ , and  $(M, D) = 1$ . Define an integer decomposition

$$M = M^+ M^-$$

such that  $(M^+, M^-) = 1$ , and  $M^+$  is divisible by  $p$  and by the prime divisors of  $M/p$  which are split in  $K$ . Assume that:

- (1)  $M^-$  is squarefree,
- (2)  $M^-$  is the product of an even number of primes.

Let  $X = X_{M^+, M^-}$  be the Shimura curve attached in Section 5.1 to an Eichler order  $\mathcal{R}$  of level  $M^+$  in the indefinite quaternion algebra  $\mathcal{B}$  of discriminant  $M^-$ . For all  $m \geq 0$ , there is a point  $P_m$  on  $X$ , which has complex multiplication by the order  $\mathcal{O}_m$  of  $K$  of conductor  $p^{m+1}$  (introduced after the proof of Theorem 5.17). More precisely, let  $(A_m, \iota_m, C_m)$  be a triple corresponding to  $P_m$  via the moduli interpretation of  $X$  given in Section 5.1. Write  $\text{End}(P_m)$  for the ring of endomorphisms of  $A_m$  which commute with the action  $\iota_m$  and respect the level structure  $C_m$ . Then,

$$\text{End}(P_m) \simeq \mathcal{O}_m.$$

The point  $P_m$  is called a *Heegner point* of level  $m$ . By the theory of complex multiplication,  $P_m$  is defined over the field  $\tilde{K}_m$  (defined after the proof of Theorem 5.17). Choose the  $P_m$  so that they are *compatible*, in the sense that for all  $m \geq 0$ , there is an isogeny of degree  $p^2$  from  $A_m$  to  $A_{m+1}$ , which is equivariant for the actions  $\iota_m$  and  $\iota_{m+1}$ , and preserves the level structures  $C_m$  and  $C_{m+1}$ .

One has the following interpretation of the sequence  $\{P_m\}$  in terms of the Bruhat-Tits tree at  $p$ . Given a point  $P$  on the Shimura curve  $X_{M^+/p, M^-}$ , corresponding to a triple  $(A, \iota, C)$ , let  $\mathcal{T}^{(P)}$  denote the tree of  $p$ -isogenies of  $P$ . The vertices of  $\mathcal{T}^{(P)}$  correspond to points of  $X_{M^+/p, M^-}$  representing moduli related to  $(A, \iota, C)$  by an isogeny of  $p$ -power degree. Two vertices of  $\mathcal{T}^{(P)}$  are adjacent if the corresponding moduli are related by an isogeny of degree  $p^2$ . Thus, the oriented edges of  $\mathcal{T}^{(P)}$  are naturally identified with points on  $X$ . The tree  $\mathcal{T}^{(P)}$  is isomorphic to the Bruhat-Tits tree  $\mathcal{T}$ , and has a distinguished vertex  $v_P$  corresponding to  $P$ . There is a unique point  $P$  on  $X_{M^+/p, M^-}$  such that:

- (1)  $\text{End}(P) \simeq \mathcal{O}_K$ ,
- (2)  $P_0$  corresponds to an edge of the tree  $\mathcal{T}^{(P)}$ , with origin in  $P$ .

Then, the points  $P_m$  determine a half line of  $\mathcal{T}^{(P)}$  originating from  $P$ , with no back-trackings.

## 7. Construction of the Euler system

Notation and assumptions being as in Section 5.6, the construction of Section 6 yields a compatible family of Heegner points

$$P_m \in X^{(\ell)}(\tilde{K}_m)$$

for all  $m \geq 0$  (the notation  $X^{(\ell)}$  was defined right before Lemma 5.16). View  $P_m$  as an element of the Picard group  $\text{Pic}(X^{(\ell)})(\tilde{K}_m)$ . Since the ideal  $\mathcal{I}_{f_\ell}$  is not Eisenstein, the natural inclusion

$$J^{(\ell)}(\tilde{K}_m)/\mathcal{I}_{f_\ell} \rightarrow \text{Pic}(X^{(\ell)})(\tilde{K}_m)/\mathcal{I}_{f_\ell}$$

is an isomorphism. Let  $\alpha_p$  be the unit root of Frobenius at  $p$ . Write  $P_m^*$  for the image of  $\alpha_p^{-m} P_m$  in  $J^{(\ell)}(\tilde{K}_m)/\mathcal{I}_{f_\ell}$ . The points  $P_m^*$  are norm-compatible. Hence their images by the coboundary maps

$$J^{(\ell)}(\tilde{K}_m)/\mathcal{I}_{f_\ell} \rightarrow H^1(\tilde{K}_m, \text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell})$$

yield a sequence of cohomology classes which are compatible under the corestriction maps. The choice of an isomorphism of  $\text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$  with  $T_{f,n}$ , which exists by Theorem 5.17, gives a class  $\tilde{\kappa}(\ell)$  in  $\hat{H}^1(\tilde{K}_\infty, T_{f,n})$ , where  $\hat{H}^1(\tilde{K}_\infty, T_{f,n})$  denotes the inverse limit under the corestriction maps of the groups  $H^1(\tilde{K}_m, T_{f,n})$ . Define  $\kappa(\ell)$  in  $\hat{H}^1(K_\infty, T_{f,n})$  to be the corestriction from  $\tilde{K}_\infty$  to  $K_\infty$  of  $\tilde{\kappa}(\ell)$ . In other words, when  $Q_m$  is the norm from  $\tilde{K}_\infty$  to  $K_\infty$  of  $P_m^*$ , the class  $\kappa(\ell)$  is the natural image in  $\hat{H}^1(K_\infty, T_{f,n})$  of the sequence  $Q_m$  via the coboundary maps.

## 8. The first explicit reciprocity law

This section is devoted to the proof of Theorem 4.1, with notation as in the previous sections. Recall that the class  $\kappa(\ell)$  is constructed from a family of points on the Shimura curve  $X^{(\ell)}$ ; hence, it can be viewed as an element of the usual (compactified) Selmer group of  $J^{(\ell)}$  over  $K_\infty$  relative to the Galois module  $\text{Ta}_p(J^{(\ell)})/\mathcal{I}_{f_\ell}$ . This shows that  $\kappa(\ell)$  belongs to  $\hat{H}_\ell^1(K_\infty, T_{f,n})$ .

Recall the groups  $\hat{\Phi}_\ell$  and  $\tilde{\Phi}_\ell$ , and the maps  $\hat{\partial}_\ell$  and  $\tilde{\partial}_\ell$ , which were defined in Section 5.6 (see equations (59) and (60)).

LEMMA 8.1. *Theorem 4.1 is implied by the equality*

$$\tilde{\partial}_\ell(\{P_m^*\}) = \tilde{\mathcal{L}}_f \pmod{p^n}.$$

*Remark.* Note that the terms in the equality of Lemma 8.1 are well defined only up to multiplication by elements of  $\mathbb{Z}_p^\times$  and of  $\tilde{G}_\infty$ .

*Proof.* The element  $\hat{\partial}_\ell(\{Q_m\})$  is mapped to  $\partial_\ell(\kappa(\ell))$  by the isomorphism of Corollary 5.18. Since  $\hat{\partial}_\ell(\{Q_m\})$  is the norm of  $\tilde{\partial}_\ell(\{P_m^*\})$ , and  $\mathcal{L}_f$  is the natural image of  $\tilde{\mathcal{L}}_f$  in  $\Lambda$ , the claim follows.

In view of Lemma 8.1, one is reduced to studying the specialization of the Heegner points  $P_m$  to connected components. To begin with, it is necessary to recall the  $\ell$ -adic description of the point  $P_m$  given in Section 5 of [BD3], and based on Drinfeld's moduli interpretation of the  $\ell$ -adic upper half-plane. Let  $\bar{P}_m$  denote the reduction of  $P_m$  modulo a fixed prime above  $\ell$ , and let

$$\text{End}(P_m) \rightarrow \text{End}(\bar{P}_m)$$

be the map obtained by reduction of endomorphisms. Recall that  $\text{End}(P_m)$  is isomorphic to the quadratic order  $\mathcal{O}_m$ . Moreover, the ring  $\text{End}(\bar{P}_m)[\frac{1}{\ell}]$  is isomorphic to  $\underline{R}[\frac{1}{\ell}]$ , where  $\underline{R}$  denotes an Eichler order of level  $N^+$  in the definite quaternion algebra  $B$  of discriminant  $N^-$ , which can be chosen to be independent of  $m$ . Extending scalars by  $\mathbb{Z}[\frac{1}{\ell}]$ , one obtains an injection

$$\Psi_m^0 : \mathcal{O}_m[\frac{1}{\ell}] \rightarrow \underline{R}[\frac{1}{\ell}].$$

It can be shown that  $\Psi_m^0$  is well defined up to conjugation by elements in  $\underline{R}[\frac{1}{\ell}]_1^\times$ . Therefore,  $\Psi_m^0$  can be identified with an element of the space

$$(\text{Hom}(K, B) \times \mathcal{V}(\mathcal{T}_\ell)) / \Gamma_{\ell,1},$$

by mapping  $\Psi_m^0$  to the pair  $(\Psi_m, v_m)$ , where  $\Psi_m$  is the extension of scalars of  $\Psi_m^0$ , and  $v_m$  is the vertex of  $\mathcal{T}_\ell$  corresponding to the unique maximal order of  $B_\ell$  containing  $\Psi_m(\mathcal{O}_m)$ . The embedding  $\Psi_m$  induces an action of  $K_\ell^\times$  on  $\mathcal{H}_\ell$ , having two fixed points which belong to  $K_\ell - \mathbb{Q}_\ell$ , and which are conjugate by the generator of  $\text{Gal}(K_\ell/\mathbb{Q}_\ell)$ . Then,  $P_m$  is identified with the image in  $\mathcal{H}_\ell/\Gamma_{\ell,1}$  of one of these two points, via the isomorphism of Theorem 5.1. (A suitable condition of normalization specifies which point corresponds to  $P_m$ , but this will not be of use here.) Let

$$r_\ell : \mathcal{H}_\ell(\mathbb{C}_\ell) / \Gamma_{\ell,1} \longrightarrow \mathcal{V}(\mathcal{G}_\ell) \cup \mathcal{E}(\mathcal{G}_\ell) = (\mathcal{V}(\mathcal{T}_\ell) \cup \mathcal{E}(\mathcal{T}_\ell)) / \Gamma_{\ell,1}$$

be the reduction map (see for example Chapter I of [BC], and also Sections 1 and 6 of [BD3]). Here  $\mathcal{G}_\ell$  is the dual graph  $\mathcal{T}_\ell/\Gamma_{\ell,1}$  of  $X^{(\ell)}$  at  $\ell$ . Given  $P$  in  $X^{(\ell)}(\mathbb{C}_\ell)$ , viewed as a point in  $\mathcal{H}_\ell(\mathbb{C}_\ell)/\Gamma_{\ell,1}$ ,  $r_\ell(P)$  is equal to a vertex  $v$  if the reduction of  $P$  modulo  $\ell$  lands in the single irreducible component of the fiber  $X_{\mathbb{F}_{\ell^2}}^{(\ell)}$  corresponding to  $v$ , and  $r_\ell(P)$  is equal to an edge  $e$  if  $P$  reduces to the singular point corresponding to  $e$ . The above description of  $P_m$  in terms of an  $\ell$ -adic argument in  $\mathcal{H}_\ell$  shows that the image of  $P_m$  by  $r_\ell$  is equal to the vertex  $v_m \pmod{\Gamma_{\ell,1}}$ . This follows from the  $\mathbf{GL}_2(\mathbb{Q}_\ell)$ -equivariance of the reduction map, combined with the fact that  $P_m$  corresponds to a fixed point for the action of  $\Psi_m(K_\ell^\times)$  on  $\mathcal{H}_\ell(\mathbb{C}_\ell)$  and  $v_m$  is the unique fixed point for the action of  $\Psi_m(K_\ell^\times)$  on  $\mathcal{V}(\mathcal{T}_\ell) \cup \mathcal{E}(\mathcal{T}_\ell)$ . Hence, the reduction of  $P_m$  modulo  $\ell$  lands in the single irreducible component of the fiber  $X_{\mathbb{F}_{\ell^2}}^{(\ell)}$  defined by  $v_m \pmod{\Gamma_{\ell,1}}$ . Furthermore, by Proposition 5.14,  $v_m \pmod{\Gamma_{\ell,1}}$  computes the image of  $P_m$  in the group of connected components  $\Phi_{\ell,m}/\mathcal{I}_{f_\ell}$ .

On the other hand, recalling that  $p$  divides the level of the order  $\underline{R}$ , we see that strong approximation (see [Vi, p. 61]) gives an identification

$$(\mathrm{Hom}(K, B) \times \mathcal{V}(\mathcal{T}_\ell)) / \Gamma_{\ell,1} = (\mathrm{Hom}(K, B) \times \vec{\mathcal{E}}(\mathcal{T}_p)) / \Gamma_p$$

where, in the notations of Section 1.1,  $\mathcal{T}_p = \mathcal{T}$  and  $\Gamma_p = \Gamma$ . The compatibility condition on the Heegner points  $P_m$  translates to the condition that they may be represented by pairs  $(\Psi, e_m)$  in  $\mathrm{Hom}(K, B) \times \vec{\mathcal{E}}(\mathcal{T}_p)$ , where  $\Psi$  arises from an embedding

$$\Psi^0 : \mathcal{O}_K[1/p] \rightarrow \underline{R}[1/p]$$

which does not depend on  $m$ , and where the  $e_m$  determine a half line with no back-trackings in  $\mathcal{T}_p$ . The results of [BD3, §5] show that the action of  $\tilde{G}_\infty$  on the  $P_m$  is compatible with the action of  $\tilde{G}_\infty$  on the edges  $e_m$ , which was defined in Section 1.2 using the embedding  $\Psi$ . Fix a prime  $\lambda_\infty$  of  $\tilde{K}_\infty$  above  $\ell$ , and set  $\lambda_m = \lambda_\infty \cap \tilde{K}_m$ . For  $\sigma \in \tilde{G}_\infty$ , write  $\partial_{\lambda_m}(P_m^\sigma)$  for the natural image of  $P_m^\sigma$  in the group of connected components  $\Phi_{\lambda_m} / \mathcal{I}_{f_\ell} \simeq \mathbb{Z}/p^n\mathbb{Z}$ . The use of Proposition 5.14 described above, together with the identification coming from strong approximation, shows that the map  $\partial_{\lambda_m}$  on the points  $P_m^\sigma$  can be viewed as a function

$$\vec{\mathcal{E}}(\mathcal{T}_p) / \Gamma_p \longrightarrow \mathbb{Z}/p^n\mathbb{Z}.$$

By multiplicity one, this function is equal modulo  $p^n$  to the eigenform  $f$  introduced in Section 5.6 (up to multiplication by an element of  $(\mathbb{Z}/p^n\mathbb{Z})^\times$ ). By comparing with definition (17), one finds that the equality

$$(61) \quad \partial_{\lambda_m}(P_m^\sigma) = [\sigma, e_m]_f \pmod{p^n}$$

holds for a suitable choice of  $\lambda_\infty$ . (Note that the possible choices of  $\lambda_\infty$  are permuted by  $\tilde{G}_\infty$ . Similarly, the choices of embedding  $\Psi$  and of a sequence of edges in the definition of the  $p$ -adic  $L$ -function given in Section 1.2 show that  $\tilde{\mathcal{L}}_f$  is well defined only up to multiplication by elements of  $\tilde{G}_\infty$ .) It follows from (61) that

$$\partial_{\lambda_m}(\sigma P_m^*) = \alpha_p^{-m} [\sigma, e_m]_f \pmod{p^n}.$$

In view of equation (20) and of the definition of  $\tilde{\mathcal{L}}_f$ , this concludes the proof.

## 9. The second explicit reciprocity law

This section is devoted to the proof of Theorem 4.2. Recall that  $\ell_1$  and  $\ell_2$  are distinct  $n$ -admissible primes relative to  $f$ , such that  $p^n$  divides  $\ell_1 + 1 - \varepsilon_1 a_{\ell_1}(f)$  and  $\ell_2 + 1 - \varepsilon_2 a_{\ell_2}(f)$ , with  $\varepsilon_1$  and  $\varepsilon_2$  equal to  $\pm 1$ . Let  $\mathbb{T}_{\ell_1}$  be the Hecke algebra acting on  $X^{(\ell_1)}$ . As in the previous sections, equip  $J^{(\ell_1)}$  with the action of  $\mathbb{T}_{\ell_1}$  induced by Picard functoriality. Since  $f$  is  $p$ -isolated, the assumptions of Theorem 5.15 are satisfied in the current setting. Thus, by

Theorem 5.17,  $\mathrm{Ta}_p(J^{(\ell_1)})/\mathcal{I}_{f_{\ell_1}}$  is isomorphic to  $T_{f,n}$  as a Galois module. Fix such an isomorphism. Then, the map

$$J^{(\ell_1)}(K_{\ell_2})/\mathcal{I}_{f_{\ell_1}} \rightarrow H^1(K_{\ell_2}, \mathrm{Ta}_p(J^{(\ell_1)})/\mathcal{I}_{f_{\ell_1}})$$

arising from Kummer theory yields a map

$$(62) \quad J^{(\ell_1)}(K_{\ell_2})/\mathcal{I}_{f_{\ell_1}} \rightarrow H^1(K_{\ell_2}, T_{f,n}).$$

The image of (62) is equal to  $H_{\mathrm{fin}}^1(K_{\ell_2}, T_{f,n})$ , since  $T_{f,n}$  is unramified at  $\ell_2$  and  $\ell_2$  is a prime of good reduction for  $J^{(\ell_1)}$ . Since  $p \neq \ell_2$ , the map induced by reduction modulo  $\ell_2$

$$(63) \quad J^{(\ell_1)}(K_{\ell_2})/\mathcal{I}_{f_{\ell_1}} \rightarrow J^{(\ell_1)}(\mathbb{F}_{\ell_2^2})/\mathcal{I}_{f_{\ell_1}}$$

is an isomorphism. Hence, by composing the inverse of (63) with (62), and fixing an identification of  $H_{\mathrm{fin}}^1(K_{\ell_2}, T_{f,n})$  with  $\mathbb{Z}/p^n\mathbb{Z}$  (use Lemma 2.6), one obtains a surjective map

$$(64) \quad J^{(\ell_1)}(\mathbb{F}_{\ell_2^2})/\mathcal{I}_{f_{\ell_1}} \rightarrow \mathbb{Z}/p^n\mathbb{Z}.$$

Let  $\mathcal{S}_{\ell_2} \subset X^{(\ell_1)}(\mathbb{F}_{\ell_2^2})$  denote the set of supersingular points of  $X^{(\ell_1)}$  in characteristic  $\ell_2$ , and let  $\mathrm{Div}(\mathcal{S}_{\ell_2})$ , respectively  $\mathrm{Div}^0(\mathcal{S}_{\ell_2})$ , be the module of formal divisors, respectively, degree zero formal divisors with  $\mathbb{Z}$ -coefficients supported on  $\mathcal{S}_{\ell_2}$ .

One may define two different actions of  $\mathbb{T}_{\ell_1}$  on  $\mathrm{Div}(\mathcal{S}_{\ell_2})$  and  $\mathrm{Div}^0(\mathcal{S}_{\ell_2})$ , by using either Picard or Albanese functoriality (see the remark at the beginning of Section 5.4, which explains that these two actions differ by an Atkin-Lehner involution  $w_{N^+,1}$ ). The latter action is the usual action defined on supersingular points, and is more natural if one views the supersingular points as being points in  $X^{(\ell_1)}(\mathbb{F}_{\ell_2^2})$ . The former action is more natural when viewing divisors on supersingular points as giving rise to points in  $\mathrm{Pic}(X^{(\ell_1)})(\mathbb{F}_{\ell_2^2})$  and  $J^{(\ell_1)}(\mathbb{F}_{\ell_2^2})$ , on which the action of  $\mathbb{T}_{\ell_1}$  was defined via Picard functoriality. Since  $f_{\ell_1}$  is an eigenform for  $w_{N^+,1}$ , which acts via multiplication by  $\pm 1$ , the choice of a specific Hecke action makes no difference for the purpose of establishing the  $\mathbb{T}_{\ell_1}$ -equivariance of the maps defined below; let us establish the convention that the Hecke action on supersingular points be the Albanese one.

Since the inclusion of  $\mathrm{Div}^0(\mathcal{S}_{\ell_2})$  in  $\mathrm{Div}(\mathcal{S}_{\ell_2})$  induces an identification of  $\mathrm{Div}^0(\mathcal{S}_{\ell_2})/\mathcal{I}_{f_{\ell_1}}$  with  $\mathrm{Div}(\mathcal{S}_{\ell_2})/\mathcal{I}_{f_{\ell_1}}$  (use the fact that  $\mathcal{I}_{f_{\ell_1}}$  is not Eisenstein), one obtains a natural map

$$(65) \quad \mathrm{Div}(\mathcal{S}_{\ell_2}) \rightarrow J^{(\ell_1)}(\mathbb{F}_{\ell_2^2})/\mathcal{I}_{f_{\ell_1}}.$$

The composition of (65) with the surjection (64) yields a map

$$\gamma : \mathrm{Div}(\mathcal{S}_{\ell_2}) \rightarrow \mathbb{Z}/p^n\mathbb{Z}.$$

Write  $T_q$  ( $q \nmid N\ell_1$ ) and  $U_q$  ( $q \mid N\ell_1$ ) for the  $q^{\mathrm{th}}$  Hecke operator in  $\mathbb{T}_{\ell_1}$ , and  $\bar{T}_q$  and  $\bar{U}_q$  for the natural image of  $T_q$  and  $U_q$ , respectively, in  $\mathbb{T}_{\ell_1}/\mathcal{I}_{f_{\ell_1}} = \mathbb{Z}/p^n\mathbb{Z}$ .

Thus, the following equalities modulo  $p^n$  hold:  $\bar{T}_q \equiv a_q(f)$  for  $q \nmid N\ell_1$ ,  $\bar{U}_q \equiv a_q(f)$  for  $q \mid N$ , and  $\bar{U}_{\ell_1} \equiv \varepsilon_1$ .

LEMMA 9.1. *The relations*

$$\begin{aligned} \gamma(T_q x) &= \bar{T}_q \gamma(x) \quad (q \nmid N\ell_1\ell_2), & \gamma(U_q x) &= \bar{U}_q \gamma(x) \quad (q \mid N\ell_1), \\ \gamma(T_{\ell_2} x) &= \bar{T}_{\ell_2} \gamma(x), & \gamma(\text{Frob}_{\ell_2} x) &= \varepsilon_2 \gamma(x) \end{aligned}$$

hold for  $x \in \text{Div}(\mathcal{S}_{\ell_2})$ .

*Proof.* As observed in the proof of Lemma 2.6,  $H_{\text{fin}}^1(K_{\ell_2}, T_{f,n})$  is identified with the module  $T_{f,n}/(\text{Frob}_{\ell_2}^2 - 1)T_{f,n}$  of  $G_{K_{\ell_2}}$ -coinvariants of  $T_{f,n}$ . Therefore, the map  $\gamma$  is defined by sending a point  $x$  to the image of  $((\text{Frob}_{\ell_2}^2 - 1)/p^n)x$  in  $T_{f,n}/(\text{Frob}_{\ell_2}^2 - 1)T_{f,n}$ . The first two identities are a direct consequence of this description of  $\gamma$ . As for the last two identities, note that the Eichler-Shimura relations identify  $T_{\ell_2}$  with the correspondence  $\text{Frob}_{\ell_2} + \text{Frob}_{\ell_2}^\vee$ , where  $\text{Frob}_{\ell_2}^\vee$  denotes the transpose of  $\text{Frob}_{\ell_2}$ . Furthermore, on points defined over  $\mathbb{F}_{\ell_2^2}$ , one has  $\text{Frob}_{\ell_2}^\vee x = \ell_2 \text{Frob}_{\ell_2} x$ , and hence  $T_{\ell_2} x = (\ell_2 + 1)\text{Frob}_{\ell_2} x$ . The claim follows from the fact that  $\text{Frob}_{\ell_2}$  acts on  $T_{f,n}$  with eigenvalues  $\varepsilon_2$  and  $\varepsilon_2 \ell_2$ , which implies that  $\text{Frob}_{\ell_2}$  acts as  $\varepsilon_2$  on the quotient  $T_{f,n}/(\text{Frob}_{\ell_2}^2 - 1)T_{f,n}$ .

PROPOSITION 9.2. *The map  $\gamma$  is surjective.*

*Proof.* Recall that  $p$  exactly divides  $N^+$ , and that the Shimura curve  $X = X^{(\ell_1)}$  was defined in Section 5.1 in terms of an Eichler order  $\mathcal{R}$  of level  $N^+$  in the indefinite quaternion algebra  $\mathcal{B}$  of discriminant  $N^-\ell_1$ . Let  $J = J^{(\ell_1)}$  be the jacobian of  $X$ . Write  $J(\mathbb{F}_{\ell_2^2})^{ss}$  for the subgroup of  $J(\mathbb{F}_{\ell_2^2})$  generated by divisors supported on supersingular points. Since the map (64) is surjective, it is enough to show that the natural image of  $J(\mathbb{F}_{\ell_2^2})^{ss}$  in the group  $J(\mathbb{F}_{\ell_2^2})/\mathcal{I}_{f_{\ell_1}}$  is equal to the whole group. Let  $\Gamma^{(\ell_2)}$  denote the group of norm one elements in  $\mathcal{R}[1/\ell_2]^\times/\{\pm 1\}$ . Let  $\tilde{X}$  be the Shimura curve defined in the same way as  $X$  but imposing an extra  $\Gamma_1(p)$ -level structure. Let  $\tilde{\Gamma}^{(\ell_2)}$  be the finite index subgroup of  $\Gamma^{(\ell_2)}$  consisting of elements which are congruent to the standard unipotent matrices modulo  $p$ . Write  $\tilde{J}(\mathbb{F}_{\ell_2^2})$  for the  $\mathbb{F}_{\ell_2^2}$ -points of the jacobian of  $\tilde{X}$ , and  $\tilde{J}(\mathbb{F}_{\ell_2^2})^{ss}$  for the subgroup generated by divisors supported on supersingular points. Since  $\tilde{\Gamma}^{(\ell_2)}$  is torsion-free, the results of [I2] (see in particular Remark G, page 19) establish a canonical isomorphism

$$(66) \quad \tilde{J}(\mathbb{F}_{\ell_2^2})/\tilde{J}(\mathbb{F}_{\ell_2^2})^{ss} \simeq (\tilde{\Gamma}^{(\ell_2)})^{ab},$$

where  $(\tilde{\Gamma}^{(\ell_2)})^{ab}$  denotes the abelianization of  $\tilde{\Gamma}^{(\ell_2)}$ . By fixing an embedding of  $\mathcal{B}$  in  $M_2(\mathbb{Q}_{\ell_2})$ , one obtains an action of  $\tilde{\Gamma}^{(\ell_2)}$  on the Bruhat-Tits tree  $\mathcal{T}_{\ell_2}$ . Let  $v_0$  be the vertex of  $\mathcal{T}_{\ell_2}$  such that the stabilizer  $\tilde{\Gamma}_{v_0}^{(\ell_2)}$  of  $v_0$  in  $\tilde{\Gamma}^{(\ell_2)}$  is the subgroup of integral elements (relative to the fixed embedding of  $\tilde{\Gamma}^{(\ell_2)}$  in  $M_2(\mathbb{Q}_{\ell_2})$ ). Let

$e_0$  be the edge originating from  $v_0$  such that the stabilizer  $\tilde{\Gamma}_{e_0}^{(\ell_2)}$  of  $e_0$  is the subgroup of  $\tilde{\Gamma}_{v_0}^{(\ell_2)}$  whose elements are upper triangular modulo  $\ell_2$  relative to this embedding. Write  $v_1$  for the target of  $e_0$ . Note that  $\tilde{\Gamma}_{v_0}^{(\ell_2)}$ , respectively  $\tilde{\Gamma}_{e_0}^{(\ell_2)}$ , can be identified with the discrete subgroup of  $\mathbf{SL}_2(\mathbb{R})$  which defines the Shimura curve  $\tilde{X}$ , respectively, the Shimura curve  $\tilde{X}_0(\ell_2)$  defined in the same way as  $\tilde{X}$ , but with an extra  $\Gamma_0(\ell_2)$ -level structure imposed. The group  $\tilde{\Gamma}^{(\ell_2)}$  acts on the tree  $\mathcal{T}_{\ell_2}$  with the closed edge attached to  $e_0$  as a fundamental region. Hence the exact cohomology sequence in Proposition 13 of Section II.2.8 of [Se1], in the case  $i = 1$ ,  $M = \mathbb{F}_p$  and  $G = \tilde{\Gamma}^{(\ell_2)}$  becomes

$$(67) \quad 0 \longrightarrow \mathrm{Hom}(\tilde{\Gamma}^{(\ell_2)}, \mathbb{F}_p) \longrightarrow \mathrm{Hom}(\tilde{\Gamma}_{v_0}^{(\ell_2)}, \mathbb{F}_p) \oplus \mathrm{Hom}(\tilde{\Gamma}_{v_1}^{(\ell_2)}, \mathbb{F}_p) \\ \xrightarrow{d} \mathrm{Hom}(\tilde{\Gamma}_{e_0}^{(\ell_2)}, \mathbb{F}_p).$$

The modules appearing as the source and target of  $d$  are identified by duality with two copies or one copy of the  $p$ -torsion in the jacobian of  $\tilde{X}$  and  $\tilde{X}_0(\ell_2)$ , respectively. Moreover, the map  $d$  corresponds under these identifications to the map denoted  $\alpha_p$  in the statement of Theorem 2, p. 451 of [DT]. This theorem (the analogue of Ihara's lemma in the setting of Shimura curves) states that the action of  $G_{\mathbb{Q}}$  on each Jordan-Hölder constituent of the kernel of  $d$  factors through an abelian quotient of  $G_{\mathbb{Q}}$ . View  $f_{\ell_1}$  as a mod  $p^n$  modular form (with trivial character) on  $\tilde{X}$ , and write  $\tilde{\mathcal{I}}_{f_{\ell_1}}$  for the associated ideal in the Hecke algebra  $\tilde{\mathbb{T}}_{\ell_1}$ . Let  $\tilde{\mathfrak{m}}_{f_{\ell_1}}$  be the maximal ideal of  $\tilde{\mathbb{T}}_{\ell_1}$  containing  $\tilde{\mathcal{I}}_{f_{\ell_1}}$ . Since  $\tilde{\mathfrak{m}}_{f_{\ell_1}}$  corresponds to an irreducible mod  $p$  Galois representation, it follows from the sequence (67), combined with the semisimplicity result of [BoLeRi] and the above mentioned theorem of [DT], that  $\mathrm{Hom}(\tilde{\Gamma}^{(\ell_2)}, \mathbb{F}_p)[\tilde{\mathfrak{m}}_{f_{\ell_1}}]$  is trivial, and therefore that  $(\tilde{\Gamma}^{(\ell_2)})^{ab}/\tilde{\mathfrak{m}}_{f_{\ell_1}}$  is trivial. By Nakayama's lemma, this shows that

$$(68) \quad (\tilde{\Gamma}^{(\ell_2)})^{ab}/\tilde{\mathcal{I}}_{f_{\ell_1}} = 0.$$

Equation (66) then implies that the natural image of  $\tilde{J}(\mathbb{F}_{\ell_2^2})^{ss}$  in  $\tilde{J}(\mathbb{F}_{\ell_2^2})/\tilde{\mathcal{I}}_{f_{\ell_1}}$  fills the whole group. Finally, the cokernel of the natural map

$$(69) \quad \tilde{J}^{(\ell_1)}(\mathbb{F}_{\ell_2^2}) \rightarrow J^{(\ell_1)}(\mathbb{F}_{\ell_2^2})$$

can be identified with an abelian quotient of the (finite) image of  $\Gamma_0(p)$  in  $\mathbf{SL}_2(\mathbb{Z}/p\mathbb{Z})$ , and hence has order dividing  $p - 1$ ; see for example the results of Chapter 7 of [Co], particularly in pages 107 and 110. It follows in particular that the composition of the map (69) with the projection from  $J^{(\ell_1)}(\mathbb{F}_{\ell_2^2})$  to  $J^{(\ell_1)}(\mathbb{F}_{\ell_2^2})/p^n$  is surjective. When combined with the fact that  $\tilde{J}(\mathbb{F}_{\ell_2^2})^{ss}$  maps surjectively to  $\tilde{J}(\mathbb{F}_{\ell_2^2})/\tilde{\mathcal{I}}_{f_{\ell_1}}$  (as noted after equation (68)), this implies that  $J(\mathbb{F}_{\ell_2^2})^{ss}$  maps surjectively to  $J(\mathbb{F}_{\ell_2^2})/\mathcal{I}_{f_{\ell_1}}$ , as was to be shown.

*Remark.* The proof of Proposition 9.2 relies crucially on the analogue of Ihara's lemma proved in [DT]. In turn, Proposition 9.2 provides the key step

in the proof of Theorem 9.3, a raising-the-level result which is a theorem of [DT] (extended to mod  $p^n$  modular forms).

Following the notation of Proposition 3.12, let  $B'$  be the definite quaternion algebra of discriminant  $N^-\ell_1\ell_2$ ,  $R'$  an Eichler  $\mathbb{Z}[1/p]$ -order of level  $N^+$  in  $B$ , and  $\Gamma'$  the group  $(R')^\times/\mathbb{Z}[1/p]^\times$ . The next result contains the part of the statement of Proposition 3.12 that remains to be proved.

**THEOREM 9.3.** *There exists an eigenform  $g \in S_2(\mathcal{T}/\Gamma', \mathbb{Z}/p^n\mathbb{Z})$  such that*

$$\begin{aligned} T_q g &= a_q(f)g \quad (q \nmid N\ell_1\ell_2), & U_q g &= a_q(f)g \quad (q|N), \\ U_{\ell_1} g &= \varepsilon_1 g, & U_{\ell_2} g &= \varepsilon_2 g. \end{aligned}$$

*Proof.* Write  $\mathbb{T}_{\ell_1} = \mathbb{T}_{N^+, N^-\ell_1}$  for the Hecke algebra acting on cusp forms on  $\Gamma_0(N\ell_1)$  which are new at  $N^-\ell_1$ , and  $\mathbb{T}_{\ell_2, \ell_1} = \mathbb{T}_{N^+\ell_2, N^-\ell_1}$  for the Hecke algebra acting on cusp forms on  $\Gamma_0(N\ell_1\ell_2)$  which are new at  $N^-\ell_1$ . By Theorem 5.15, there is a mod  $p^n$  modular form

$$f_{\ell_1} : \mathbb{T}_{\ell_1} \rightarrow \mathbb{Z}/p^n\mathbb{Z}$$

such that  $f_{\ell_1}(T_q) = f(t_q)$  for all  $q \nmid N\ell_1$ ,  $f_{\ell_1}(U_q) = f(u_q)$  for all  $q | N$ , and  $f_{\ell_1}(U_{\ell_1}) = \varepsilon_1 f_{\ell_1}$ . Recall from the proof of Theorem 5.15 that the homomorphism  $f_{\ell_1}$  arises from the group of connected components at  $\ell_1$  attached to the Shimura curve  $X^{(\ell_1)} = X_{N^+, N^-\ell_1}$ . Consider now the Shimura curve  $X^{(\ell_2, \ell_1)} := X_{N^+\ell_2, N^-\ell_1}$ , on which the elements of  $\mathbb{T}_{\ell_2, \ell_1}$  act as correspondences. By Proposition 5.3, the character group  $\mathcal{X}_{\ell_2}$  of  $X^{(\ell_2, \ell_1)}$  at  $\ell_2$  is identified with the module  $\text{Div}^0(\mathcal{S}_{\ell_2})$  defined at the beginning of this section; here  $\mathcal{S}_{\ell_2}$  is viewed as the set of supersingular points in  $X^{(\ell_2, \ell_1)}(\mathbb{F}_{\ell_2^2})$  by the results of Section 5.1. Furthermore, the action of  $\mathbb{T}_{\ell_2, \ell_1}$  on  $\mathcal{X}_{\ell_2}$  induced from the action on  $\text{Pic}^0(X^{(\ell_2, \ell_1)})$  defined by Picard functoriality is compatible with the standard (Albanese) action of  $\mathbb{T}_{\ell_2, \ell_1}$  via correspondences on the set of supersingular points (see [Ri2, p. 445]). The results of [Wa] on the endomorphisms of supersingular abelian surfaces, combined with strong approximation, yield a  $\mathbb{T}_{\ell_2, \ell_1}$ -compatible identification of  $\mathcal{S}_{\ell_2}$  with  $\vec{\mathcal{E}}(\mathcal{T})/\Gamma'$ . (Note that  $\mathcal{X}_{\ell_2} = \text{Div}^0(\mathcal{S}_{\ell_2})$  pertains to the  $\ell_2$ -new part of the jacobian  $J^{(\ell_2, \ell_1)}$  of  $X^{(\ell_2, \ell_1)}$ , being  $\mathcal{X}_{\ell_2}$  the character group of the maximal torus of  $J^{(\ell_2, \ell_1)}$  over  $\mathbb{F}_{\ell_2^2}$ .) Therefore, the map  $\gamma$  defined above can also be viewed as a  $\mathbb{Z}/p^n\mathbb{Z}$ -valued map on  $\vec{\mathcal{E}}(\mathcal{T})/\Gamma'$ , whose values are not contained in any proper subgroup of  $\mathbb{Z}/p^n\mathbb{Z}$ , by Proposition 9.2. This map defines the sought-for modular form  $g$  on  $\Gamma'$ , as can be seen by appealing to Lemma 9.1. More precisely, let  $T_q^*$  (for  $q$  not dividing  $N\ell_1\ell_2$ ) and  $U_q^*$  (for  $q$  dividing  $N\ell_1\ell_2$ ) denote the Hecke operators in  $\mathbb{T}_{\ell_2, \ell_1}$ . (Note the discrepancy between the notation for the Hecke operators used in this proof and that used in the statement of Proposition 3.12, which is also reproduced in the statement of Theorem 9.3; in this proof, and throughout the section, the

symbols  $T_q$  and  $U_q$  are used to indicate the Hecke operators in  $\mathbb{T}_{\ell_1}$ .) Since the Hecke operators at  $q \neq \ell_2$  in  $\mathbb{T}_{\ell_1}$  and  $\mathbb{T}_{\ell_2, \ell_1}$  act in the same way on  $\gamma$  and  $g$ , respectively, Lemma 9.1 implies directly the relations of Theorem 9.3 for all  $q \neq \ell_2$ :

$$T_q^* g = T_q \gamma = a_q(f)g \quad (q \nmid N\ell_1\ell_2), \quad U_q g = U_q \gamma = a_q(f)g \quad (q \mid N\ell_1)$$

(in particular,  $T_{\ell_1}^* g = \varepsilon_1 g$ ). As for the operator  $U_{\ell_2}^*$ , it is known that  $U_{\ell_2}^* x = \text{Frob}_{\ell_2} x$  for  $x \in \mathcal{S}_{\ell_2}$  (see [Ri2, Prop. 3.8]). Hence, Lemma 9.1 yields

$$(U_q^* g)(x) = \gamma(\text{Frob}_{\ell_2} x) = \varepsilon_2 g(x).$$

This concludes the proof.

The proof of Theorem 9.3 implies the following result.

**COROLLARY 9.4.** *Under the identification of  $\mathcal{S}_{\ell_2}$  with  $\vec{\mathcal{E}}(\mathcal{T})/\Gamma'$ , the map  $\gamma$  corresponds to an eigenform  $g \in S_2(\mathcal{T}/\Gamma', \mathbb{Z}/p^n\mathbb{Z})$  satisfying the conclusions of Theorem 9.3.*

Consider the sequence  $\{P_m\}$  of Heegner points  $P_m \in X^{(\ell_1)}(\tilde{K}_m)$ , constructed in Section 6. Fix a prime  $\lambda_\infty$  of  $\tilde{K}_\infty$  above  $\ell_2$ , and let  $\lambda_m = \lambda_\infty \cap \tilde{K}_m$ . Since  $\ell_2$  is inert in  $K$ , the point  $P_m$  reduces modulo  $\lambda_m$  to a supersingular point  $\bar{P}_m \in X^{(\ell_1)}(\mathbb{F}_{\lambda_m})$ . When  $\mathbb{F}_{\lambda_m}$  is identified with  $\mathbb{F}_{\ell_2}$ ,  $\bar{P}_m$  can be viewed as an element of  $\mathcal{S}_{\ell_2}$ . Identifying  $\mathcal{S}_{\ell_2}$  with  $\vec{\mathcal{E}}(\mathcal{T})/\Gamma'$ , we can describe the sequence  $\{\bar{P}_m\}$  by a sequence of consecutive edges  $\{e_m\}$  in  $\vec{\mathcal{E}}(\mathcal{T})$ , modulo  $\Gamma'$ , in such a way that the map  $\text{End}(P_m) \rightarrow \text{End}(\bar{P}_m)$  of reduction of endomorphisms modulo  $\lambda_m$  induces by extension of scalars an embedding

$$\Psi : K \rightarrow B',$$

which is independent of  $m$ . Then, the natural Galois action of  $\tilde{G}_\infty$  on  $P_m$  is compatible with the action of  $\tilde{G}_\infty$  on  $e_m$  via  $\Psi$ , which was defined in Section 1.2. When

$$\tilde{\mathcal{L}}_{g,m} := \alpha_p^{-m} \sum_{\sigma \in \tilde{G}_m} g(\sigma \bar{P}_m) \cdot \sigma^{-1} \in \mathbb{Z}/p^n\mathbb{Z}[\tilde{G}_m],$$

it follows that the sequence  $\{\tilde{\mathcal{L}}_{g,m}\}$  defines an element of  $\mathbb{Z}/p^n\mathbb{Z}[[\tilde{G}_\infty]]$ , equal to  $\tilde{\mathcal{L}}_g$ . Define the local cohomology groups

$$H_{\text{fin}}^1(\tilde{K}_{m,\ell_2}, T_{f,n}) := \bigoplus_{\lambda \mid \ell_2} H_{\text{fin}}^1((\tilde{K}_m)_\lambda, T_{f,n}),$$

where the sum is taken over all the primes of  $\tilde{K}_m$  dividing  $\ell_2$ , and

$$\hat{H}_{\text{fin}}^1(\tilde{K}_{\infty,\ell_2}, T_{f,n}) := \varprojlim_{\bar{m}} H_{\text{fin}}^1(\tilde{K}_{m,\ell_2}, T_{f,n}),$$

where the inverse limit is taken with respect to the natural corestriction maps. The fixed identification of  $H_{\text{fin}}^1(K_{\ell_2}, T_{f,n})$  with  $\mathbb{Z}/p^n\mathbb{Z}$ , together with the choice

of the prime  $\lambda_\infty$ , yields the identifications

$$H_{\text{fin}}^1(\tilde{K}_{m,\ell_2}, T_{f,n}) = \mathbb{Z}/p^n[\tilde{G}_m], \quad \hat{H}_{\text{fin}}^1(\tilde{K}_{\infty,\ell_2}, T_{f,n}) = \mathbb{Z}/p^n[\tilde{G}_\infty].$$

In view of Corollary 9.4 and the definition of the map  $\gamma$ , the image of  $P_m^*$  in  $H_{\text{fin}}^1(\tilde{K}_{m,\ell_2}, T_{f,n})$  corresponds to  $\tilde{\mathcal{L}}_{g,m} \pmod{p^n}$ , and the image of the compatible sequence  $\{P_m^*\}$  in  $\hat{H}_{\text{fin}}^1(\tilde{K}_{\infty,\ell_2}, T_{f,n})$  corresponds to  $\tilde{\mathcal{L}}_g \pmod{p^n}$ , under the above identifications. Recall the class  $\tilde{\kappa}(\ell_1)$ , defined in Section 7 as the image of the sequence  $\{P_m^*\}$  in  $\hat{H}^1(\tilde{K}_\infty, T_{f,n})$  by the coboundary map. The value  $v_{\ell_2}(\tilde{\kappa}(\ell_1))$  at  $\ell_2$  of  $\tilde{\kappa}(\ell_1)$  is naturally an element of  $\hat{H}_{\text{fin}}^1(\tilde{K}_{\infty,\ell_2}, T_{f,n})$ , and is equal to the image of  $\{P_m^*\}$ , and hence to  $\tilde{\mathcal{L}}_g \pmod{p^n}$ . Since  $\mathcal{L}_g$  is the image in  $\Lambda$  of  $\tilde{\mathcal{L}}_g$ , and  $\kappa(\ell_1)$  is the corestriction from  $\tilde{K}_\infty$  to  $K_\infty$  of  $\tilde{\kappa}(\ell_1)$ , Theorem 4.2 follows.

*Remark.* The result proved in this section can be viewed as a generalization of the main result of [BD5]. The proof given here follows closely the approach in [Va2], avoiding the study of certain groups of connected components which was involved in the methods of [BD5].

UNIVERSITY OF MILANO, MILANO, ITALY  
E-mail address: Massimo.Bertolini@mat.unimi.it

MCGILL UNIVERSITY, MONTREAL, QC, CANADA  
E-mail address: darmon@math.mcgill.ca

#### REFERENCES

- [BC] J-F. BOUTOT and H. CARAYOL, Uniformisation  $p$ -adique des courbes de Shimura: les théorèmes de Cerednik et de Drinfeld, *Astérisque* **196–97** (1991), 45–158.
- [BD0] M. BERTOLINI and H. DARMON, Derived heights and generalized Mazur-Tate regulators, *Duke Math. J.* **76** (1994), 75–111.
- [BD $\frac{1}{2}$ ] ———, Derived  $p$ -adic heights, *Amer. J. Math.* **117** (1995), 1517–1554.
- [BD1] ———, Heegner points on Mumford-Tate curves, *Invent. Math.* **126** (1996), 413–456.
- [BD2] ———, A rigid-analytic Gross-Zagier formula and arithmetic applications, *Ann. of Math.* **146** (1997), 111–147.
- [BD3] ———, Heegner points,  $p$ -adic  $L$ -functions, and the Cerednik-Drinfeld uniformization, *Invent. Math.* **131** (1998), 453–491.
- [BD4] ———,  $p$ -adic periods,  $p$ -adic  $L$ -functions and the  $p$ -adic uniformization of Shimura curves, *Duke Math. J.* **98** (1999), 305–334.
- [BD5] ———, Euler systems and Jochnowitz congruences, *Amer. J. Math.* **121** (1999), 259–281.
- [BLR] S. BOSCH, W. LÜTKEBOHMERT, and M. RAYNAUD, *Néron Models*, *Ergebnisse der Mathematik und ihrer Grenzgebiete* **21**, Springer-Verlag, New York, 1990.
- [BoLeRi] N. BOSTON, H. LENSTRA, and K. RIBET, Quotients of groups rings arising from two-dimensional representations, *C. R. Acad. Sci. Paris* **312**, Série I (1991), 323–328.

- [Bu] K. BUZZARD, Integral models of certain Shimura curves, *Duke Math. J.* **87** (1998), 591–612.
- [Co] C. CORNUT, Réduction de familles de points CM, Ph.D. Thesis, Université Louis Pasteur, Strasbourg, 2000.
- [Dag] H. DAGHIGH, Modular forms, quaternion algebras, and special values of  $L$ -functions, Ph.D. thesis, McGill University, 1997.
- [Da] H. DARMON, A refined conjecture of Mazur-Tate type for Heegner points, *Invent. Math.* **110** (1992), 123–146.
- [DDT] H. DARMON, F. DIAMOND, and R. TAYLOR, Fermat's last theorem, in *Current Developments in Mathematics, 1995* (Cambridge, MA), 1–154, Internat. Press, Cambridge, MA, 1994.
- [DR] P. DELIGNE and M. RAPOPORT, *Les Schémas de modules des courbes elliptiques*, *Lecture Notes in Math.* **349**, Springer-Verlag, New York, 1973, 143–316.
- [Dr] V. G. DRINFELD, Coverings of  $p$ -adic symmetric regions (in Russian), *Funkts. Anal. Prilozn.* **10** (1976), 29–40; Transl. in *Funct. Anal. Appl.* **10** (1976), 107–115.
- [DT] F. DIAMOND and R. TAYLOR, Nonoptimal levels of mod  $\ell$  modular representations, *Invent. Math.* **115** (1994), 435–462.
- [Ed] B. EDIXHOVEN, Appendix in [BD2].
- [Ei] M. EICHLER, The basis problem for modular forms and the traces of the Hecke operators, in *Modular Functions of One Variable, I* (*Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972*), 75–151. *Lecture Notes in Math.* **320**, 75–151, Springer-Verlag, New York, 1973.
- [Gr1] B. H. GROSS, Heights and the special values of  $L$ -series, in *Number Theory* (Montreal, Que., 1985), 115–187, *CMS Conf. Proc.* **7**, A. M. S., Providence, RI, 1987.
- [Gr2] B. H. GROSS and D. B. ZAGIER, Heegner points and derivatives of  $L$ -series, *Invent. Math.* **84** (1986), 225–320.
- [Groth] A. GROTHENDIECK, *Groupes de Monodromie en Géométrie Algébrique, SGA VII*, *Lecture Notes in Math.* **288**, Springer-Verlag, New York, 1972.
- [GvdP] L. GERRITZEN and M. VAN DER PUT, *Schottky Groups and Mumford Curves*, *Lecture Notes in Math.* **817**, Springer-Verlag, New York, 1980.
- [I1] Y. IHARA, *On Congruence Monodromy Problems*, *Lecture Notes* **1**, Department of Mathematics, University of Tokyo, Tokyo, 1968.
- [I2] ———, Shimura curves over finite fields and their rational points, *Contemporary Math.* **245** (1999), 15–23.
- [JL] H. JACQUET and R. P. LANGLANDS, *Automorphic forms on  $GL(2)$* , *Lecture Notes in Math.* **114**, Springer-Verlag, New York, 1970.
- [JoLi1] B. W. JORDAN and R. LIVNÉ, Local Diophantine properties of Shimura curves, *Math. Ann.* **270** (1985), 235–248.
- [JoLi2] ———, On the Néron model of Jacobians of Shimura curves, *Compositio Math.* **60** (1986), 227–236.
- [JoLi3] ———, Integral Hodge theory and congruences between modular forms, *Duke Math. J.* **80** (1995), 419–484.
- [KM] N. KATZ and B. MAZUR, *Arithmetic Moduli of Elliptic Curves*, *Ann. of Math. Studies* **108**, Princeton Univ. Press, Princeton, NJ, 1985.
- [M] Y. I. MANIN,  $p$ -adic automorphic functions, *J. Soviet Math.* **5** (1976), 279–333.

- [Ma1] B. MAZUR, On the arithmetic of special values of  $L$  functions, *Invent. Math.* **55** (1979), 207–240.
- [MR] B. MAZUR and K. RUBIN, *Kolyvagin Systems*, *Mem. Amer. Math. Soc.* **168**, A. M. S., Providence, RI (2004).
- [MTT] B. MAZUR, J. TATE, and J. TEITELBAUM, On  $p$ -adic analogues of the conjectures of Birch and Swinnerton-Dyer, *Invent. Math.* **84** (1986), 1–48.
- [Ram] R. RAMAKRISHNA, Lifting Galois representations, *Invent. Math.* **138** (1999), 537–562.
- [Ray] M. RAYNAUD, Spécialisation du foncteur de Picard, *Publ. Math. Inst. Hautes Etudes Sc.* **38** (1970), 27–76.
- [Ri1] K. RIBET, Bimodules and abelian surfaces, *Adv. Stud. Pure Math.* **17** (1989), 359–407.
- [Ri2] ———, On modular representation of  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$  arising from modular forms, *Invent. Math.* **100** (1990), 431–476.
- [Ro] D. ROBERTS, Shimura curves analogous to  $X_0(N)$ , Ph.D. Thesis, Harvard University, 1989.
- [Ru] K. RUBIN, *Euler Systems*, *Ann. of Math. Studies* **147**, Princeton Univ. Press, Princeton, NJ, 2000.
- [Se1] J-P. SERRE, *Trees* (translated from the French by John Stilwell), Springer-Verlag, New York, 1980.
- [Se2] ———, *Abelian  $\ell$ -Adic Representations and Elliptic Curves*, Addison-Wesley Publ., Redwood City, CA, 1989.
- [Sh] G. SHIMURA, *Introduction to the Arithmetic Theory of Automorphic Functions* (reprint of the 1971 original), *Publication of the Mathematical Society of Japan* **11**, *Kanô Memorial Lectures* **1**, Princeton Univ. Press, Princeton, NJ, 1994.
- [Va1] V. VATSAL, Uniform distribution of Heegner points, *Invent. Math.* **148** (2002), 1–48.
- [Va2] ———, Special values of anticyclotomic  $L$ -functions, *Duke Math. J.* **116** (2003), 219–261.
- [Vi] M.-F. VIGNERAS, *Arithmétique des Algèbres des Quaternions*, *Lecture Notes in Math.* **800**, Springer-Verlag, New York, 1980.
- [Wa] W. WATERHOUSE, Abelian varieties over finite fields, *Ann. Sci. École Norm. Sup. Série* **4** (1969), 521–560.
- [W] A. WILES, Modular elliptic curves and Fermat’s last theorem, *Ann. of Math.* **141** (1995) 443–551.
- [Zh] S. ZHANG, Gross-Zagier formula for  $\text{GL}_2$ , *Asian J. Math.* **5** (2001), 183–290.

(Received June 19, 2001)