

189-235A: Basic Algebra I

Assignment 4

Due: Wednesday, October 6

1. Let R be a commutative ring. Is the set S of matrices of the form $\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}$, with $a, b, c \in R$, a subring of $M_2(R)$?
2. Same question as 1, but with S replaced by the set of matrices of the form $\begin{pmatrix} a & b \\ 0 & a \end{pmatrix}$, with $a, b \in R$.
3. Show that the ring \mathbf{Q} of rational numbers has no subrings which are finite sets.
4. Let A be a commutative ring, let $R = A \times A$, and let S be the subset of elements of R of the form $(a, 0)$, with $a \in A$. Show that S is a ring which is isomorphic to A . Show that S is *not* a subring of R . (According to the definition of subring given in class, which differs from the definition given in Hungerford's book for example.)
5. Let R be a ring, and let a be an element of R which is not a zero divisor. Show that the cancellation law can be applied to a , i.e., for all $x, y \in R$, if $ax = ay$ then $x = y$.
6. Let $R = \mathbf{Q}(\sqrt{2})$ be the ring of elements of the form $a + b\sqrt{2}$, with $a, b \in \mathbf{Q}$. Show that the function which sends $a + b\sqrt{2}$ to $a - b\sqrt{2}$ is an isomorphism from R to itself.
7. Let R be a ring. Show that there is a *unique* ring homomorphism f from \mathbf{Z} to R .
8. Given a ring R , let $R[i]$ denote the set of pairs (a, b) with $(a, b) \in R$.

Define an addition and multiplication on $R[i]$ by the rules:

$$(a_1, b_1) + (a_2, b_2) = (a_1 + a_2, b_1 + b_2).$$

$$(a_1, b_1)(a_2, b_2) = (a_1a_2 - b_1b_2, a_1b_2 + b_1a_2).$$

Show that these rules equip $R[i]$ with the structure of a ring.

9. Show that the subset S of the ring $R[i]$ of exercise 8 consisting of elements of the form $(r, 0)$ with $r \in R$ is a subring of $R[i]$ which is isomorphic to R . Produce an element i in $R[i]$ satisfying $i^2 = -1$.

10. Let \mathbf{R} be the field of real numbers. Retaining the notations of exercise 8, show that $\mathbf{R}[i]$ is isomorphic to the ring \mathbf{C} of complex numbers.

Optional questions.

11. Keeping the notations of exercise 8, show that the ring $\mathbf{C}[i]$ is isomorphic to $\mathbf{C} \times \mathbf{C}$.

12. Show that a ring R which is a finite set and an integral domain (has no zero-divisors) is necessarily a field.