

## MATH 556 - EXERCISES 3

### *Not for Assessment.*

1. Suppose  $X$  is a random variable, with mgf  $M_X(t)$  defined on  $(-h, h)$  for some  $h > 0$ . Show that

$$P_X[X \geq a] \leq e^{-at} M_X(t) \quad \text{for } 0 < t < \delta$$

For the *cumulant generating function*  $K_X(t) = \log M_X(t)$ , verify that

$$\frac{d}{dt} \{K_X(t)\}_{t=0} = \mathbb{E}_X[X] \qquad \frac{d^2}{dt^2} \{K_X(t)\}_{t=0} = \text{Var}_X[X]$$

2. The *non-central chi-square* distribution arises as the distribution of the square of a normal random variable. That is, if  $X \sim \text{Normal}(\mu, 1)$ , then  $Y = X^2$  has the non-central chi-square distribution with one degree of freedom and non-centrality parameter  $\lambda$ , denoted  $Y \sim \chi_\nu^2(\lambda)$ , where  $\nu = 1$  and  $\lambda = \mu^2$ . In this setting,

- (a) Find the pdf of  $Y$ , and show that it can be expressed in the form

$$f_Y(y) = e^{-\lambda/2} \sum_{j=0}^{\infty} \frac{(\lambda/2)^j}{j!} f_{Z_{2j+1}}(y) \quad y > 0$$

where  $f_{Z_m}$  is the pdf of a random variable  $Z_m$  which has a chi-square distribution with  $m$  degrees of freedom (that is,  $Z_m \sim \text{Gamma}(m/2, 1/2)$ ).

- (b) Find the characteristic function  $\varphi_Y(t)$ .  
 (c) Find the *Laplace transform*  $\mathcal{L}_Y(t)$ , defined for  $t \geq 0$  by

$$\mathcal{L}_Y(t) = \int_0^{\infty} e^{-ty} dF_Y(y) = \mathbb{E}_Y[e^{-tY}].$$

Note that  $\mathcal{L}_Y(t)$  is well-defined provided  $Y \geq 0$  with probability 1.

- (d) Find the expectation and variance of  $Y$ .  
 (e) Find the distribution of

$$S = \sum_{i=1}^n Y_i$$

where  $Y_1, \dots, Y_n$  are independent, with  $Y_i \sim \chi_{\nu_i}^2(\lambda_i)$ ,  $i = 1, \dots, n$ .

3. If  $\mathcal{L}_X(t)$  is the Laplace transform (see question above) of a nonnegative random variable  $X$ , show that for  $r = 1, 2, \dots$

$$(-1)^r \frac{d^r}{dt^r} \{\mathcal{L}_X(t)\} \geq 0 \quad t \geq 0.$$

If  $F_X$  is the corresponding cdf, show that

$$\mathcal{L}_X(t) = t \int_0^{\infty} \exp\{-tx\} F_X(x) dx.$$

4. Suppose that  $X_1 \sim \text{Gamma}(\alpha_1, \beta_1)$  and  $X_2 \sim \text{Gamma}(\alpha_2, \beta_2)$  are independent random variables. Characterize the distribution of  $Y = X_1 - X_2$ .

5. Suppose that  $\{\varphi_k(t)\}_{k=1}^n$  is a sequence of characteristic functions, and  $\{c_k\}_{k=1}^n$  is a sequence of non-negative real valued constants, with

$$\sum_{k=1}^n c_k = 1.$$

Show that

$$\sum_{k=1}^n c_k \varphi_k(t)$$

is also a characteristic function, and identify the distribution to which it corresponds. Does the result extend to the case where  $n \rightarrow \infty$ ? Justify your answer.

6. If

$$\varphi_1(t) = \exp(-4t^2) \quad \varphi_2(t) = (3 + \cos(t) + \cos(2t))/5$$

identify the distribution with cf

$$\frac{\varphi_1(t) + \varphi_2(t)}{2}.$$

7. Suppose  $X_1$  and  $X_2$  are independent random variables, and suppose also that  $X_1$  and  $X_1 - X_2$  are independent. Show that

$$P_{X_1}[X_1 = c] = 1$$

for some constant  $c$ .

*Hint: write  $X_2 = X_1 + (X_2 - X_1)$ , and recall that if  $\varphi(t)$  is an arbitrary cf, then  $\varphi(t)$  is continuous for all  $t$ .*

8. Suppose that mgf  $M_X(t)$  is defined, for a suitable neighbourhood of zero  $(-h, h)$ , as

$$M_X(t) = \frac{9e^{-t}}{(3 + 2t)^2}.$$

Find an expression for  $\mathbb{E}_X[X^r]$ , for  $r = 1, 2, \dots$ .

9. Suppose that  $X \sim \text{Binomial}(n, \theta)$  for integer  $n \geq 1$ , and  $0 < \theta < 1$ . Let

$$Z_n = \frac{(X - n\theta)}{\sqrt{n\theta(1 - \theta)}}.$$

Find the first two non-zero terms in the power series expansion of the cumulant generating function of  $Z_n$ , and the order of approximation (in terms of  $n$ ) when truncating the expansion at the second term, for large  $n$ .

Recall that

$$\log\{(1 + z)^n\} = n\{z - z^2/2 + \dots\}.$$