

ASSIGNMENT 3 - MATH 251, WINTER 2008

Submit by Monday, February 4, 12:00

- (1) Read §4.5 in the notes and write a proof for Proposition 4.5.2.
- (2) **Coding theory.** Let $V = \mathbb{F}_2^n$ and let C be a code (= a subspace) of dimension k . Let d be the distance of C . Prove that $d \leq n - k + 1$.
- (3) **Coding theory.** Let G be a bipartite regular graph, whose set of left vertices is L and right vertices is R . For a set $S \subset L$ denote by $\partial S := \{r \in R : r \text{ is connected to a vertex in } S\}$. Suppose that $|L| = n$, $|R| = 3n/4$ and n is “large”.

Suppose that G has the following expansion property. For every $S \subset L$ such that $|S| \leq \frac{n}{10d_L}$ we have $|\partial S| \geq \frac{5d_L}{8}|S|$. Prove that for every such S there is a vertex r_S (many, in fact) such that r_S is a neighbor of exactly one element of S .

Consider now the linear code defined by the “half adjacency matrix M ” whose columns are indexed by the elements of L and rows by the elements of R , having 1 as an entry if the corresponding vertices are connected and 0 otherwise. (Refer to the previous assignment). Prove that if x is a non-zero vector in the code then x has more than $\frac{n}{10d_L}$ non-zero coordinates. Conclude that we get a code where the distance between any two code words is at least $\frac{n}{10d_L}$ and whose rate is at least $1/4$.

- (4) (a) Find a linear map $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ whose image is generated by $(1, 1, 0)$ and $(0, 1, 1)$. Here ‘find’ means represent by a matrix with respect to the standard basis.
(b) Find a linear map $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ whose kernel is generated by $(1, 2, 0, 4), (0, 1, 0, 1)$.
- (5) Let $W = \{(x, y, z, w) : x + y + z + w = 0, x - y - w = 0\}$, $U_1 = \{(x, x, x, x) : x \in \mathbb{R}\}$ be subspaces of \mathbb{R}^4 . Find a subspace $U \supset U_1$ such that $\mathbb{R}^4 = U \oplus W$. Let T be the projection of \mathbb{R}^4 on U along W . Write the matrix representing T with respect to the standard basis.
- (6) Let V and W be any two finite dimensional vector spaces over a field \mathbb{F} . Let $T : V \rightarrow W$ be a linear map. Prove that there are bases of V and W such that with respect to those bases T is represented by a matrix composed of 0’s and 1’s only. If T is an isomorphism, prove that with respect to suitable bases it is represented by the identity matrix.