

On Dirichlet series for sums of squares

Kwok-Kwong Stephen Choi (choi@cecm.sfu.ca)

Simon Fraser University

Department of Mathematics

8888 University Drive

Burnaby, BC V5A 1S6

Canada

Abstract. In their famous book on the theory of numbers, Hardy and Wright recorded elegant closed forms for the generating functions of the divisor functions $\sigma_k(n)$ and $\sigma_k^2(n)$ in the terms of Riemann Zeta function $\zeta(s)$ only. In this talk, we explore other arithmetical functions enjoying this remarkable property. We are able to generalize this result and prove that if f_i and g_i are completely multiplicative, then we have

$$\sum_{n=1}^{\infty} \frac{(f_1 * g_1)(n) \cdot (f_2 * g_2)(n)}{n^s} = \frac{L_{f_1 f_2}(s) L_{g_1 g_2}(s) L_{f_1 g_2}(s) L_{g_1 f_2}(s)}{L_{f_1 f_2 g_1 g_2}(2s)}$$

where $L_f(s) := \sum_{n=1}^{\infty} f(n)n^{-s}$ is the Dirichlet series corresponding to f .

Let $r_N(n)$ be the number of solutions of $x_1^2 + \cdots + x_N^2 = n$ and $r_{2,P}(n)$ be the number of solutions of $x^2 + Py^2 = n$. One of the applications of our theorem above is to obtain closed forms, in terms of $\zeta(s)$ and Dirichlet L -functions, for the generating functions of $r_N(n)$, $r_N^2(n)$, $r_{2,P}(n)$ and $r_{2,P}(n)^2$ for certain N and P . We also use these generating functions to obtain asymptotic estimates of the average values for each function for which we obtain a Dirichlet series. This is a joint work with Jonathan Borwein.

