

Do any 8 of the following 10 problems. Every problem are worth 10 points.

Problem 1.

- Let X be a metric space with the distance d_1 . Prove that $d_2(x, y) = d_1(x, y)/(1 + d_1(x, y))$ also defines the distance on X . Prove that open sets and Cauchy sequences for d_1 and d_2 coincide.
- Prove the same results for $d_3(x, y) = \min\{d_1(x, y), 1\}$.
- Define the distance on the set X of all sequences $x = (x_1, x_2, \dots)$ of real numbers by the formula

$$d(x, y) = \sum_{k=1}^{\infty} \frac{|x_k - y_k|}{2^k(1 + |x_k - y_k|)}.$$

Prove that d defines a distance on X , and that X is complete with respect to d . Is X separable?

Problem 2. Let $X = \prod_i X_i$ be a product of metric spaces, with the distance d_i . Consider the product topology on X (a basis of open sets is given by $\prod_i U_i$, where $U_i = X_i$ except for finitely many i -s). Let $\rho_i = d_i/(1 + d_i)$; it preserves the topology of X_i by Problem 1a. Prove that

$$\rho(x, y) = \sum_{j=1}^{\infty} \frac{\rho_j(x_j, y_j)}{2^j}$$

defines a distance on X , and that the topology given by ρ coincides with the product topology. Hint: Let U be an open set in the basis for the product topology, and let $x \in U$. Prove that there exists $r > 0$ s.t. $B_\rho(x, r) \subset U$. Conversely, let $y \in B_\rho(x, r)$. Prove that there exists a basis set U for the product topology s.t. $y \in U \subset B_\rho(x, r)$.

Problem 3. Let $C^\infty[a, b]$ denote the space of infinitely differentiable functions on $[a, b]$ (all the derivatives exist and are continuous). Let

$$d(f, g) = \sum_{k=1}^{\infty} \frac{\max_{x \in [a, b]} |f^{(k)}(x) - g^{(k)}(x)|}{2^k(1 + \max_{x \in [a, b]} |f^{(k)}(x) - g^{(k)}(x)|)}.$$

Prove that d defines the distance on $C^\infty[a, b]$, and that the resulting metric space is complete.

Problem 4. Hausdorff distance. Let $A, B \subset X$, where X is a metric space. Let $U_r(Z) := \{x \in X : d(x, Z) \leq r\}$. Define the *Hausdorff distance*

$$d_H(A, B) = \inf\{r > 0 : A \subset U_r(B), \text{ and } B \subset U_r(A)\}.$$

- Prove that $d_H \geq 0$, is symmetric and satisfies the triangle inequality.
- Prove that $d_H = \max(\sup_{a \in A} d(a, B), \sup_{b \in B} d(b, A))$.
- Prove that $d_H(A, B) \leq r$ if and only if $d(a, B) \leq r$, for all $a \in A$ and $d(b, A) \leq r$ for all $b \in B$.
- $d_H(A, \bar{A}) = 0$, where \bar{A} is the closure of A .

e) Show that if A, B are closed subsets of X , and $d_H(A, B) = 0$, then $A = B$.

According to e), d_H defines a distance on the set $\mathcal{M}(X)$ of all *closed* subsets of X .

Problem 5. Let $A_n \in \mathcal{M}(X)$ be a sequence of closed subsets of X , and let $d_H(A_n, A) \rightarrow 0$ as $n \rightarrow \infty$, i.e. let $A_n \rightarrow A$ in the metric space $(\mathcal{M}(X), d_H)$. Prove that

a) A is the set of limits of all converging subsequences $\{a_n\}$ in X , s.t. $a_n \in A_n$ for all n .

b) $A = \bigcap_{n=1}^{\infty} (\text{closure of } \bigcup_{m=n}^{\infty} A_m)$.

Next, let X be compact, and $\{A_i\}$ be a sequence of its compact subspaces. Prove that

c) If $A_{i+1} \subset A_i$ for all i , then $A_k \rightarrow \bigcap_{k=1}^{\infty} A_k$ in $\mathcal{M}(X)$.

d) If $A_i \subset A_{i+1}$ for all i , then A_i converges to the closure of $\bigcup_{i=1}^{\infty} A_i$.

Problem 6. Consider the orthogonal group $O(n)$ consisting of all $n \times n$ orthogonal matrices, i.e. matrices whose columns form an orthonormal basis v_1, \dots, v_n of \mathbf{R}^n . We introduce the topology on $O(n)$ by considering it as a subspace of \mathbf{R}^{n^2} (consider the matrix entries as coordinates). Show that

a) $O(n)$ is a closed subset of $\text{Mat}_n(\mathbf{R}) \simeq \mathbf{R}^{n^2}$, by considering the dot products (v_i, v_j) of the columns of matrices in $O(n)$ as functions from $\text{Mat}_n(\mathbf{R})$ into \mathbf{R} .

b) Show that $O(n)$ is compact.

c) Prove that $O(n)$ is a group, i.e. if $A, B \in O(n)$, then $AB \in O(n)$, and $A^{-1} \in O(n)$.

Problem 7. Let $X = C^m[0, 1]$ denote the space of m times continuously differentiable functions on $[0, 1]$. Define the norm on X by

$$\|f\| = \sum_{k=0}^m \max_{x \in [0, 1]} |f^{(k)}(x)|.$$

Prove that $(X, \|\cdot\|)$ is a complete metric space. Is it separable?

Problem 8.

a) Compute the area $A(r)$ of the ball of radius r in \mathbf{R}^2 , S^2 , and \mathbf{H}^2 . Hint: the volume element in polar coordinates (r, θ) is given by $rdrd\theta$ in \mathbf{R}^2 ; $\sin rdrd\theta$ on S^2 ; and $\sinh rdrd\theta$ in \mathbf{H}^2 . Where does the volume grow faster? Compute the first 3 terms in the Taylor series expansion of the volume as $r \rightarrow 0$; what do you get?

b) Next, compute the length $L(r)$ of the circle of radius r in \mathbf{R}^2 , S^2 , and \mathbf{H}^2 . Hint: the length element in polar coordinates (r, θ) is given by $dr^2 + r^2d\theta^2$ in \mathbf{R}^2 ; $dr^2 + \sin^2 rd\theta^2$ on S^2 ; and $dr^2 + \sinh^2 rd\theta^2$ in \mathbf{H}^2 .

c) Describe the behavior of the ratio $A(r)/L(r)$ as $r \rightarrow 0$.

d) Describe the behavior of the ratio $L(r)/A(r)$ as $r \rightarrow \infty$ in \mathbf{R}^2 and \mathbf{H}^2 ; and as $r \rightarrow \pi$ in S^2 .

Problem 9.

a) Compute $L(r)$ and $A(r)$ on an infinite k -regular tree, $k \geq 2$. Describe the behavior of the ratio $L(r)/A(r)$ as $r \rightarrow \infty$.

b) Do the same for the graph \mathbf{Z}^2 .

Problem 10. Every real number in $x \in [0, 1]$ can be expanded into a (finite or infinite) *continued fraction*

$$x = \frac{1}{n_1 + \frac{1}{n_2 + \frac{1}{n_3 + \dots}}},$$

sometimes denoted by $x = [n_1, n_2, n_3, \dots]$.

- a) Prove that finite continued fractions correspond to rational numbers, while infinite fractions correspond to irrational numbers.
- b) Prove that the function f in part a) can be written as a *shift map*,

$$f([n_1, n_2, n_3, \dots]) = [n_2, n_3, \dots].$$

- c) Describe all the *periodic* continued fractions, $x = [n_1, \dots, n_k, n_1, \dots, n_k, \dots]$.