

COHOMOLOGICAL DIMENSION AND SCHREIER'S FORMULA IN GALOIS COHOMOLOGY

JOHN LABUTE, NICOLE LEMIRE[†], JÁN MINÁČ[‡], AND JOHN SWALLOW

ABSTRACT. Let p be a prime and F a field containing a primitive p th root of unity. Then for $n \in \mathbb{N}$, the cohomological dimension of the maximal pro- p -quotient G of the absolute Galois group of F is $\leq n$ if and only if the corestriction maps $H^n(H, \mathbb{F}_p) \rightarrow H^n(G, \mathbb{F}_p)$ are surjective for all open subgroups H of index p . Using this result we derive a surprising generalization to $\dim_{\mathbb{F}_p} H^n(H, \mathbb{F}_p)$ of Schreier's formula for $\dim_{\mathbb{F}_p} H^1(H, \mathbb{F}_p)$.

For a prime p , let $F(p)$ denote the maximal p -extension of a field F . One of the fundamental questions in the Galois theory of p -extensions is to discover useful interpretations of the cohomological dimension $\text{cd}(G)$ of the Galois group $G = \text{Gal}(F(p)/F)$ in terms of the arithmetic of p -extensions of F . When $\text{cd}(G) = 1$, for instance, we know that G is a free pro- p -group [S1, §3.4], and when $\text{cd}(G) = 2$ we have important information on the G -module of relations in a minimal presentation [K, §7.3].

For a fixed $n > 2$, however, little is known about the structure of p -extensions when $\text{cd}(G) = n$. Now when $n = 1$ and G is finitely generated as a pro- p -group, we have Schreier's well-known formula

$$(1) \quad h_1(H) = 1 + [G : H](h_1(G) - 1)$$

for each open subgroup H of G , where

$$h_1(H) := \dim_{\mathbb{F}_p} H^1(H, \mathbb{F}_p).$$

(See, for instance, [K, Example 6.3].)

Date: May 10, 2005.

2000 Mathematics Subject Classification. Primary 12G05, 12G10.

Key words and phrases. cohomological dimension, Schreier's formula, Galois theory, p -extensions, pro- p -groups.

[†]Research supported in part by NSERC grant R3276A01.

[‡]Research supported in part by NSERC grant R0370A01, by the Mathematical Sciences Research Institute, Berkeley, and by a 2004/2005 Distinguished Research Professorship at the University of Western Ontario.

Observe that from basic properties of p -groups it follows that for each open subgroup H of G there exists a chain of subgroups

$$G = G_0 \supset G_1 \supset \cdots \supset G_k = H$$

such that G_{i+1} is normal in G_i and $[G_i : G_{i+1}] = p$ for each $i = 0, 1, \dots, k-1$. Since closed subgroups of free pro- p -groups are free [S1, Corollary 3, §I.4.2], Schreier's formula (1) is equivalent to the seemingly weaker statement that the formula holds for all open subgroups H of G of index p :

$$(2) \quad h_1(H) = 1 + p(h_1(G) - 1).$$

Using the Bloch-Kato Conjecture (see [V1], [V2], [Su], and [R]), we deduce a remarkable generalization of Schreier's formula for each $n \in \mathbb{N}$, as follows. Let F^\times denote the nonzero elements of a field F , and for $c \in F^\times$, let $(c) \in H^1(G, \mathbb{F}_p)$ denote the corresponding class. For $\alpha \in H^n(G, \mathbb{F}_p)$ abbreviate by $\text{ann}_n \alpha$ the annihilator

$$\text{ann}_n \alpha = \{\beta \in H^n(G, \mathbb{F}_p) \mid \alpha \cup \beta = 0\}.$$

Finally, set $h_n(G) = \dim_{\mathbb{F}_p} H^n(G, \mathbb{F}_p)$. Observe that the hypothesis on the surjectivity of the corestriction on degree n cohomology holds for all pro- p -groups of cohomological dimension n [NSW, Proposition 3.3.8]; conversely, in section 1, we show that this hypothesis for all subgroups of index p implies $\text{cd}(G) \leq n$.

We denote by ξ_p a p th root of unity of order p .

Theorem 1. *Suppose that $\xi_p \in F$, $\text{cd}(G) = n$, and $h_n(G) < \infty$. Let H be an open subgroup of G of index p , with fixed field $F(\sqrt[p]{a})$, and suppose that the corestriction map $H^n(H, \mathbb{F}_p) \rightarrow H^n(G, \mathbb{F}_p)$ is surjective. Then*

$$h_n(H) = a_{n-1}(G, H) + p(h_n(G) - a_{n-1}(G, H)),$$

where $a_{n-1}(G, H)$ is the codimension of $\text{ann}_{n-1}(a)$:

$$a_{n-1}(G, H) := \dim_{\mathbb{F}_p} \left(H^{n-1}(G, \mathbb{F}_p) / \text{ann}_{n-1}(a) \right).$$

The proof of Theorem 1 brings additional insight into the structure of Schreier's formula; in fact, it makes Schreier's formula transparent for any $n \in \mathbb{N}$. In section 1, we derive several interpretations for the statement $\text{cd}(G) = n$. First, we prove in Theorem 2 that if F contains a primitive p th root of unity ξ_p then $\text{cd}(G) \leq n$ if and only if the corestriction maps $\text{cor} : H^n(H, \mathbb{F}_p) \rightarrow H^n(G, \mathbb{F}_p)$ are surjective for all open subgroups H of G of index p . As a corollary, we show that the

corresponding cohomology groups $H^{n+1}(H, \mathbb{F}_p)$ are all free as $\mathbb{F}_p[G/H]$ -modules if and only if $\text{cd}(G) \leq n$, under the additional hypothesis that $F = F^2 + F^2$ when $p = 2$. Finally, we show in Theorem 3 that if G is finitely generated, then $\text{cd}(G) \leq n$ if and only if a single corestriction map, from the Frattini subgroup $\Phi(G) = G^p[G, G]$ of G , is surjective. In section 2 we prove Theorem 1.

For basic facts about Galois cohomology and maximal p -extensions of fields, we refer to [K] and [S1]. In particular, we work in the category of pro- p -groups.

1. WHEN IS $\text{cd}(G) = n$?

From the Bloch-Kato Conjecture ([V1], [V2], [Su], and [R]), we have the following interesting translation of the statement $\text{cd}(G) \leq n$ for a given $n \in \mathbb{N}$. Observe that when $\text{cd}(G) \leq n$, the corestriction maps $\text{cor} : H^n(H, \mathbb{F}_p) \rightarrow H^n(G, \mathbb{F}_p)$ are surjective for all open subgroups H of G of index p [NSW, Proposition 3.3.8].

Theorem 2. *Suppose that $\xi_p \in F$. Then for each $n \in \mathbb{N}$ we have $\text{cd}(G) \leq n$ if and only if*

$$\text{cor} : H^n(H, \mathbb{F}_p) \rightarrow H^n(G, \mathbb{F}_p)$$

is surjective for every open subgroup H of G of index p .

Proof. Suppose that F satisfies the conditions of the theorem, and let $G_{F(p)}$ be the absolute Galois group of $F(p)$.

Observe that since F contains ξ_p , the maximal p -extension $F(p)$ is closed under taking p th roots and hence $H^1(G_{F(p)}, \mathbb{F}_p) = \{0\}$. By the Bloch-Kato Conjecture [V2, Theorem 7.1], the subring of the cohomology ring $H^*(G_{F(p)}, \mathbb{F}_p)$ consisting of elements of positive degree is generated by cup-products of elements in $H^1(G_{F(p)}, \mathbb{F}_p)$. Hence $H^n(G_{F(p)}, \mathbb{F}_p) = \{0\}$ for $n \in \mathbb{N}$. Then, considering the Lyndon-Hochschild-Serre spectral sequence associated to the exact sequence

$$1 \rightarrow G_{F(p)} \rightarrow G_F \rightarrow G \rightarrow 1,$$

we have that

$$(3) \quad \text{inf} : H^*(G, \mathbb{F}_p) \rightarrow H^*(G_F, \mathbb{F}_p)$$

is an isomorphism.

Now suppose that $\text{cor} : H^n(H, \mathbb{F}_p) \rightarrow H^n(G, \mathbb{F}_p)$ is surjective for all open subgroups H of G of index p . Let K be the fixed field of such a subgroup H . Then $K = F(\sqrt[p]{a})$ for some $a \in F^\times$. From [V1, Lemma 6.11 and §7] and [V2, §6 and Theorem 7.1], as well as [V1, Proposition 5.2], modified in [LMS1, Theorem 5] and translated to G from G_F via the inflation maps (3) above, we obtain the following exact sequence:

$$(4) \quad H^n(H, \mathbb{F}_p) \xrightarrow{\text{cor}} H^n(G, \mathbb{F}_p) \xrightarrow{-\cup(a)} H^{n+1}(G, \mathbb{F}_p) \xrightarrow{\text{res}} H^{n+1}(H, \mathbb{F}_p).$$

Therefore $\text{res} : H^{n+1}(G, \mathbb{F}_p) \rightarrow H^{n+1}(H, \mathbb{F}_p)$ is injective for every open subgroup H of G of index p .

Now consider an arbitrary element

$$\alpha = (a_1) \cup \cdots \cup (a_{n+1}) \in H^{n+1}(G, \mathbb{F}_p),$$

where $a_i \in F^\times$ and (a_i) is the element of $H^1(G, \mathbb{F}_p)$ associated to a_i , $i = 1, 2, \dots, n+1$. Suppose that $(a_1) \neq 0$, and set $K = F(\sqrt[p]{a_1})$ and $H = \text{Gal}(F(p)/K)$. We have $0 = \text{res}(\alpha) \in H^{n+1}(H, \mathbb{F}_p)$. Since res is injective, $\alpha = 0$. Again by the Bloch-Kato conjecture [V1, Theorem 7.1], we know that $H^{n+1}(G, \mathbb{F}_p)$ is generated by the elements α above. Hence $H^{n+1}(G, \mathbb{F}_p) = \{0\}$ and therefore $\text{cd}(G) \leq n$. (See [K, page 49].)

Conversely, if $\text{cd}(G) \leq n$ then by [NSW, Proposition 3.3.8] we conclude that $\text{cor} : H^n(H, \mathbb{F}_p) \rightarrow H^n(G, \mathbb{F}_p)$ is surjective for open subgroups H of G of index p . \square

Using conditions obtained in [LMS2] for $H^n(H, \mathbb{F}_p)$ to be a free $\mathbb{F}_p[G/H]$ -module, we obtain the following corollary. We observe the convention that $\{0\}$ is a free $\mathbb{F}_p[G/H]$ -module.

Corollary. *Suppose that $\xi_p \in F$ and if $p = 2$ suppose also that $F = F^2 + F^2$. Then for each $n \in \mathbb{N}$, we have that $H^{n+1}(H, \mathbb{F}_p)$ is a free $\mathbb{F}_p[G/H]$ -module for every open subgroup H of G of index p if and only if $\text{cd}(G) \leq n$.*

Observe that the condition $F = F^2 + F^2$ is satisfied in particular when F contains a primitive fourth root of unity i : for all $c \in F^\times$, $c = ((c+1)/2)^2 + ((c-1)i/2)^2$.

Proof. Assume that F is as above, $n \in \mathbb{N}$, and that $H^{n+1}(H, \mathbb{F}_p)$ is a free $\mathbb{F}_p[G/H]$ -module for every open subgroup H of G of index p . If

$p > 2$, then it follows from [LMS2, Theorem 1] that the corestriction maps $\text{cor} : H^n(H, \mathbb{F}_p) \rightarrow H^n(G, \mathbb{F}_p)$ are surjective for all such subgroups H .

If $p = 2$, then we consider open subgroups H of index 2 with corresponding fixed fields $K = F(\sqrt{a})$. From [LMS2, Theorem 1] we obtain that $\text{ann}_n(a) = \text{ann}_n((a) \cup (-1))$. It follows from the hypothesis $F = F^2 + F^2$ that $(c) \cup (-1) = 0 \in H^2(G, \mathbb{F}_2)$ for each $c \in F^\times$ and in particular for $c = a$. Hence $\text{ann}_n(a) = H^n(G, \mathbb{F}_2)$. But then from exact sequence (4) above, we deduce that $\text{cor} : H^n(H, \mathbb{F}_2) \rightarrow H^n(G, \mathbb{F}_2)$ is surjective.

Since our analysis holds for all open subgroups H of index p , by Theorem 2 we conclude that $\text{cd}(G) \leq n$.

Assume now that $\text{cd}(G) \leq n$. Then by Serre's theorem in [S2] we find that $\text{cd}(H) \leq n$ for every open subgroup H of G . Hence $H^{n+1}(H, \mathbb{F}_p) = \{0\}$ which, by our convention, is a free $\mathbb{F}_p[G/H]$ -module, as required. \square

Remark. When $p = 2$ and $F \neq F^2 + F^2$, the statement of the corollary may fail. Consider the case $F = \mathbb{R}$. Then the only subgroup H of index 2 in $G = \mathbb{Z}/2\mathbb{Z}$ is $H = \{1\}$. Then for all $n \in \mathbb{N}$, $H^{n+1}(H, \mathbb{F}_2) = \{0\}$ and is free as an $\mathbb{F}_2[G/H]$ -module. However, $\text{cd}(G) = \infty$.

Under the additional assumption that G is finitely generated, we show that the surjectivity of a single corestriction map is equivalent to $\text{cd}(G) \leq n$.

Theorem 3. *Suppose that $\xi_p \in F$ and G is finitely generated. Then for each $n \in \mathbb{N}$ we have $\text{cd}(G) \leq n$ if and only if*

$$\text{cor} : H^n(\Phi(G), \mathbb{F}_p) \rightarrow H^n(G, \mathbb{F}_p)$$

is surjective.

Proof. Because G is finitely generated, the index $[G : \Phi(G)]$ is finite, and we may consider a suitable chain of open subgroups

$$G = G_0 \supset G_1 \supset \cdots \supset G_k = \Phi(G)$$

such that $[G_i : G_{i+1}] = p$ for each $i = 0, 1, \dots, k-1$.

By Serre's theorem in [S2], $\text{cd}(H) = \text{cd}(G)$ for every open subgroup H of G . Hence if $\text{cd}(G) \leq n$ we may iteratively apply Theorem 2 to

the chain of open subgroups to conclude that

$$\text{cor} : H^n(\Phi(G), \mathbb{F}_p) \rightarrow H^n(G, \mathbb{F}_p)$$

is surjective.

Assume now that $\text{cor} : H^n(\Phi(G), \mathbb{F}_p) \rightarrow H^n(G, \mathbb{F}_p)$ is surjective. For each open subgroup H of G of index p we have a commutative diagram of corestriction maps

$$\begin{array}{ccc} H^n(\Phi(G), \mathbb{F}_p) & \longrightarrow & H^n(H, \mathbb{F}_p) \\ & \searrow & \downarrow \\ & & H^n(G, \mathbb{F}_p) \end{array}$$

since $\Phi(G) \subset H$. We obtain that $\text{cor} : H^n(H, \mathbb{F}_p) \rightarrow H^n(G, \mathbb{F}_p)$ is surjective, and by Theorem 2 we deduce that $\text{cd}(G) \leq n$, as required. \square

2. SCHREIER'S FORMULA FOR H^n

We now prove Theorem 1. Suppose that H is an open subgroup of G of index p and the corestriction map $\text{cor} : H^n(H, \mathbb{F}_p) \rightarrow H^n(G, \mathbb{F}_p)$ is surjective. Let $K = F(\sqrt[p]{a})$ be the fixed field of H .

We claim that $\text{ann}_{n-1}((a) \cup (\xi_p)) = H^{n-1}(G, \mathbb{F}_p)$. Suppose that $\alpha \in H^{n-1}(G, \mathbb{F}_p)$. By the surjectivity hypothesis there exists $\beta \in H^n(H, \mathbb{F}_p)$ such that $\text{cor} \beta = (\xi_p) \cup \alpha$. From [V1, Proposition 5.2] modified in [LMS1, Theorem 5], $(a) \cup (\text{cor} \beta) = 0$ and hence $(a) \cup (\xi_p) \cup \alpha = 0$. Therefore the claim is established. By [LMS1, Theorem 1], we obtain the decomposition

$$H^n(H, \mathbb{F}_p) = X \oplus Y,$$

where X is a trivial $\mathbb{F}_p[G/H]$ -module and Y is a free $\mathbb{F}_p[G/H]$ -module. (Because $\text{ann}_{n-1}((a) \cup (\xi_p)) = H^{n-1}(G, \mathbb{F}_p)$ there are no 2-dimensional summands, and by the surjectivity of the corestriction map, the fourth summand Z in [LMS1], a trivial $\mathbb{F}_p[G/H]$ -module, is also $\{0\}$.) Moreover

$$x := \dim_{\mathbb{F}_p} X = \dim_{\mathbb{F}_p} H^{n-1}(G, \mathbb{F}_p) / \text{ann}_{n-1}(a) = a_{n-1}(G, H), \quad \text{and}$$

$$y := \text{rank } Y = \dim_{\mathbb{F}_p} H^n(G, \mathbb{F}_p) / (a) \cup H^{n-1}(G, \mathbb{F}_p).$$

Therefore $h_n(H) = \dim_{\mathbb{F}_p} H^n(H, \mathbb{F}_p) = x + py$.

Now, considering the exact sequence

$$0 \rightarrow \frac{H^{n-1}(G, \mathbb{F}_p)}{\text{ann}_{n-1}(a)} \xrightarrow{-\cup(a)} H^n(G, \mathbb{F}_p) \rightarrow \frac{H^n(G, \mathbb{F}_p)}{(a) \cup H^{n-1}(G, \mathbb{F}_p)} \rightarrow 0,$$

we see that $\dim_{\mathbb{F}_p} H^n(H, \mathbb{F}_p)$ is equal to the sum of the dimension x of the kernel and p times the dimension y of the cokernel, and the theorem follows.

Observe that our formula $h_n(H) = x + py$ holds without the assumption that $h_n(G)$ is finite. (This assumption is used only in the formulation of Theorem 1 where we subtract $a_{n-1}(G, H)$ from $h_n(G)$.)

When $n = 1$, $\text{ann}_{n-1}(a) = \{0\}$ so that $a_{n-1}(G, H) = 1$. Therefore when G is finitely generated we recover Schreier's formula (2):

$$h_1(H) = 1 + p(h_1(G) - 1).$$

REFERENCES

- [K] H. Koch, *Galois theory of p -extensions*. Berlin: Springer-Verlag, 2002.
- [LMS1] N. Lemire, J. Mináč, and J. Swallow. Galois module structure of Galois cohomology. ArXiv:math.NT/0409484 (2004).
- [LMS2] N. Lemire, J. Mináč, and J. Swallow. When is Galois cohomology free or trivial? ArXiv:math.NT/0410617 (2004).
- [NSW] J. Neukirch, A. Schmidt, and K. Wingberg. *Cohomology of number fields*. Berlin: Springer-Verlag, 2000.
- [R] M. Rost. Norm varieties and the chain lemma. Lectures given at the Institute for Advanced Study, Princeton, January–March 2005. Notes by C. Haesemeyer. www.mathematik.uni-bielefeld.de/~rost/snv.html.
- [S1] J.-P. Serre. *Galois cohomology*. Heidelberg: Springer-Verlag, 1997.
- [S2] J.-P. Serre. Sur la dimension cohomologique des groupes profinis. *Topology* **3** (1965), 413–420.
- [Su] A. Suslin. Norm varieties. Lectures given at the Institute for Advanced Study, Princeton, October–December 2004. Notes by S. Joukhovitski. www.mathematik.uni-bielefeld.de/~rost/snv.html.
- [V1] V. Voevodsky. Motivic cohomology with $\mathbb{Z}/2$ -coefficients. *Publ. Inst. Hautes Études Sci.*, No. 98 (2003), 59–104.
- [V2] V. Voevodsky. On motivic cohomology with \mathbb{Z}/l coefficients. K-theory preprint archive 639 (2003). www.math.uiuc.edu/K-theory/0639/.

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCGILL UNIVERSITY,
BURNSIDE HALL, 805 SHERBROOKE STREET WEST, MONTREAL, QUEBEC
H3A 2K6 CANADA

E-mail address: labute@math.mcgill.ca

DEPARTMENT OF MATHEMATICS, MIDDLESEX COLLEGE, UNIVERSITY OF
WESTERN ONTARIO, LONDON, ONTARIO N6A 5B7 CANADA

E-mail address: nlemire@uwo.ca

E-mail address: minac@uwo.ca

DEPARTMENT OF MATHEMATICS, DAVIDSON COLLEGE, BOX 7046, DAVIDSON,
NORTH CAROLINA 28035-7046 USA

E-mail address: joswallow@davidson.edu