



Cal I (S) (Maths 201–NYA)

(Marks)

**Justify** all your answers—just having the correct answer is not sufficient.

Pace yourself—a rough guide is to spend less than  $2m$  minutes or so on a question worth  $m$  marks.


- (6) 1. For a continuous function  $y = f(x)$ , suppose you know its derivative is positive for  $-\infty < x < 1$  and for  $3 < x < +\infty$ , and that its derivative is negative for  $1 < x < 3$ . Furthermore suppose you know that its second derivative is positive for  $-\infty < x < 0$  and  $2 < x < 4$ , and is negative for  $0 < x < 2$  and  $4 < x < +\infty$ . Finally, suppose you know the  $x$ -axis is a horizontal asymptote and that  $(2, 0)$  is the only  $x$ -intercept. Draw a rough sketch of the function showing all these properties.

- (7) 2. For the following function

$$y = \left(1 + \frac{1}{x}\right)^2$$

graph the function, identifying all intercepts, asymptotes, local extrema, and inflection points. Specify intervals where the graph is increasing, decreasing, concave up, and concave down. Show all your work.

- (6) 3. A rectangle is to be inscribed in a half circle of radius = 1 as shown in the figure; find the dimensions that give the rectangle with the largest possible area.



- (3×5) 4. Evaluate the following:

(a)  $\int_0^{\frac{\pi}{3}} \left(x^2 - \frac{1}{\cos^2 x}\right) dx$

(b)  $\int_1^2 \left(t - \frac{1}{t}\right)^2 dt$

(c)  $\int (e^x + \sqrt[5]{x^3} - e^2) dx$

- (5) 5. Given  $f''(x) = 2 + \cos x$  satisfying  $f'(\pi) = 2\pi$ ,  $f(\pi) = 0$ , find the function  $f$ .
- (3) 6. For the function  $f(x) = \int_0^{x^2} \frac{1}{1+t^4} dt$ , what is the derivative  $f'(x)$ ? What is the minimum value of  $f(x)$ ?
- (4) 7. The graph of  $y = \frac{3}{4}x^4 - 6x^2 + 12$  has exactly two  $x$ -intercepts (at  $\pm 2$ ), and so there is a closed region between them, the curve, and the  $x$ -axis. Find the area of that region. (You may—or may not!—find that sketching a very rough graph will help you solve this problem.)
- (4) 8. Express the integral  $\int_0^2 x 2^{2x} dx$  as the limit of a Riemann sum. (You do not need to evaluate the limit!)

(Total: 50)