



Cal II (S) (Maths 201–NYB)

Answers

Answers

- (a) 1 (b) ∞
- (a) D (*n*TT) (b) C (to 1)
- (a) (LCT) with $\sum \frac{1}{n^2}$ (C *p*S): $\lim \frac{\sin(1/n^2)}{1/n^2} = 1$ so \sum converges
(b) (RT): $\lim \frac{a_{n+1}}{a_n} = \infty > 1$: \sum diverges
(c) (CT) with $\sum \frac{1}{n^{3/2}}$ (C *p*S), or
(*f*T): f cont, pos, decr; $\int_2^\infty f dx = \frac{1}{2} \ln 2 + \frac{1}{2}$ converges: \sum converges
(d) ($\sqrt[n]{}$ T): $\lim \sqrt[n]{a_n} = \lim \frac{25}{n} = 0 < 1$: \sum converges
- (a) (*n*TT): $\lim \frac{2^n}{n^2} = \infty \neq 0$: \sum diverges (could use (RT) instead)
(b) (LCT) with $\sum \frac{1}{n^{5/4}}$ (C *p*S): $\lim \frac{a_n}{b_n} = \frac{1}{\sqrt[4]{2}} \neq 0, \neq \infty$ so AC.
- $-6 \leq x < 2$ ($R = 4$) (Use (AST) and (LCT) at endpoints)
- (a) $a_n \rightarrow 0$ (*n*TT)
(b) It converges: (LCT) with $\sum a_n$.
- $\sum_{n=1}^{\infty} (-1)^{n+1} x^n$; converges for $-1 < x < 1$.