

Assignment # 2: Turán- and Ramsey-type problems.

Due in class on Thursday, March 12th.

1. Let  $G$  be a graph on  $n$  vertices for some  $n \geq 3$  with  $|G| \geq \lfloor \frac{n^2}{4} \rfloor + 1$ .
  - a) Show that  $G$  contains at least  $\lfloor \frac{n}{2} \rfloor$  triangles.
  - b) Show that the bound in a) is tight: For every  $n \geq 3$  there exists a graph  $G$  on  $n$  vertices with  $|G| = \lfloor \frac{n^2}{4} \rfloor + 1$  containing exactly  $\lfloor \frac{n}{2} \rfloor$  triangles.

2. Show that for every positive integer  $t$  there exists  $\delta > 0$  such that the following holds. If  $G$  is a graph not containing  $K_t$  on  $n$  vertices and every vertex of  $G$  belongs to at least  $\left(\frac{t-2}{t-1} - \delta\right)n$  edges then  $G$  is  $(t-1)$ -colorable.

3. **Bollobás. 8.7.** Let  $K_4^{(3)}$  denote the complete 3-graph on 4 vertices, i.e. the 3-graph isomorphic to  $[4]^{(3)}$ . Following de Caen (1983), we give an upper bound on  $\pi(K_4^{(3)})$ . Let  $\mathcal{F} \subseteq [n]^{(3)}$  be a hypergraph containing no  $K_4^{(3)}$  with  $|\mathcal{F}| = m$ . For  $x, y \in [n]$ ,  $x \neq y$  let

$$A(x, y) := \{z \in [n] \mid \{x, y, z\} \in \mathcal{F}\},$$

and let  $a_{xy} := |A(x, y)|$ . Note that if  $\{x, y, z\} \in \mathcal{F}$  then  $A(x, y) \cap A(y, z) \cap A(z, x) = \emptyset$  and so

$$a_{xy} + a_{yz} + a_{zx} \leq 2n - 3.$$

Summing over all edges of  $\mathcal{F}$  deduce that

$$\sum_{\{x,y\} \in [n]^{(2)}} a_{xy}^2 \leq (2n - 3)m.$$

Using convexity of  $x^2$  show that the left hand side is at least  $(3m)^2 / \binom{n}{2}$  and deduce that  $m \leq \frac{2n-3}{9} \binom{n}{2}$  and  $\pi(K_4^{(3)}) \leq 2/3$ .

4. Let  $G$  be a graph with  $V(G) = [17]$  and  $x, y \in V(G)$  adjacent if and only if

$$(x - y) \bmod 17 \in \{\pm 1, \pm 2, \pm 4, \pm 8\}.$$

a) Show that neither  $G$  nor the complement of  $G$  contains a  $K_4$  subgraph.

b) Deduce that  $R(4, 4) = 18$ .

5. **Hypergraph Ramsey theorem.** Show that for all positive integers  $r, k_1$  and  $k_2$  there exists a positive integer  $n = R^{(r)}(k_1, k_2)$  so that the following holds. If elements of  $[n]^{(r)}$  are colored in colors red and blue then there is a set  $Z \subseteq [n]$  such that either  $|Z| = k_1$  and all elements of  $Z^{(r)}$  are red, or  $|Z| = k_2$  and all elements of  $Z^{(r)}$  are blue.

(Hint: Use induction on  $r$  and, for given  $r$ , induction on  $k_1 + k_2$ . Consider all hyperedges containing a given vertex and attempt to imitate the proof of Ramsey's theorem.)

6. Show that for each  $\varepsilon > 0$  there exists  $N$  with the following property. For each real  $\alpha > 0$  there exist integers  $q$  and  $p$  such that  $1 \leq q \leq N$  and

$$|q^2 \alpha - p| \leq \varepsilon.$$

(Hint: Use van der Waerden's theorem.)