

MATH 550: Combinatorics. Winter 2016.

Assignment # 3: Discrete Geometry.

Due by e-mail (sent to snorine@gmail.com) by Monday, April 18th.

1. For $X \subseteq \mathbb{R}^d$ define $S(X)$ as a set of all points which lie on segments with ends in X . Let $S_2(X) := S(S(X))$ and, more generally, $S_{k+1}(X) = S(S_k(X))$. Show that $S_{\lceil \log_2(d+1) \rceil}(X)$ is always convex.

2.

- (a) Show that if $x, y, z \in \mathbb{R}^2$ are three points at pairwise distance at most 1 then there exists a disk in \mathbb{R}^2 of radius $1/\sqrt{3}$ containing x, y and z .
- (b) Show that if $X \subseteq \mathbb{R}^2$ is a finite set of diameter at most 1 then X is contained in some disk of radius $1/\sqrt{3}$.
- (c) Find the minimum c such that every finite set of diameter at most 1 in \mathbb{R}^3 is contained in some ball of radius c .

3. Matoušek. 1.3.4. A *strip of width w* is a part of the plane bounded by two parallel lines at distance w . The *width* of a set $X \subseteq \mathbb{R}^2$ is the smallest width of a strip containing X .

- (a) Show that every compact convex set of width 1 contains a segment of length 1 in every direction.
- (b) Let C_1, C_2, \dots, C_n be closed convex sets in the plane, $n \geq 3$, such that the intersection of every 3 of them has width at least 1. Show that $\bigcap_{i=1}^n C_i$ has width at least 1.

4.

- (a) Prove that if a collection of n convex sets in \mathbb{R}^2 has the property that out of every 4 sets some three have a point in common then there is a point that belongs to at least $n/12$ sets in the collection.
- (b) Prove that for all positive integers p, d so that $p \geq d + 1$ there exists a constant $c = c(d, p) > 0$ so that if a family of $n \geq p$ convex sets in \mathbb{R}^d has the property that among any p sets some $d + 1$ have a point in common then some point belongs to at least cn sets in the family.
- (c) Prove that for every positive integer d there is a constant $c = c(d)$ such that if a family \mathcal{F} of n convex sets in \mathbb{R}^d has the property that out of any $d + 2$ sets in \mathcal{F} some $d + 1$ have a point in common, then \mathcal{F} can be partitioned into at most $c \log n$ intersecting sub-families.

5. Matoušek. 4.1.5 (a). Use the Szemerédi-Trotter theorem to show that n points in the plane determine at most $O(n^{7/3})$ triangles of unit area.

6. Tao-Vu. 8.2.6. (Beck's theorem.) Let $P \subseteq \mathbb{R}^2$ be finite. Show that there either exists a line incident with $\Omega(|P|)$ points in P or there exist $\Omega(|P|^2)$ lines incident with at least 2 points in P .