

Math 340: Discrete Structures II

Practice Exam 1


Instructions. The exam is 3 hours long and contains 6 questions. Write your answers clearly in the notebook provided. You may quote any result/theorem seen in the lectures or in the assignments without proving it (unless, of course, that is what you are asked to prove).

1. *Matchings.*

- (a) What is a perfect matching?
- (b) True or false: a non-bipartite graph cannot contain a perfect matching.
- (c) Take a bipartite graph $G = (V, E)$ where the two parts of V in the bipartition are X and Y , where $|X| = |Y|$.
 - i. State Hall's Theorem.
 - ii. Prove Hall's Theorem.

2. *Planar Graphs.*

- (a)
 - i. State Kuratowski's theorem.
 - ii. State Euler's formula.
- (b) Prove that any graph with at most eight edges is planar.
- (c) Explain whether or not the following graph is planar.



3. *Probability.*

- (a) Two dice are rolled. At least one of the dice is a 6. What is the probability that the sum of the dice is 8?
- (b) We repeatedly roll a fair die until any number appears twice in a row. What is the expected number of rolls until we stop?
- (c) A coin is tossed until the first head appears. You win 2^n dollars if the first head appears on the n th toss.
 - i. What are the expected winnings if you play the game?
 - ii. Are you willing to pay this amount to play the game?

4. *Probability.*

- (a) State the Chernoff bound.
- (b) Suppose we have n boxes and we start randomly and independently throwing balls into the boxes.
 - i. If we throw exactly n balls, give an upper bound on the probability that Box #1 contains more than $2 \ln n$ balls.
 - ii. What is the expected number of balls we need to throw before *every* bin contains at least one ball?

5. *Combinatorics.*

- (a) State the binomial theorem.
- (b) Consider the equality

$$2^n = \sum_{k=0}^n \binom{n}{k}$$

- i. Prove this equality using the binomial theorem.
 - ii. Prove it combinatorially.
- (c) Prove combinatorially that, for $n \geq 1$,

$$\sum_{0 \leq k \leq n: k \text{ odd}} \binom{n}{k} = \sum_{0 \leq k \leq n: k \text{ even}} \binom{n}{k}$$

6. *Combinatorics.*

- (a) How many different ways are there to make up 22 cents using coins of denomination 1, 5 and 10 cents?
- (b) Let $f(n)$ be the number of ways to make up n cents using coins of denomination 1, 5 and 10 cents if we can use at most four 1 cent coins. Give the ordinary generating function $F(x)$.
- (c) Using b) or otherwise, obtain a simple expression for $f(n)$.