

THE CHU CONSTRUCTION

MICHAEL BARR

Transmitted by

Dedicated to the memory of Robert W. Thomason, 1952–1995

ABSTRACT. We take another look at the Chu construction and show how to simplify it by looking at it as a module category in a trivial Chu category. This simplifies the construction substantially, especially in the case of a non-symmetric biclosed monoidal category. We also show that if the original category is accessible, then for any of a large class of “polynomial-like” functors, the category of coalgebras has cofree objects.

1. Introduction

In a recent paper, I showed how the Chu construction, given originally in [Chu, 1979] for symmetric monoidal closed categories, could be adapted to monoidal biclosed (but not necessarily symmetric) categories. The construction, although well motivated by the necessity of providing a doubly infinite family of duals, was rather complicated with many computations involving indices. Recently I have discovered that the $*$ -autonomous structure of Chu categories can be put into the familiar context of bimodules over a not necessarily commutative “algebra” object (really, it is a monoid object, but the dual has always been called a coalgebra). It is a familiar fact that over an ordinary ring or algebra, the category of bimodules is a monoidal biclosed category. It was, in fact, the motivating example that led Eilenberg and Kelly to include the non-symmetric case of closed categories in their original paper in 1966.

We begin with a rational reconstruction of how the Chu construction might have been discovered. This is very far from how it actually happened, of course, which we describe briefly at the end. We then extend this construction to the non-symmetric case, which is rather easy. In fact, this whole idea of relating the Chu construction to bimodules was discovered in the non-symmetric case, where the analogy with bimodules became compelling. This turns out to be no analogy, just an instance. We then show how one instance of this non-symmetric Chu construction described in [Barr, 1995].

One of the interesting properties of the symmetric Chu construction is that if the original category is locally presentable (that is, accessible and complete), then the Chu category has cofree cocommutative coalgebras, which means there is a $!$ construction. This is interesting especially in light of the fact that the Chu category of a complete category

This research has been supported by grants from the NSERC of Canada and the FCAR du Québec

AMS subject classification (1990): 18D15.

Key words and phrases: Chu category, bimodules, cofree coalgebras.

© Michael Barr . Permission to copy for private use granted.

cannot be accessible unless the original category is a poset. In the non-symmetric case, the notion of cocommutativity is not even definable, so that a ! cannot be constructed that way. Nonetheless, it is interesting to observe that an interesting class of functors R , the category of R coalgebras has a cofree cotriple. This class consists of the smallest class of functors closed under product, sum and tensor product. Such functors are polymorphic over the class of locally presentable monoidal categories, a crucial point in the proof, which is by induction on the structural complexity.

2. A rational reconstruction

2.1. The symmetric case. For simplicity, we deal first with the symmetric case, although this works perfectly well in the non-symmetric case and was, in fact, first discovered in that context.

During the year I spent at the ETH in 1975-76, I was interested in self-dual categories. Now it is very easy to find self-dual categories; $\mathcal{V} \times \mathcal{V}^{\text{op}}$ is self dual regardless of the nature of \mathcal{V} . So more specifically, I was interested in finding closed self-dual categories in which the dual was given as the internal hom into some “dualizing object”. Suppose \mathcal{V} is a monoidal category. What about $\mathcal{V} \times \mathcal{V}^{\text{op}}$? If you make no particular requirements on the tensor (except that it be a bifunctor), you can simply define $(U, U') \otimes (V, V') = (U \otimes V, U' \otimes V')$. But if \mathcal{V} is closed, $\mathcal{V} \times \mathcal{V}^{\text{op}}$ is unlikely to be closed since the tensor will commute with limits in the second coordinate, instead of colimits.

So let us suppose that the original category \mathcal{V} is symmetric closed monoidal and try to see what the corresponding autonomous structure on $\mathcal{V} \times \mathcal{V}^{\text{op}}$ ought to be. Let us suppose that we have $(U, U') \otimes (V, V') = (U \otimes V, X)$ and we are trying to determine X . Let us also suppose that the first coordinate of $(U, U') \multimap (V, V')$ is the internalization of $\text{Hom}((U, U'), (V, V')) = \text{Hom}(U, V) \times \text{Hom}(V', U')$. This suggests that $(U, U') \multimap (V, V') = ((U \multimap V) \times (V' \multimap U'), Y)$ where Y is to be determined. Finally, we observe that if the duality on the category determined by the internal hom into a dualizing object \perp , then we must have

$$\begin{aligned} ((U, U') \otimes (V, V'))^\perp &= (U, U') \otimes (V, V') \multimap \perp \\ &= (U, U') \multimap ((V, V') \multimap \perp) \\ &= (U, U') \multimap (V', V) = ((U \multimap V') \times (V \multimap U'), Y) \end{aligned}$$

from which we conclude that

$$(U, U') \otimes (V, V') = (U \otimes V, (U \multimap V') \otimes (V \multimap U'))$$

Similarly,

$$(U, U') \multimap (V, V') = ((U \multimap V) \times (V' \multimap U'), U \otimes V')$$

Does this work? It certainly does. A map

$$(U, U') \otimes (V \otimes V') \longrightarrow (W, W')$$

is equivalent to a pair of arrows $U \otimes V \rightarrow W$ and $W' \rightarrow (U \multimap V') \times (V \multimap U')$, which is equivalent to three arrows $U \otimes V \rightarrow W$, $W' \rightarrow U \multimap V'$ and $W' \rightarrow V \multimap U'$. Similarly, a map $(U, U') \rightarrow (V, V') \multimap (W, W')$ is a map $(U, U') \rightarrow ((V \multimap W) \times (W' \multimap V'), V \otimes W')$, which corresponds to three maps $U \rightarrow V \multimap W$, $U \rightarrow W' \multimap V'$ and $V \otimes W' \rightarrow U'$. These two sets of arrows are clearly in one-one correspondence.

2.2. The non-symmetric case. With some care, this also extends to the case of a biclosed monoidal category. We suppose that \mathcal{V} is such a category with two internal homs \multimap and \multimap that satisfy

$$\mathrm{Hom}(U \otimes V, W) \cong \mathrm{Hom}(V, U \multimap W) \cong \mathrm{Hom}(V, W \multimap U)$$

We note that there is some arbitrariness in which one to denote \multimap and which one is \multimap . The final choice was suggested by the example of a group considered as a biclosed monoidal category in which the elements of the group are the objects, the only maps are identities and the group multiplication is the tensor. In that case, $u \multimap v = u^{-1}v$ and $v \multimap u = vu^{-1}$, while if the other choice were made, we would have to interchange the elements. Incidentally, there is more to this analogy. For example, the internalizations of those hom isomorphisms are

$$U \otimes V \multimap W \cong V \multimap (U \multimap W) \quad W \multimap (U \otimes V) \cong (W \multimap V) \multimap U$$

However, there is one further isomorphism that turns out to be important.

2.3. PROPOSITION.

$$(U \multimap W) \multimap V \cong U \multimap (W \multimap V)$$

PROOF. We have, for any object T ,

$$\begin{aligned} \mathrm{Hom}(T, (U \multimap W) \multimap V) &\cong \mathrm{Hom}(T \otimes V, U \multimap W) \cong \mathrm{Hom}(U \otimes T \otimes V, W) \\ &\cong \mathrm{Hom}(U \otimes T, W \multimap V) \cong \mathrm{Hom}(T, U \multimap (W \multimap V)) \end{aligned}$$

■

This proof clearly depends on the associativity of the tensor and the proposition gives a kind of associativity between the left and right internal homs. We will often write $U \multimap W \multimap V$ for either one, just as with the tensor.

We will now suppose that \mathcal{V} is a biclosed monoidal category in this sense and also that $\sigma: \mathcal{V} \rightarrow \mathcal{V}$ is a given isomorphism that preserves all the structure. The role of σ will be explained later; for the time being, you can take it to be the identity.

We now define structures on $\mathcal{V} \times \mathcal{V}^{\mathrm{op}}$:

$$\begin{aligned} (U, U') \otimes (V, V') &= (U \otimes V, (V \multimap U') \times (V' \multimap \sigma^{-1}U)) \\ (U, U') \multimap (V, V') &= ((U \multimap V) \times (U' \multimap V'), V' \otimes \sigma^{-1}U) \\ (V, V') \multimap (U, U') &= ((V \multimap U) \times (\sigma V' \multimap \sigma U'), U \otimes V') \\ (U, U')^\perp &= (U', \sigma^{-1}U) \\ {}^\perp(U, U') &= (\sigma U', U) \end{aligned} \tag{*}$$

2.4. THEOREM. *The structures defined by (*) give a *-autonomous category.*

PROOF. First we show that it is a monoidal biclosed category, which means that for all objects (U, U') , (V, V') and (W, W') , we have

$$\begin{aligned} \text{Hom}((U, U') \otimes (V, V'), (W, W')) &\cong \text{Hom}((V, V'), (U, U') \multimap (W, W')) \\ &\cong \text{Hom}((U, U'), (W, W') \multimap (V, V')) \end{aligned}$$

For the first isomorphism, we calculate that an arrow

$$(U, U') \otimes (V, V') = (U \otimes V, (V \multimap U') \times (V' \multimap \sigma^{-1}U)) \rightarrow (W, W')$$

is given by three arrows

$$U \otimes V \rightarrow W, \quad W' \rightarrow (V \multimap U'), \quad W' \rightarrow (V' \multimap \sigma^{-1}U)$$

which transpose to

$$U \otimes V \rightarrow W, \quad V \otimes W' \rightarrow U', \quad W' \otimes \sigma^{-1}U \rightarrow V'$$

An arrow

$$(V, V') \rightarrow (U, U') \multimap (W, W') = ((U \multimap W) \times (U' \multimap W'), W' \otimes \sigma^{-1}U)$$

is given by three arrows

$$V \rightarrow (U \multimap W), \quad V \rightarrow (U' \multimap W'), \quad W' \otimes \sigma^{-1}U \rightarrow V'$$

which transpose to

$$V \otimes U \rightarrow W), \quad V \otimes W' \rightarrow U', \quad W' \otimes \sigma^{-1}U \rightarrow V'$$

which are the same data. For the second isomorphism, we find that a map

$$(U, U') \rightarrow (W, W') \multimap (V, V') = ((W \multimap V) \times (\sigma W' \multimap \sigma V'), V \otimes W')$$

is given by three arrows

$$U \rightarrow W \multimap V, \quad U \rightarrow \sigma W' \multimap \sigma V', \quad V \otimes W' \rightarrow U'$$

which transpose to

$$U \otimes V \rightarrow W, \quad \sigma W' \otimes U \rightarrow \sigma V', \quad V \otimes W' \rightarrow U'$$

Since σ is an isomorphism that preserves all structure, an arrow $\sigma W' \otimes U \rightarrow \sigma V'$ is the same as an arrow $W' \otimes \sigma^{-1}U \rightarrow V'$.

This demonstrates the closed monoidal structure. For the rest of the $*$ -autonomous structure, we must show that for any object \mathbf{U} of the Chu category, $\mathbf{U}^\perp = \mathbf{U} \multimap \top^\perp$ and that the second dual map $\mathbf{U} \rightarrow {}^\perp(\mathbf{U}^\perp)$ is an isomorphism. We see that $\top^\perp = (1, \top)$, which we will denote \perp . Then

$$(U, U') \multimap (1, \top) = (U \multimap 1 \times U' \multimap \top, \top \otimes \sigma^{-1}U) \cong (U', \sigma^{-1}U)$$

as required. For the second dual, we must show that the map $\mathbf{U} \rightarrow \perp \multimap (\mathbf{U} \multimap \perp)$, gotten by twice transposing the identity $\mathbf{U} \multimap \perp \rightarrow \mathbf{U} \multimap \perp$, is an isomorphism.

We begin by working out what map $\mathbf{U} \rightarrow \mathbf{V} \multimap \mathbf{W}$ corresponds to the second transpose of a given map $\mathbf{W} \rightarrow \mathbf{U} \multimap \mathbf{V}$. Assuming $\mathbf{U} = (U, U')$ and so on, the latter comes, as we have seen, from three arrows:

$$W \xrightarrow{f} U \multimap V, \quad W \xrightarrow{g} U' \multimap V', \quad V' \otimes \sigma^{-1}U \xrightarrow{h} W'$$

This corresponds to the map $\mathbf{U} \rightarrow \mathbf{V} \multimap \mathbf{W}$:

$$U \rightarrow V \multimap W, \quad U \rightarrow (\sigma V' \multimap \sigma W'), \quad W \otimes V' \rightarrow U'$$

described as follows. The first is the second transpose of f , the second is σ applied to the second transpose of h and the third is the transpose of g . Now we consider the special case that $\mathbf{V} = \perp$, $\mathbf{W} = \mathbf{U} \multimap \perp$ and we begin with the identity map. Then we begin with

$$\begin{aligned} p_1: (U \multimap 1) \times (U' \multimap \top) &\rightarrow U \multimap 1 \\ p_2: (U \multimap 1) \times (U' \multimap \top) &\rightarrow U' \multimap \top \\ \text{id}: \top \otimes \sigma^{-1}U &\rightarrow \top \otimes \sigma^{-1}U \end{aligned}$$

The corresponding maps $\mathbf{U} \rightarrow \perp \multimap (\mathbf{U} \multimap \perp)$ are

$$\begin{aligned} U &\rightarrow 1 \multimap [(U \multimap 1) \times (U' \multimap \top)] = 1 \\ U &\rightarrow \top \multimap (\sigma \top \otimes \sigma \sigma^{-1}U) \\ (U' \multimap \top) \otimes \top &\rightarrow U' \end{aligned}$$

The first map is the only map $U \rightarrow 1$, the second is gotten by applying σ to the second transpose of $\text{id}: \top \otimes \sigma^{-1}U \rightarrow \top \otimes \sigma^{-1}U$. The coherence identities involving transposition and \top make this an isomorphism. The result is that the combined map

$$U \rightarrow (1 \multimap [(U \multimap 1) \times (U' \multimap \top)]) \times (\top \multimap (\sigma \top \otimes \sigma \sigma^{-1}U))$$

is an isomorphism. The coherence identities also force the second transpose of $\text{id}: U' \multimap \top \rightarrow U' \multimap \top$ to be an isomorphism. It follows from Definition B of Section 2 of [Barr, 1995] that $\mathcal{V} \times \mathcal{V}$ is, with the given structure, a $*$ -autonomous category.

2.5. Algebras and modules. The notion of algebra (associative, unitary) makes sense in any tensored category. Let us work out what is required for (U, U') to be an algebra in $\mathcal{V} \times \mathcal{V}^{\text{op}}$. The tensor unit is $(\top, 1)$ so a unit is a map $(\top, 1) \rightarrow (U, U')$, which is nothing but a map $\top \rightarrow U$ in \mathcal{V} . A multiplication is a map

$$(U, U') \otimes (U, U') = (U \otimes U, U \multimap U' \times U' \multimap \sigma^{-1}U) \rightarrow (U, U')$$

which corresponds to maps $U \otimes U \rightarrow U$, $U' \rightarrow U \multimap U'$ and $U' \rightarrow U' \multimap \sigma^{-1}U$ in \mathcal{V} . The first of these, together with the unit, makes U into an algebra in \mathcal{V} . The second arrow defines a left U -module structure on U and the third defines a right $\sigma^{-1}U$ -module structure on U' .

Since $\top = \sigma^{-1}(\top)$ is obviously an algebra object in \mathcal{V} —the initial algebra object, in fact—and every object of \mathcal{V} is a bimodule over \top , it follows immediately that every object of the form (\top, \perp) is an algebra, regardless of the nature of \perp . These are the ones that will be of interest to us later.

3. The biclosed monoidal structure

In this section, we describe how the category of bimodules for an algebra object in a monoidal biclosed category is monoidal biclosed and $*$ -autonomous when the original category is. Let \mathcal{V} be a monoidal biclosed category and suppose that K is an algebra object, associative and unitary. The development parallels the familiar one for rings. We will talk of right, left and two-sided modules, always meaning with respect to the algebra K . These facts are essentially known, see [Street, 1983] or [Kosłowski, forthcoming]. Each of these papers gives the proofs in a more general context. In Street's paper, the algebra has many objects and in Kosłowski's the category of left K , right L -bimodules is the hom category in a bicategory whose objects are algebras. We give the constructions and omit the proofs.

3.1. The basic definitions. Suppose K is an algebra object, M is a right K -module and N a left K -module. Let the two actions be denoted $\rho(M, K): M \otimes K \rightarrow M$ and $\lambda(K, N): K \otimes N \rightarrow N$. Define $M \otimes_K N$ so that

$$M \otimes K \otimes N \begin{array}{c} \xrightarrow{\rho(M, K) \otimes N} \\ \xrightarrow{M \otimes \lambda(K, N)} \end{array} M \otimes N \rightarrow M \otimes_K N$$

is a coequalizer.

For the next construction, suppose that M and N are left K -modules. Then define $M \xrightarrow{K} \circ N$ so that

$$M \xrightarrow{K} \circ N \rightarrow M \multimap N \rightrightarrows (K \otimes M) \multimap N$$

is an equalizer. One of the two maps $M \multimap N \rightarrow (K \otimes M) \multimap N$ is just induced by $\lambda(K, M): K \otimes M \rightarrow M$ and the other is the transpose of the map

$$K \otimes M \otimes (M \multimap N) \xrightarrow{\text{eval}} K \otimes N \xrightarrow{\lambda(K, N)} N$$

Thus this is the internalization of the set of arrows that preserve the K structure.

The third construction begins with right K -modules M and N and defines $N \circ_{\overline{K}} M$ so that

$$N \circ_{\overline{K}} M \rightarrow N \circ - M \rightrightarrows N \circ -(M \otimes K)$$

is an equalizer. The two arrows are defined similarly to the preceding case.

There appears to be no way of defining $M \xrightarrow{\overline{K}} N$ for two right modules or $N \circ_{\overline{K}} M$ for two left modules. This is fortunate, since it means that when M and N are bimodules, the notations $M \xrightarrow{\overline{K}} N$ and $N \circ_{\overline{K}} M$ are unambiguous. It is analogous to the unambiguous nature of $M \otimes_K N$, using the right structure on M and the left one on N . Another point to note is that, given that the given tensor is non-symmetric, there appears to be no way of defining an opposite algebra K^{op} in such a way that right K -modules are left K^{op} -modules. Similarly, there is no notion, given two algebras K and L , of K, L -bimodule. What there is is a notion of left K , right L -bimodule, to which we now turn.

3.2. Bimodules. In this section, we are primarily interested in left and right K -bimodules. However we investigate the notion of left K , right L bimodule, not so much for the added generality, but also it is less confusing and certainly no harder.

By a left K , right L bimodule, we mean an object M that is both a left K -module and a right L -module such that the square

$$\begin{array}{ccc} K \otimes M \otimes L & \xrightarrow{K \otimes \rho(M, L)} & M \otimes L \\ \lambda(K, M) \otimes L \downarrow & & \downarrow \rho(M, L) \\ K \otimes M & \xrightarrow{\lambda(K, M)} & M \end{array}$$

commutes. For convenience, we will use the notation ${}_K M_L$ to indicate that M is a left K , right L -bimodule. If ${}_K M_L$ and ${}_K N_L$, I will write $\text{Hom}_K(M, N)_L$ to denote the set of left K , right L morphisms. If K or L is the tensor unit, it can be omitted from the notation.

3.3. PROPOSITION. *Suppose that J, K and L are algebras and that ${}_J M_K$ and ${}_K N_L$. Then $M \otimes_K N$ gets the structure of a left J , right L -bimodule.*

As mentioned above, we omit the proof, but give the construction. Since \otimes is a left adjoint, it preserves coequalizers so that the top line of the diagram

$$\begin{array}{ccccc} J \otimes M \otimes K \otimes N \otimes I & \rightrightarrows & J \otimes M \otimes N \otimes I & \longrightarrow & J \otimes M \otimes_K N \otimes I \\ \lambda(J, M) \otimes K \otimes \rho(N, I) \downarrow & & \lambda(J, M) \otimes \rho(N, I) \downarrow & & \\ M \otimes K \otimes N & \rightrightarrows & M \otimes N & \longrightarrow & M \otimes_K N \end{array}$$

is a coequalizer. The diagram serially commutes and thus induces an arrow $J \otimes M \otimes_K N \otimes I \rightarrow M \otimes_K N$, which gives the structure.

3.4. PROPOSITION. *Suppose J , K and L are algebras. Suppose ${}_K M_J$ and ${}_K N_L$. Then $M \xrightarrow[\underline{K}]{} \circ N$ has the structure of a left J , right L bimodule.*

Here is the construction. The composite

$$M \otimes J \otimes (M \multimap N) \otimes L \rightarrow M \otimes (M \multimap N) \otimes L \rightarrow N \otimes L \rightarrow N$$

has as exponential transpose an arrow $J \otimes (M \multimap N) \otimes L \rightarrow M \multimap N$. Similarly, the composite

$$K \otimes M \otimes J \otimes ((K \otimes M) \multimap N) \otimes L \rightarrow K \otimes M \otimes ((K \otimes M) \multimap N) \otimes L \rightarrow N \otimes L \rightarrow N$$

has as exponential transpose an arrow $J \otimes ((K \otimes M) \multimap N) \otimes L \rightarrow (K \otimes M) \multimap N$. These are the vertical arrows in the diagram

$$\begin{array}{ccccc} J \otimes (M \xrightarrow[\underline{K}]{} \circ N) \otimes L & \longrightarrow & J \otimes (M \multimap N) \otimes L & \xRightarrow{\quad} & J \otimes ((K \otimes M) \multimap N) \otimes L \\ & & \downarrow & & \downarrow \\ (M \xrightarrow[\underline{K}]{} \circ N) & \longrightarrow & (M \multimap N) & \xRightarrow{\quad} & ((K \otimes M) \multimap N) \end{array}$$

The diagram is serially commutative and the bottom row is an equalizer, so there is induced an arrow $J \otimes (M \xrightarrow[\underline{K}]{} \circ N) \otimes L \rightarrow M \xrightarrow[\underline{K}]{} \circ N$ which is the induced structure giving the left J , right L structure on the internal homset.

By interchanging left and right, we also have,

3.5. PROPOSITION. *Suppose J , K and L are algebras. Suppose ${}_L M_K$ and ${}_J N_K$. Then $N \circ \xrightarrow[\underline{K}]{} M$ has the structure of a left J , right L bimodule.*

These structures are related by the following theorem. Although it is stated in terms of objects, these are actually instances of natural equivalences, natural in M , N and P .

3.6. THEOREM. *Suppose J , K and L are algebra objects and ${}_J M_K$, ${}_K N_L$ and ${}_J P_L$. Then*

$$\mathrm{Hom}_J(M \otimes_K N, P)_L \cong \mathrm{Hom}_K(N, M \xrightarrow[\underline{J}]{} \circ P)_L \cong \mathrm{Hom}_J(M, P \circ \xrightarrow[\underline{L}]{} N)_K$$

There are also isomorphisms of the internal homs that look similar to the above. They are not really internalizations of the above because they concern homs of only left or right structure, not the bimodule structure.

3.7. THEOREM.

1. *In the situation ${}_J M_K$, ${}_K N_I$, ${}_J P_L$, we have*

$$M \otimes_K N \xrightarrow[\underline{J}]{} \circ P \cong N \xrightarrow[\underline{K}]{} \circ (M \xrightarrow[\underline{J}]{} \circ P)$$

as left I , right L -modules.

2. In the situation ${}_J M_K, {}_K N_I, {}_L P_I$, we have

$$P \circ_{\overline{I}} M \otimes_K N \cong (P \circ_{\overline{I}} N) \circ_{\overline{K}} M$$

as left L , right J -bimodules.

PROOF. We prove the first, the second is similar. Let Q be any left I , right L -bimodule. Then

$$\begin{aligned} \mathrm{Hom}_I(Q, M \otimes_K N \xrightarrow{J} P)_L &\cong \mathrm{Hom}_J(M \otimes_K N \otimes_I Q, P)_L \\ &\cong \mathrm{Hom}_K(N \otimes_I Q, M \xrightarrow{J} P)_L \\ &\cong \mathrm{Hom}_I(Q, N \xrightarrow{K} (M \xrightarrow{J} P))_L \end{aligned}$$

Since these isomorphisms are evidently natural in Q , the conclusion follows from Yoneda. ■

Street proved these results under somewhat stronger hypotheses, but he allowed his “algebras” to have many objects. Our hypotheses of equalizers and coequalizers suffice for this case.

What Koslowski actually proves is that if you begin with a biclosed monoidal category there is a biclosed monoidal bicategory whose objects are the algebra objects in the original category and for which the hom category between the algebra objects K and L has as objects left K , right L bimodules and morphisms thereof. The monoidal structure and the biclosed structure is exactly as described above and Koslowski shows that this all works as advertised.

3.8. The symmetric case. Suppose that the tensor product in \mathcal{V} is symmetric. This means that there are natural isomorphisms $c = c(U, V): U \otimes V \rightarrow V \otimes U$ that satisfy the usual coherence isomorphisms, including that $c(V, U) \circ c(U, V) = \mathrm{id}$. In that case, we can consider the case that K is a commutative algebra object and then define a module to be symmetric when the left and right actions coincide in the sense that

$$\begin{array}{ccc} K \otimes V & \xrightarrow{c(K, V)} & V \otimes K \\ \lambda(K, V) \searrow & & \nearrow \rho(V, K) \\ & V & \end{array}$$

commutes. The subcategory of symmetric modules can be shown to be closed under tensor and right and left internal hom and the last two are also isomorphic.

4. The non-symmetric Chu construction

In this section, we will show how the non-symmetric Chu construction of [Barr, 1995] comes as special case of what we have done. Begin with a monoidal biclosed category

\mathcal{V} . Let \mathcal{W} be the category of \mathbf{Z} -graded \mathcal{V} objects. That is, an object of \mathcal{W} is a doubly infinite sequence $(\dots, V_{-1}, V_0, V_1, \dots)$ of objects of \mathcal{V} . An arrow is simply a doubly infinite sequence of arrows, subject to no conditions. In fact, this is just the cartesian product of countably many copies of \mathcal{V} , although we number it using \mathbf{Z} , not \mathbf{N} . With coordinate-wise operations, this is also a monoidal biclosed category. We let $\sigma: \mathcal{W} \rightarrow \mathcal{W}$ be the automorphism that renumbers everything one higher. That is, if \mathbf{V} is the object above, whose n th component \mathbf{V}_n is V_n , then $\sigma(\mathbf{V})_n = V_{n-1}$. An object of $\text{Chu}(\mathcal{W}, \perp)$ is a pair $(\mathbf{V}, \mathbf{V}')$ equipped with pairings $\mathbf{V} \otimes \mathbf{V}' \rightarrow \perp$ and $\sigma\mathbf{V}' \otimes \mathbf{V} \rightarrow \perp$. This translates to $V_n \otimes V'_n \rightarrow \perp$ and $V'_{n-1} \otimes V_n \rightarrow \perp$. We relate this to the theory of [Barr, 1995] by defining a graded object \mathbf{W} by $W_{2n} = V_n$ and $W_{2n+1} = V'_n$. Then the map $V_n \otimes V'_n \rightarrow \perp$ becomes a map $W_{2n} \otimes W_{2n+1} \rightarrow \perp$, while the map $V'_{n-1} \otimes V_n \rightarrow \perp$ becomes a map $W_{2n-1} \otimes W_{2n} \rightarrow \perp$. Thus we see that we get simply $W_m \otimes W_{m+1} \rightarrow \perp$ for all $m \in \mathbf{Z}$. But this was the definition used in the earlier paper. It is easy to verify that all the other definitions used in that paper correspond to what we get here. It is a fact that the formulas of this paper were derived by looking at the earlier construction and seeing how they looked in this framework. This was how I determined the placement of σ and σ^{-1} was determined in Equation (*) at the end of Section 2.2.

5. Cofree coalgebras

In this section, we will suppose that \mathcal{V} is a locally presentable, that is accessible and complete, category. See [Makkai and Paré, 1990] as well as [Gabriel and Ulmer, 1971]. In the symmetric case, this implies that there are cofree objects in the category of commutative, coassociative, counitary coalgebras, which gives a natural model for the ! construction of linear logic, [Barr, 1991]. In the non-symmetric case, cocommutativity cannot be defined and we cannot model ! in this way. Nonetheless, it could be useful to know that cofree coalgebras exist. We will actually show here that cofree coalgebras of the form $C \rightarrow RC$ exist for a large class of functors R .

5.1. THEOREM. *Suppose \mathcal{V} is a locally presentable category and \perp is an object of \mathcal{V} . Let \mathcal{R} denote the least class of functors that is closed under arbitrary products and sums and under finite (including empty) tensor products. Then for any $R \in \mathcal{R}$, the category of R -coalgebras in $\text{Chu}(\mathcal{V}, \perp)$ has cofree coalgebras.*

The proof will occupy the rest of this section. Note that the class \mathcal{R} makes sense in any bicomplete monoidal category. This is why we did not include all the constants, although it includes the initial and terminal objects as well as the tensor unit as constant functors and the identity functor as a unary sum (or product or tensor).

We begin with the fact that the functors in \mathcal{R} are accessible on \mathcal{V} . Since \mathcal{V} is an accessible category (which $\text{Chu}(\mathcal{V}, \perp)$ is not), this shows that for any $R \in \mathcal{R}$, there are cofree R -coalgebras in \mathcal{V} .

5.2. PROPOSITION. *The functors in \mathcal{R} are accessible on \mathcal{V} .*

PROOF. It is standard that the sum and product of accessible functors is accessible, so we need worry only about tensor products. But the tensor with a fixed object has a right adjoint, hence commutes with arbitrary colimits. We require the following, whose proof is left to the reader.

5.3. LEMMA. *Suppose α is an infinite cardinal and \mathcal{I} is an α -filtered category. Then the diagonal $\Delta: \mathcal{I} \rightarrow \mathcal{I} \times \mathcal{I}$ is cofinal. \blacksquare*

Now if $D: \mathcal{I} \rightarrow \mathcal{V}$ is an α -filtered diagram and $R_1, R_2: \mathcal{V} \rightarrow \mathcal{V}$ are α -accessible, then

$$\begin{aligned} \operatorname{colim}_{i \in \mathcal{I}} (R_1 \otimes R_2)Di &\cong \operatorname{colim}_{i \in \mathcal{I}} (R_1Di \otimes R_2Di) \cong \operatorname{colim}_{i \in \mathcal{I}} \operatorname{colim}_{j \in \mathcal{I}} (R_1Di \otimes R_2Dj) \\ &\cong \operatorname{colim}_{i \in \mathcal{I}} (R_1Di \otimes R_2(\operatorname{colim}_{j \in \mathcal{I}} Dj)) \cong R_1(\operatorname{colim}_{i \in \mathcal{I}} Di) \otimes R_2(\operatorname{colim}_{j \in \mathcal{I}} Dj) \end{aligned}$$

It follows from [Makkai, Paré, 1990] that for any $R \in \mathcal{R}$, the underlying functor from the category of R -coalgebras in \mathcal{V} to \mathcal{V} has a right adjoint. We will denote this functor by G_R . It is the functor part of a cotriple $\mathbf{G}_R = (G_R, \epsilon_R, \delta_R)$ on \mathcal{V} for which the category of coalgebras is just the category of R -coalgebras and homomorphisms.

5.4. PROPOSITION. *The functor R on $\operatorname{Chu}(\mathcal{V}, \perp)$ has the form*

$$R(V, V') = (RV, \tilde{R}(V, V'))$$

where $\tilde{R}: \mathcal{V}^{\text{op}} \times \mathcal{V} \rightarrow \mathcal{V}$ is a functor.

PROOF. Since $\prod(V_i, V'_i) = (\prod V_i, \sum V'_i)$ and $\sum(V_i, V'_i) = (\sum V_i, \prod V'_i)$ the assertion follows by structural induction for sums and products of such functions. As for tensor products, the conclusion follows from

$$(U, U') \otimes (V, V') = (U \otimes V, (V \multimap U') \times_{V \multimap \perp \multimap \sigma^{-1}U} (V' \multimap \sigma^{-1}U))$$

in which the first coordinate is just the tensor product of the first coordinates and the second is functorial in both, contravariant in the first and covariant in the second. Thus if $R = R_1 \otimes R_2$, then

$$\tilde{R}(V, V') = R_2(V) \multimap \tilde{R}_1(V, V') \times_{R_2(V) \multimap \perp \multimap \sigma^{-1}R_1(V)} \tilde{R}_2(V, V') \multimap \sigma^{-1}R_1(V)$$

5.5. PROPOSITION. *Let \mathcal{V} be a monoidal biclosed accessible category. For a fixed object U of \mathcal{V} , the functors $U \multimap -$ and $- \multimap U$ are accessible.*

PROOF. It is clearly sufficient to do either one, say $U \multimap -$. Let Γ be a set of generators for \mathcal{V} and let α be the sup of the presentation rank of all objects of the form W and $U \otimes W$, where $W \in \Gamma$. Now let $V = \operatorname{colim} V_i$, the colimit taken over an α filtered diagram. Then for any $W \in \Gamma$, we have

$$\begin{aligned} \operatorname{Hom}(W, U \multimap V) &\cong \operatorname{Hom}(U \otimes W, V) \cong \operatorname{colim} \operatorname{Hom}(U \otimes W, V_i) \\ &\cong \operatorname{colim} \operatorname{Hom}(W, U \multimap V_i) \cong \operatorname{Hom}(W, \operatorname{colim}(U \multimap V_i)) \end{aligned}$$

Since W ranges over a generating set, it follows that the induced map $\text{colim}(U \multimap V_i) \rightarrow U \multimap V$ is an isomorphism. \blacksquare

5.6. PROPOSITION. *For a fixed object V of \mathcal{V} and $R \in \mathcal{R}$, the corresponding functor \tilde{R} is accessible as a functor $\mathcal{V} \rightarrow \mathcal{V}$.*

PROOF. If $R = \prod R_i$ and \tilde{R}_i is the functor corresponding to R_i , then

$$R(V, V') = \left(\prod R_i(V), \sum \tilde{R}_i(V, V') \right)$$

so that $\tilde{R}(V, V') = \sum \tilde{R}_i(V, V')$ and the sum of accessible functors is accessible. A similar argument, since the product of accessible functors is accessible, allows us to draw the same consequence for sums. Finally, we have to deal with the tensor product. Suppose that $R = R_1 \otimes R_2$, $R_1 = (R_1, \tilde{R}_1)$ and $R_2 = (R_2, \tilde{R}_2)$ and we know that \tilde{R}_1 and \tilde{R}_2 are accessible in their second variable. Then

$$R(V, V') = (RV, (R_1V \multimap \tilde{R}_2(V, V')) \times_{R_1V \multimap \perp \multimap \sigma R_2V} (\tilde{R}_1(V, V') \multimap \sigma R_2(V)))$$

The objects $R_1(V)$ and $R_2(V)$ are fixed. Thus by the preceding proposition, it follows that \tilde{R} is accessible in the second variable. \blacksquare

We introduce some temporary notation in order to reduce the notational complication of the rest of the argument. If V is an object of \mathcal{V} , let $V^* = V \multimap \perp \times \perp \multimap \sigma^{-1}V$. This is not exactly a dual, but it is the case that there is a one-one correspondence between arrows $W \rightarrow V^*$ and $\sigma^{-1}V \rightarrow W^*$, although we will not use that fact. What we will use are the obvious facts that a Chu structure on (V, V') is given by an arrow $V' \rightarrow V^*$ and that $(f, f'): (V, V') \rightarrow (W, W')$ is a pair of arrows for which the square

$$\begin{array}{ccc} W' & \xrightarrow{f'} & V' \\ \downarrow & & \downarrow \\ W^* & \xrightarrow{f^*} & V^* \end{array}$$

commutes.

We now fix an $R \in \mathcal{R}$. Since R is accessible, there is a cofree R -coalgebra cotriple on \mathcal{V} that we will denote $\mathbf{G} = (G, \epsilon, \delta)$. If (V, V') is an object of $\text{Chu}(\mathcal{V}, \perp)$ there is a natural Chu structure on (GV, V') , namely

$$\begin{aligned} GV \otimes V' &\xrightarrow{\epsilon V \otimes 1} V \otimes V' \rightarrow \perp \\ \sigma V' \otimes GV &\xrightarrow{1 \otimes \epsilon V} \sigma V' \otimes V \rightarrow \perp \end{aligned}$$

We denote the R -coalgebra structure on GV by $\rho = \rho V: GV \rightarrow RGV$. If (C, C') is an R -coalgebra and $(f, f'): (C, C') \rightarrow (V, V')$ is an arrow in $\text{Chu}(\mathcal{V}, \perp)$, then C is an R -coalgebra in \mathcal{V} and $f: C \rightarrow V$ is an arrow. Thus f factors as $f = \epsilon V \circ \hat{f}$ where $\hat{f}: C \rightarrow GV$

is a coalgebra morphism. It is immediate that (f, f') factors as $(f, f') = (\epsilon V, 1) \circ (\widehat{f}, f')$. We claim that each of these maps is in $\text{Chu}(\mathcal{V}, \perp)$. These follow from the commutativity of

$$\begin{array}{ccc}
 V' & \xrightarrow{f'} & C' \\
 \downarrow & & \downarrow \\
 V^* & & C^* \\
 (\epsilon V)^* \downarrow & \searrow f^* & \downarrow \\
 (GV)^* & \xrightarrow{(\widehat{f})^*} & C^*
 \end{array}
 \qquad
 \begin{array}{ccc}
 V' & \xrightarrow{\text{id}} & V' \\
 \downarrow & & \downarrow \\
 V^* & \xrightarrow{(\epsilon V)^*} & (GV)^*
 \end{array}$$

In the left diagram, the upper trapezoid commutes because $(C, C') \rightarrow (V, V')$ is an arrow in $\text{Chu}(\mathcal{V}, \perp)$, while the lower triangle commutes from the application of the contravariant functor $(\)^*$ to the identity $\epsilon V \circ \widehat{f} = f$. The right hand square commutes by the definition of the Chu structure on (GV, V') .

Now define an object V^\sharp of \mathcal{V} so that

$$\begin{array}{ccc}
 V^\sharp & \xrightarrow{\widetilde{f}} & C' \\
 \downarrow & & \downarrow \\
 (GV)^* & \xrightarrow{(\widehat{f})^*} & C^*
 \end{array}$$

is a pullback.

5.7. PROPOSITION. *Let V^\sharp be constructed as above. Then*

1. (GV, V^\sharp) is an object of $\text{Chu}(\mathcal{V}, \perp)$;
2. $(\widehat{f}, f') : (C, C') \rightarrow (GV, V')$ factors through (GV, V^\sharp) ;
3. (GV, V^\sharp) is an R -coalgebra;
4. the first factor of this factorization is by an R -coalgebra morphism.

PROOF. We give it the structure of a Chu object by using the pullback diagram

$$\begin{array}{ccccc}
 V' & & & & \\
 \searrow & & & & \\
 & g & f' & & \\
 & \searrow & \searrow & & \\
 & & V^\sharp & \xrightarrow{\tilde{f}} & C' \\
 & & \downarrow & & \downarrow \\
 & & (GV)^* & \xrightarrow{(\hat{f})^*} & C^*
 \end{array}$$

to define g . The other figure commutes because (\hat{f}, f') is a Chu morphism. This gives the structure of a Chu object. The commutativity of the left hand triangle shows that the arrow (id, g) is in $\text{Chu}(\mathcal{V}, \perp)$. The commutativity of the square shows that (\hat{f}, \tilde{f}) is in $\text{Chu}(\mathcal{V}, \perp)$, while the commutativity of the upper triangle gives the factorization $(\hat{f}, f') = (\text{id}, g) \circ (\hat{f}, \tilde{f})$.

A coalgebra structure on an object (C, C') is determined by arrows $c: C \rightarrow RC$ and $c': \tilde{R}(C, C') \rightarrow C'$ such that the diagram

$$\begin{array}{ccc}
 \tilde{R} & \xrightarrow{c'} & C' \\
 \downarrow & & \downarrow \\
 (RC)^* & \xrightarrow{c^*} & C^*
 \end{array}$$

commutes. Consider the diagram

$$\begin{array}{ccc}
 \tilde{R}G(V, V^\sharp) & \xrightarrow{\quad} & \tilde{R}(C, C') \\
 \downarrow & & \swarrow \\
 & & V^\sharp \xrightarrow{\quad} C' \\
 & & \downarrow \quad \downarrow \\
 & & (GV)^* \xrightarrow{\quad} C^* \\
 \downarrow & \swarrow & \searrow \\
 (RGV)^* & \xrightarrow{\quad} & (RC)^*
 \end{array}$$

The outer square commutes because it just expresses the fact that there is a morphism $(C, C') \rightarrow (V, V^\sharp)$ is a morphism and hence so is $R(C, C') \rightarrow R(V, V^\sharp)$. The right

hand trapezoid commutes because (C, C') is an R -coalgebra and the bottom trapezoid commutes since $C \rightarrow GV$ is an R -coalgebra morphism in \mathcal{V} . It readily follows since the inner square is a pullback that there is induced a map $\tilde{R}(GV, V^\sharp) \rightarrow V^\sharp$ that makes the other two trapezoids commute. The commutation of the left one implies that (GV, V^\sharp) is an R -coalgebra and of the top and bottom one together that $(C, C') \rightarrow (GV, V^\sharp)$ is a morphism of R -coalgebras. \blacksquare

This proposition shows that the class of coalgebras of the form (GV, U) constitute a cofinal class among all those mapping to (V, V') . It is understood, of course, that the R -coalgebra structure restricts to $\rho: GV \rightarrow RGV$ on the first coordinate. We have not yet cut that class down to a set because the second component is still unrestricted.

A coalgebra of the form (GV, U) can be described as a map $U \rightarrow (GV)^*$ and a map $\tilde{R}(GV, U) \rightarrow U$ such that the square

$$\begin{array}{ccc} \tilde{R}(GV, U) & \longrightarrow & U \\ \downarrow & & \downarrow \\ (RGV)^* & \xrightarrow{(\rho GV)^*} & (GV)^* \end{array}$$

commutes. Here the arrow $\tilde{R}(GV, U) \rightarrow (RGV)^*$ is the one that results from the Chu structure on $R(GV, U)$. Another way of saying this is to say that U and $(RGV)^*$ are objects of the slice category $\mathcal{V}/(GV)^*$ and that \tilde{R} induces a functor on that category as indicated.

Therefore, let \mathcal{U} denote the category $\mathcal{V}/(GV)^*$ and let $S: \mathcal{U} \rightarrow \mathcal{U}$ be the functor defined by

$$S(U \rightarrow (GV)^*) = \tilde{R}(GV, U) \rightarrow (RGV)^* \rightarrow (GV)^*$$

Then an S -coalgebra in \mathcal{U} is exactly the same thing as an R -coalgebra of the form (GV, U) , whose first component is ρ . Since colimits in \mathcal{U} are created by the forgetful functor $\mathcal{U} \rightarrow \mathcal{V}$, it follows from Proposition 5.6 that S is an accessible category and hence that there is a small solution set for the existence of cofree S -coalgebras. In particular there is a small cofinal set for $V \rightarrow (GV)^*$, which implies that there is a cofree coalgebra generated by (V, V') . This completes the proof of Theorem 5.1.

6. The true history of Chu categories

Here is, insofar as I am aware, the true history of Chu categories. As I was preparing the preliminary work on $*$ -autonomous categories, for example, [Barr, 1976], I read a number of books on topological vector spaces, for example, [Schaeffer, 1970]. One thing these books always mentioned and usually developed in some detail was the theory of pairs of spaces. A pair (E, E') consisted of two topological vector spaces and a bilinear pairing $E \times E' \rightarrow \mathbf{K}$ (\mathbf{K} is the ground field, either the real or complex numbers). Mostly, but not

always, it was assumed that the pairing was separated and extensional. This construction was first given by Mackey, published in [1945], based on his 1942 doctoral dissertation. It is interesting to note that Mackey wrote in 1945 that the use of dual pairs, “makes it possible to regard a normable topological linear space as a linear space together with a distinguished family of linear functionals, rather than as a linear space with a topology”, which is exactly the point I have been making in [Barr, 1995] and [Barr, to appear].

Grothendieck [1973], a book based on a notes from a 1954 course, develops the idea further, dropping the extensional hypothesis that Mackey and Schaeffer assumed. The theory was developed for these objects, but, although it was evident what maps had to be, they were not made into a category. Still less, was any closed or monoidal structure considered. Nonetheless, this is an instance of $\text{Chu}(\mathcal{V}, \mathbf{K})$ and appears to have been the first. As far as I have been able to determine (based on not very extensive searching) no author actually discussed the notion of a morphism between pairs except in the separated case, where it is a subset of the set of linear maps between the first components consisting of those that induce maps on the second.

So I wrote down the obvious definition of morphism. It then occurred to me to wonder if the resultant category was closed or monoidal and quickly discovered that it was both, modulo the checking of a morass of details. Since Chu needed a topic for his master’s degree, I set him this task and the appendix to [Barr, 1979] was the result. But the story does not end there. Or rather, it would have ended there since neither I nor anyone else found any interest in this rather formal construction. But the year 1987 saw a renewed interest in $*$ -autonomous categories as models of Girard’s linear logic and then Vaughan Pratt and his student Vineet Gupta rediscovered the Chu construction and began to study it in earnest. Pratt describes this as follows.

Vineet and I had developed the category PDLat of partial distributive lattices as an extension of event structures for modeling concurrent processes, discovering along the way that it was concretely equivalent to $\text{Chu}(\mathbf{Set}, 2)$ [Pratt, 1993a,b, Gupta, 1994]. We already knew that PDLat’s were in a certain sense universal for Stone duality, and my result that the category of k -ary relational structures embeds concretely in $\text{Chu}(\mathbf{Set}, 2^k)$ led me to advance the thesis that Chu spaces were universal for concrete mathematics [Pratt, 1995]. If we regard pure first order logic as the logic just of sets as relational structures with the empty sort, and interpret linear logic propositions as Chu spaces, the implication of this thesis is that linear logic, thus far understood as a logic of resources, is better understood as the logic of the rest of concrete mathematics.

This interest eventually convinced me to look at the Chu construction again as a suitable vehicle in which to express virtually all duality theories. One of the results of this was the paper [Barr, to appear] in which much of the work of [Barr, 1979] is redone in this new light.

Probably the main conclusion to draw from this history is that, as with most mathematical discoveries, it is a mistake to attribute it any one person. It was born out of need, with a number of midwives, but no real discoverer.

References

- M. Barr (1976), *Closed categories and topological vector spaces*. Cahiers de Topologie et Géométrie Différentielle, **17**, 223–234.
- M. Barr (1979), ***-Autonomous categories**. Lecture Notes in Mathematics **752**, Springer-Verlag, Berlin, Heidelberg, New York.
- M. Barr (1990), *Accessible categories and models of linear logic*. J. Pure Applied Algebra, **69**, 219–232.
- M. Barr (1991), **-Autonomous categories and linear logic*. Math. Structures Comp. Sci., **1**, 159–178.
- M. Barr (1995), *Non-symmetric *-autonomous categories*. Theoretical Computer Science, **139** 115–130.
- M. Barr (to appear), **-Autonomous categories, revisited*. J. Pure Applied Algebra.
- C. Brown, and D. Gurr (1990), *A Categorical Linear Framework for Petri Nets*. In J. Mitchell, ed., Proceedings of Logic in Computer Science, IEEE Computer Society.
- P.-H. Chu (1979), *Constructing *-autonomous categories*. Appendix to [Barr, 1979], 103–137.
- B. Day and R. Street, *Monoidal bicategories and Hopf algebroids*. Forthcoming.
- S. Eilenberg and G.M. Kelly (1966), *Closed categories*. Proceedings of the Conference on Categorical Algebra, Springer-Verlag, Berlin, Heidelberg, New York, 421–562.
- P. Gabriel and F. Ulmer (1971), **Lokal präsentierbare Kategorien**. Lecture Notes in Mathematics **221**, Springer-Verlag, Berlin, Heidelberg, New York.
- A. Grothendieck (1973), **Topological Vector spaces**. Gordon and Breach.
- V. Gupta (1994), **Chu Spaces: A Model of Concurrency**. Ph.D. Thesis, Stanford University.
- J. Koslowski, *Monads and interpolads in bicategories*. Forthcoming.
- Y. Lafont, and T. Streicher (1991), *Games Semantics for Linear Logic*. In Proceedings 6th Annual, IEEE Symp. on Logic in Computer Science, Amsterdam.
- G. Mackey (1945), *On infinite dimensional vector spaces*. Trans. Amer. Math. Soc. **57**, 1945, 155–207.
- M. Makkai and R. Paré (1990), **Accessible Categories**. Contemporary Mathematics, **104**, Amer. Math. Soc.
- V. de Paiva (1989), *A Dialectica-like Model of Linear Logic*. In Proceedings Conf. on Category Theory and Computer Science, Manchester, Springer-Verlag, Berlin, Heidelberg, New York.

- D. Pavlović (1993), *The Chu construction and cofree models of linear logic*. Preprint, McGill University.
- V.R. Pratt (1993a), *Linear Logic for Generalized Quantum Mechanics*. In Proceedings Workshop on Physics and Computation, Dallas, IEEE Computer Society.
- V.R. Pratt (1993b), *The Second Calculus of Binary Relations*. In Proceedings of MFCS'93, Gdańsk, Poland, 142–155.
- V.R. Pratt (1995), *The Stone Gamut: A Coordinatization of Mathematics*. In Proceedings of the conference Logic in Computer Science IEEE Computer Society.
- H.H. Schaeffer (1970), **Topological Vector spaces**. Springer-Verlag, Berlin, Heidelberg, New York.
- R. Street (1983), *Enriched categories and cohomology*. *Quæstiones Math.* **6**, 265–283.

Dept. of Math. and Stats.

McGill University

805 Sherbrooke St. W

Montreal, QC H3A 2K6

Email: barr@math.mcgill.ca

THEORY AND APPLICATIONS OF CATEGORIES will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Papers may be submitted to any of the editors, who are listed below. More information is available from the journal's WWW server at URL <http://www.tac.mta.ca/tac/> or by anonymous ftp from <ftp.tac.mta.ca> in the directory `pub/tac`.

Editorial policy. Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

The method of distribution of the journal is via the Internet tools `WWW/gopher/ftp`. The journal is archived electronically and in printed paper format.

Subscription information. Individual subscribers receive (by e-mail) abstracts of articles as they are published. Full text of published articles is available in .dvi and Postscript format. Details will be e-mailed to new subscribers and are available by `WWW/gopher/ftp`. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, rrosebrugh@mta.ca.

Information for authors. The typesetting language of the journal is $\text{T}_{\text{E}}\text{X}$, and $\text{L}_{\text{A}}\text{T}_{\text{E}}\text{X}$ is the preferred flavour. $\text{T}_{\text{E}}\text{X}$ source of articles for publication should be submitted by e-mail directly to an appropriate Editor. Please obtain detailed information on submission format and style files by WWW or anonymous ftp from the sites listed above. You may also write to tac@mta.ca to receive details by e-mail.

Editorial board.

John Baez, University of California, Riverside: baez@math.ucr.edu
Michael Barr, McGill University: barr@triples.math.mcgill.ca
Lawrence Breen, Université de Paris 13: breen@math.univ-paris.fr
Ronald Brown, University of North Wales: r.brown@bangor.ac.uk
Jean-Luc Brylinski, Pennsylvania State University: jlb@math.psu.edu
Aurelio Carboni, University of Genoa: carboni@vmimat.mat.unimi.it
P. T. Johnstone, University of Cambridge: P.T.Johnstone@pmms.cam.ac.uk
G. Max Kelly, University of Sydney: kelly_m@maths.su.oz.au
Anders Kock, University of Aarhus: kock@mi.aau.dk
F. William Lawvere, State University of New York at Buffalo: mthfwl@ubvms.cc.buffalo.edu
Jean-Louis Loday, Université de Strasbourg: loday@math.u-strasbg.fr
Ieke Moerdijk, University of Utrecht: moerdijk@math.ruu.nl
Susan Niefield, Union College: niefiels@gar.union.edu
Robert Paré, Dalhousie University: pare@cs.dal.ca
Andrew Pitts, University of Cambridge: ap@c1.cam.ac.uk
Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca
Jiri Rosicky, Masaryk University: rosicky@math.muni.cz
James Stasheff, University of North Carolina: jds@charlie.math.unc.edu
Ross Street, Macquarie University: street@macadam.mpce.mq.edu.au
Walter Tholen, York University: tholen@vm1.yorku.ca
Myles Tierney, Rutgers University: tierney@math.rutgers.edu
Robert F. C. Walters, University of Sydney: walters.b@maths.su.oz.au
R. J. Wood, Dalhousie University: rjwood@cs.da.ca