

556: MATHEMATICAL STATISTICS I

SOME NOTES ON CHARACTERISTIC FUNCTIONS

The characteristic function for a random variable X with pmf/pdf f_X is defined for $t \in \mathbb{R}$ as

$$\begin{aligned}\varphi_X(t) &= \mathbb{E}_X[e^{itX}] = \mathbb{E}_X[\cos(tX) + i \sin(tX)] \\ &= \mathbb{E}_X[\cos(tX)] + i \mathbb{E}_X[\sin(tX)].\end{aligned}$$

In general $\varphi_X(t)$ is a complex-valued function. If X is discrete, taking values on $\mathbb{X} = \{x_1, x_2, \dots\}$

$$\mathbb{E}_X[\cos(tX)] = \sum_{j=1}^{\infty} \cos(tx_j) f_X(x_j)$$

$$\mathbb{E}_X[\sin(tX)] = \sum_{j=1}^{\infty} \sin(tx_j) f_X(x_j)$$

Now,

$$\sum_{j=1}^{\infty} \cos(tx_j) f_X(x_j) \leq \left| \sum_{j=1}^{\infty} \cos(tx_j) f_X(x_j) \right| \leq \sum_{j=1}^{\infty} |\cos(tx_j)| f_X(x_j) \leq \sum_{j=1}^{\infty} f_X(x_j) = 1$$

with a similar result for \sin , so the two expectations are finite, so $\varphi_X(t)$ exists. The same argument works for X continuous, where

$$\varphi_X(t) = \int_{-\infty}^{\infty} e^{itx} f_X(x) dx = \int_{-\infty}^{\infty} \cos(tx) f_X(x) dx + i \int_{-\infty}^{\infty} \sin(tx) f_X(x) dx$$

EXAMPLE Double-Exponential (or Laplace) distribution

$$f_X(x) = \frac{1}{2} e^{-|x|} \quad x \in \mathbb{R}$$

Then

$$\varphi_X(t) = \int_{-\infty}^{\infty} e^{itx} \frac{1}{2} e^{-|x|} dx = \int_0^{\infty} \cos(tx) e^{-x} dx. \quad (1)$$

Integrating by parts we have

$$\begin{aligned}\varphi_X(t) &= [-\cos(tx) e^{-x}]_0^{\infty} + \int_0^{\infty} t \sin(tx) e^{-x} dx \\ &= 1 + [-t \sin(tx) e^{-x}]_0^{\infty} - \int_0^{\infty} t^2 \cos(tx) e^{-x} dx \\ &= 1 - t^2 \varphi_X(t)\end{aligned}$$

Therefore

$$\varphi_X(t) = \frac{1}{1+t^2}$$

EXAMPLE **Normal distribution**

$$f_X(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2} \quad x \in \mathbb{R}$$

Then

$$\varphi_X(t) = \int_{-\infty}^{\infty} e^{itx} \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx.$$

Completing the square

$$\varphi_X(t) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-(x-it)^2/2} e^{-t^2/2} dx.$$

Therefore

$$\varphi_X(t) = e^{-t^2/2}.$$

The following results also hold:

- $\varphi_X(t)$ is **continuous** for all t ; this follows as \cos and \sin are continuous functions of x , and sums and integrals of continuous functions are also continuous. In fact, we can prove the stronger result that $\varphi_X(t)$ is **uniformly continuous** on \mathbb{R} .
- $\varphi_X(t)$ is **bounded in modulus** by 1, as

$$|\varphi_X(t)| \leq \mathbb{E}_X[|e^{itX}|] = \mathbb{E}_X[1] = 1$$

- The derivatives of $\varphi_X(t)$ are not guaranteed to be finite; we can consider

$$\varphi_X^{(r)}(t) = \frac{d^r}{dt^r} \{\varphi_X(t)\}$$

but this quantity may not be defined, or finite, at any given t ; if $r = 1$

$$\varphi_X^{(1)}(t) = \mathbb{E}_X[-X \sin(tX)] + i\mathbb{E}_X[X \cos(tX)].$$

but there is no guarantee that either expectation is finite. For example, for the Cauchy distribution

$$\varphi_X(t) = e^{-|t|}$$

which undefined derivative at $t = 0$.

INVERSION FORMULA

A general inversion formula in 1-D gives the method via which f_X or F_X can be computed from φ_X .

- Let $\bar{F}_X(x)$ be defined by

$$\bar{F}_X(x) = \frac{1}{2} \left\{ F_X(x) + \lim_{y \rightarrow x^-} F_X(y) \right\}.$$

Then for $a < b$

$$\bar{F}_X(b) - \bar{F}_X(a) = \frac{1}{2\pi} \lim_{T \rightarrow \infty} \int_{-T}^T \left(\frac{e^{-iat} - e^{-ibt}}{it} \right) \varphi_X(t) dt$$

- For an alternative statement, let a and $a + h$ for $h > 0$ be continuity points of F_X . Then

$$F_X(a + h) - F_X(a) = \frac{1}{2\pi} \lim_{T \rightarrow \infty} \int_{-T}^T \left(\frac{1 - e^{-ith}}{it} \right) e^{-ita} \varphi_X(t) dt$$

In certain circumstances we may compute f_X from φ_X more straightforwardly.

(I) If X is **discrete** taking values on the integers. Then

$$\varphi_X(t) = \sum_{x=-\infty}^{\infty} e^{itx} f_X(x).$$

For integer j

$$\int_{-\pi}^{\pi} e^{i(j-x)t} dt = \begin{cases} 2\pi & \text{if } x = j \\ 0 & \text{if } x \neq j \end{cases}$$

Thus for any fixed x

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ixt} \varphi_X(t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ixt} \left\{ \sum_{j=-\infty}^{\infty} e^{itj} f_X(j) \right\} dt = \frac{1}{2\pi} \sum_{j=-\infty}^{\infty} \left\{ \int_{-\pi}^{\pi} e^{i(j-x)t} dt \right\} f_X(j) = f_X(x)$$

(as only the term when $j = x$ is non-zero in the sum) so we have the inversion formula: for $x \in \mathbb{Z}$

$$f_X(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ixt} \varphi_X(t) dt.$$

(II) If X is **continuous** and **absolutely integrable**

$$\int_{-\infty}^{\infty} |\varphi_X(t)| dt < \infty$$

then

$$f_X(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-itx} \varphi_X(t) dt$$

EXAMPLE Suppose that for $t \in \mathbb{R}$,

$$\varphi_X(t) = e^{-|t|}.$$

Clearly this function is absolutely integrable, so we have

$$\begin{aligned} f_X(x) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-itx} e^{-|t|} dt = \frac{1}{\pi} \int_0^{\infty} \cos(tx) e^{-t} dt \\ &= \frac{1}{\pi} \frac{1}{1+x^2} \end{aligned}$$

by the result in equation (1). Hence $X \sim \text{Cauchy}$.

DIAGNOSING DISCRETE OR CONTINUOUS DISTRIBUTIONS

(I) If

$$\limsup_{|t| \rightarrow \infty} |\varphi_X(t)| = 1$$

then X is often a **discrete** random variable. Technically, X may also have a **singular** distribution: see, or example

but such distributions are rarely encountered in practice.

(II) If

$$\limsup_{|t| \rightarrow \infty} |\varphi_X(t)| = 0$$

then X is **continuous**; consequently, if

$$\lim_{|t| \rightarrow \infty} |\varphi_X(t)| = 0$$

then X is continuous.

INTERPRETING THE CHARACTERISTIC FUNCTION.

To get a further understanding of characteristic function, we consider the inversion formulae. For discrete random variables defined on the integers, we have

$$f_X(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ixt} \varphi_X(t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} [\cos(xt) - i \sin(xt)] \varphi_X(t) dt$$

One way to think about this integral is via a discrete approximation; fix

$$t_{j,N} = -\pi + \frac{2\pi j}{N} \quad j = 0, 1, 2, \dots, N$$

and write

$$f_X(x) \simeq \frac{1}{2\pi} \left\{ \sum_{j=0}^N \cos(xt_{j,N}) \varphi_X(t_{j,N}) - i \sum_{j=0}^N \sin(xt_{j,N}) \varphi_X(t_{j,N}) \right\}$$

(I) Suppose f_X is **degenerate** at x_0 , that is,

$$f_X(x) = \begin{cases} 1 & x = x_0 \\ 0 & x \neq x_0 \end{cases}$$

Then by elementary calculations

$$\varphi_X(t) = \cos(x_0 t) + i \sin(x_0 t)$$

so that

$$\operatorname{Re}(\varphi_X(t)) = \cos(x_0 t) \quad \operatorname{Im}(\varphi_X(t)) = \sin(x_0 t)$$

that is, pure sinusoids with period $2\pi/x_0$.

(II) Suppose f_X is **discrete**, then as above

$$\varphi_X(t) = \sum_{j=1}^{\infty} \cos(tx_j) f_X(x_j) + i \sum_{j=1}^{\infty} \sin(tx_j) f_X(x_j)$$

so that

$$\operatorname{Re}(\varphi_X(t)) = \sum_{j=1}^{\infty} \cos(tx_j) f_X(x_j) \quad \operatorname{Im}(\varphi_X(t)) = \sum_{j=1}^{\infty} \sin(tx_j) f_X(x_j)$$

that is, a weighted sum of pure sinusoids with period $2\pi/x_1, 2\pi/x_2, \dots$, with weights determined by f_X