

## 556: MATHEMATICAL STATISTICS I

### NON 1-1 TRANSFORMATIONS

Suppose that  $X$  is a continuous r.v. with range  $\mathbb{X} \equiv (0, 2\pi)$  whose pdf  $f_X$  is constant

$$f_X(x) = \frac{1}{2\pi} \quad 0 < x < 2\pi$$

and zero otherwise. This pdf has corresponding continuous cdf

$$F_X(x) = \frac{x}{2\pi} \quad 0 < x < 2\pi$$

**Example 1** Consider transformed r.v.  $Y = \sin X$ . Then the range of  $Y$ ,  $\mathbb{Y}$  is  $[-1, 1]$ , but the transformation is not 1-1. However, from first principles, we have

$$F_Y(y) = P[Y \leq y] = P[\sin X \leq y]$$

Now, by inspection of Figure 1, we can easily identify the required set  $A_y$  : it is the union of **two** disjoint intervals

$$A_y = [0, x_1] \cup [x_2, 2\pi] = [0, \arcsin y] \cup [\pi - \arcsin y, 2\pi]$$




Figure 1: Computation of  $A_y$  for  $Y = \sin X$

$$F_Y(y) = P[\sin X \leq y] = P[X \leq x_1] + P[X \geq x_2] = \{P[X \leq x_1]\} + \{1 - P[X < x_2]\}$$

$$= \left\{ \frac{1}{2\pi} \arcsin y \right\} + \left\{ 1 - \frac{1}{2\pi} (\pi - \arcsin y) \right\} = \frac{1}{2} + \frac{1}{\pi} \arcsin y$$

and hence, by differentiation

$$f_Y(y) = \frac{1}{\pi} \frac{1}{\sqrt{1-y^2}} \quad 0 \leq y \leq 1$$

and zero otherwise.

**Example 2** Consider transformed r.v.  $Y = \sin^2 X$ . Then the range of  $Y, \mathbb{Y}$ , is  $[0, 1]$ , but the transformation is not 1-1. However, from first principles, we have

$$F_Y(y) = P[Y \leq y] = P[\sin^2 X \leq y]$$

In Figure 2, we identify the required set  $A_y$  : it is the union of **three** disjoint intervals

$$A_y = [0, x_1] \cup [x_2, x_3] \cup [x_4, 2\pi]$$

where

$$x_1 = \arcsin(\sqrt{y}) \quad x_2 = \pi - \arcsin(\sqrt{y}) \quad x_3 = \pi + \arcsin(\sqrt{y}) \quad x_4 = 2\pi - \arcsin(\sqrt{y})$$




Figure 2: Computation of  $A_y$  for  $Y = \sin^2 X$

$$\begin{aligned} F_Y(y) &= P[\sin^2 X \leq y] = P[X \leq x_1] + P[x_2 < X \leq x_3] + P[x_4 < X \leq 2\pi] \\ &= F_X(x_1) + \{F_X(x_3) - F_X(x_2)\} + \{1 - F_X(x_4)\} \\ &= \frac{x_1}{2\pi} + \left\{ \frac{x_3}{2\pi} - \frac{x_2}{2\pi} \right\} + \left\{ 1 - \frac{x_4}{2\pi} \right\} = \frac{2}{\pi} \arcsin(\sqrt{y}) \end{aligned}$$

and hence, by differentiation

$$f_Y(y) = \frac{1}{\pi} \frac{1}{\sqrt{(1-y)y}} \quad 0 \leq y \leq 1$$

and zero otherwise.

**Example 3** Consider transformed r.v.  $T = \tan X$ . Then the range of  $T$ ,  $\mathbb{T}$  is  $\mathbb{R}$ , but the transformation is not 1-1. However, from first principles, we have, for  $t > 0$

$$F_T(t) = P[T \leq t] = P[\tan X \leq t]$$

Figure 3 helps identify the required set  $A_t$ : in this case, it is the union of three disjoint intervals

$$A_t = [0, x_1] \cup \left[\frac{\pi}{2}, x_2\right] \cup \left[\frac{3\pi}{2}, 2\pi\right] = [0, \tan^{-1} t] \cup \left[\frac{\pi}{2}, \pi + \tan^{-1} t\right] \cup \left[\frac{3\pi}{2}, 2\pi\right]$$

(note, for values of  $t < 0$ , the union will be of only two intervals, but the calculation proceeds identically) so that




Figure 3: Computation of  $A_t$  for  $T = \tan X$

$$\begin{aligned} F_T(t) &= P[\tan X \leq t] = P[X \leq x_1] + P\left[\frac{\pi}{2} \leq X \leq x_2\right] + P\left[\frac{3\pi}{2} \leq X \leq 2\pi\right] \\ &= \left\{\frac{1}{2\pi} \tan^{-1} t\right\} + \frac{1}{2\pi} \left\{\pi + \tan^{-1} t - \frac{\pi}{2}\right\} + \frac{1}{2\pi} \left\{2\pi - \frac{3\pi}{2}\right\} = \frac{1}{\pi} \tan^{-1} t + \frac{1}{2} \end{aligned}$$

and hence, by differentiation

$$f_T(t) = \frac{1}{\pi} \frac{1}{1+t^2} \quad t \in \mathbb{R}$$