Solutions of Assignment 8
Basic Algebra I

November 11, 2004

Solution of the problem 1. Recall that a field F has only two ideals: $\{0\}$, F. Also recall that the kernel of any ring homomorphism is an ideal. Now back to the problem, in order to show that f is injective, it is enough to show that $\ker(f) = \{0\}$. If not, then $\ker(f) = F$. So $1 \in \ker(f)$, i.e., $f(1) = 0$, which is a contradiction. Thus f is injective. The first isomorphism theorem now implies the other part of the problem:

$$F \cong F/\{0\} \cong F/\ker(f) \cong f(F).$$

Solution of the problem 2. Let $f : \mathbb{Z} \longrightarrow R$ be a ring homomorphism from $\mathbb{Z}$ to an arbitrary ring R. If $\ker(f) = \{0\}$, then the argument given in the previous problem shows that $\mathbb{Z}$ is isomorphic to its image under f. And if $\ker(f) \neq \{0\}$, then it is of the form $d\mathbb{Z}$, for some $d > 0$. Once again, the first isomorphism theorem implies that the image of $\mathbb{Z}$ under f is isomorphic to $\mathbb{Z}/d\mathbb{Z} \cong \mathbb{Z}_d$, which is a finite ring with d elements.

Solution of the problem 3. This is false. For example, $\mathbb{Z}$ is an integral domain, however, its quotient by the ideal $6\mathbb{Z}$, namely $\mathbb{Z}_6$, is not an integral domain.

Solution of the problem 4. This is true. Let J be an ideal of R/I. Recall that the natural homomorphism $\pi : R \longrightarrow R/I,\ \pi(a) = a + I$, is a surjective ring homomorphism. We now claim that the inverse image $\pi^{-1}(J) := \{a \in R :\ \pi(a) \in J\}$ is an ideal of R:

If $a, b \in \pi^{-1}(J)$, then $\pi(a + b) = \pi(a) + \pi(b) \in J$, so $a + b \in \pi^{-1}(J)$.

If $a \in \pi^{-1}(J),\ r \in R$, then $\pi(ra) = \pi(r)\pi(a) \in J$, so $ra \in \pi^{-1}(J)$.

Every ideal of R is assumed to be principal, so $\pi^{-1}(J) = (a_0) = a_0R$, for some $a_0 \in R$. Now since π is onto, we conclude that

$$J = \pi(\pi^{-1}(J)) = \pi((a_0R)) = \{\pi(a_0r) :\ r \in R\} = \{a_0r + I :\ r \in R\}$$

$$= \{(a_0 + I)(r + I) :\ r \in R\} = (a_0 + I).$$

This means that J is generated by the element $a_0 + I$. Done.

Solution of the problem 5. False. Let $R = \mathbb{Z}[x]$ and let $I = (x)$, the ideal generated by x. We first claim that $R/I \cong \mathbb{Z}$. To see this, define

$$\phi : R \longrightarrow \mathbb{Z},\ \phi(f(x)) = f(0).$$

It is apparent that ϕ is a ring homomorphism. ϕ is also surjective (every integer can be regarded as a polynomial). Also note that

$$\ker(\phi) = \{f(x) :\ \phi(f(x)) = 0\} = \{f(x) :\ f(0) = 0\} = \{f(x) :\ x \mid f(x)\} = I.$$

So, $R/I \cong \mathbb{Z}$, and the claim is proved.

Since every ideal of $\mathbb{Z}$ is principal, this in fact shows that every ideal of R/I is so. We now show that the same is not true for R by showing that the ideal $J = \{f(x) :\ 2 \mid f(0)\}$ is not principal (it is left to you to check that J is in fact an ideal). On the contrary, suppose that J is generated by some polynomial $g(x)$. Since $2, x \in J$, we would have $g(x) \mid 2,\ g(x) \mid x$. So, $g(x) = \pm 1$, which is a contradiction (why?).

Solution of the problem 6. Our first claim is that for **any prime** p,

$$\frac{\mathbb{Z}[x]}{(p, x^2+1)} \cong \frac{\mathbb{Z}_p[x]}{(x^2+1)}.$$

To see this, define $\phi : \mathbb{Z}[x] \longrightarrow \dfrac{\mathbb{Z}_p[x]}{(x^2+1)}$ by the rule

$$\phi(a_0 + a_1 x + \cdots + a_n x^n) = \bar{a}_0 + \bar{a}_1 x + \cdots + \bar{a}_n x^n + (x^2+1),$$

where $\bar{a}$ denotes the congruence class of a mod p. It is readily seen that ϕ is a surjective ring homomorphism (check this!). To find the kernel, notice that since any $f(x)$ can be written as $f(x) = a + bx + g(x)(x^2+1)$ for some $g(x)$ (division algorithm), so $f(x)$ is in the kernel $\iff \bar{a} + \bar{b}x = 0 \iff p \mid a, p \mid b \iff f(x) \in (p, x^2+1)$. The first isomorphism theorem now concludes the proof of our first claim.

Now we specialize to the case where $p = 5$ or $p = 7$.

(I) For $p = 5$, we have the factorization $x^2 + 1 = (x-3)(x-2)$. Let us now define

$$\psi : \mathbb{Z}_5[x] \longrightarrow \mathbb{Z}_5 \times \mathbb{Z}_5,\ \psi(f(x)) = (f(3), f(2)).$$

ψ is clearly a ring homomorphism with the kernel

$$\begin{aligned}
\ker(\psi) &= \{f(x) :\ f(3) = f(2) = 0\} \\
&= \{f(x) :\ x-3 \mid f(x),\ x-2 \mid f(x)\} \\
&= \{f(x) :\ x^2+1 \mid f(x)\} \\
&= (x^2+1).
\end{aligned}$$

It remains to show that ψ is surjective. Given any $(\alpha,\ \beta) \in \mathbb{Z}_5 \times \mathbb{Z}_5$, take $f(x) = (3\beta - 2\alpha) + (\alpha - \beta)x$. We then have

$$\begin{aligned}
\psi(f(x)) &= (f(3),\ f(2)) \\
&= (3\beta - 2\alpha + 3\alpha - 3\beta,\ 3\beta - 2\alpha + 2\alpha - 2\beta) \\
&= (\alpha,\ \beta).
\end{aligned}$$

Hence, by the first isomorphism theorem, we deduce that

$$\frac{\mathbb{Z}[x]}{(5, x^2+1)} \cong \frac{\mathbb{Z}_5[x]}{(x^2+1)} \cong \mathbb{Z}_5 \times \mathbb{Z}_5.$$

(II) Now suppose that $p = 7$. In contrast to 5, $x^2 + 1$ does not factor in $\mathbb{Z}_7[x]$, i.e., it is irreducible. Now we claim that $\dfrac{\mathbb{Z}_7[x]}{(x^2+1)}$ is a field. To prove this, we have to show that every nonzero class has an inverse. So, suppose that $f(x) \notin (x^2+1)$. Thus $\gcd(f(x),\ x^2+1) = 1$, and since $\mathbb{Z}_7$ is a field, we can find $g(x), h(x) \in \mathbb{Z}_7[x]$ so that $f(x)g(x) + h(x)(x^2+1) = 1$. Therefore $(f(x) + (x^2+1))(g(x) + (x^2+1)) = 1 + (x^2+1)$. In other words, the class $g(x) + (x^2+1)$ is the inverse of $f(x) + (x^2+1)$. And finally we count the number of classes in $\dfrac{\mathbb{Z}_7[x]}{(x^2+1)}$. Since every class has a unique representative of the form $a + bx + (x^2+1)$ with $0 \le a, b \le 6$ (could you explain why?), we conclude that the total number of classes is $7 \times 7 = 49$. Done!

Extra Credit

Solution of the problem 7. As usual, we define the right map and will exploit it to conclude the desired result. So, consider the

$$\phi : F[[x]] \longrightarrow F,\ \phi(\sum_{n=0}^{\infty} a_n x^n) = a_0.$$

Now we check in details that ϕ is a surjective ring homomorphism.
(i) ϕ respects addition:

$$\begin{aligned}
\phi\left(\sum_{n=0}^{\infty} a_n x^n + \sum_{n=0}^{\infty} b_n x^n\right) &= \phi\left(\sum_{n=0}^{\infty} (a_n + b_n) x^n\right) \\
&= a_0 + b_0 \\
&= \phi\left(\sum_{n=0}^{\infty} a_n x^n\right) + \phi\left(\sum_{n=0}^{\infty} b_n x^n\right).
\end{aligned}$$

(ii) ϕ respects multiplication:

$$\phi\left(\sum_{n=0}^{\infty} a_n x^n \cdot \sum_{n=0}^{\infty} b_n x^n\right) = \phi\left(\sum_{n=0}^{\infty} (a_0 b_n + a_1 b_{n-1} + \cdots + a_n b_0)\, x^n\right)$$

$$
\begin{aligned}
&= a_0 \cdot b_0 \\
&= \phi\left(\sum_{n=0}^{\infty} a_n x^n\right) \cdot \phi\left(\sum_{n=0}^{\infty} b_n x^n\right).
\end{aligned}
$$

(iii) The identity element of the ring $F[[x]]$ is the formal power series

$$1 = 1 + 0x + 0x^2 + 0x^3 + \cdots,$$

and we have $\phi(1) = 1$.

(iv) ϕ is surjective: for any $a \in F$, we have

$$\phi(a + 0x + 0x^2 + 0x^3 + \cdots) = a.$$

(v) The kernel of ϕ is the ideal generated by x:

$$f(x) = \sum_{n=0}^{\infty} a_n x^n \in \ker(\phi) \iff a_0 = 0 \iff f(x) = xg(x) \iff f(x) \in (x).$$

Therefore, the first isomorphism theorem implies that

$$R = \frac{F[[x]]}{(x)} \cong F.$$

To prove the second part, just note that by the isomorphism established above, every element of R and off the ideal (x) corresponds to a nonzero element of a field, hence it is invertible. And now the last part is immediate: if an ideal of R is not contained in $I = (x)$, it has to have an invertible element, hence it is the whole ring $F[[x]]$. Done!