

Chapter 5: Probability inequalities and theorems

Notes

Markov’s inequality

Suppose that $g(Y)$ is a function of continuous random variable Y such that

$$\mathbb{E}[|g(Y)|] < \infty.$$

For any $k > 0$, let

$$\mathcal{A}_k = \{y \in \mathbb{R} : |g(y)| \geq k\}.$$

Notes

Markov’s inequality (cont.)

Then

$$\begin{aligned} \mathbb{E}[|g(Y)|] &= \int_{-\infty}^{\infty} |g(y)|f_Y(y) \, dy \\ &= \int_{\mathcal{A}_k} |g(y)|f_Y(y) \, dy + \int_{\mathcal{A}'_k} |g(y)|f_Y(y) \, dy \\ &\geq \int_{\mathcal{A}_k} |g(y)|f_Y(y) \, dy \\ &\geq \int_{\mathcal{A}_k} k f_Y(y) \, dy \\ &= k \int_{\mathcal{A}_k} f_Y(y) \, dy \\ &= kP(|g(Y)| \geq k) \end{aligned}$$

Notes

Markov's inequality (cont.)

Thus

$$\mathbb{E}[|g(Y)|] \geq kP(|g(Y)| \geq k)$$

or equivalently

$$P(|g(Y)| \geq k) \leq \frac{\mathbb{E}[|g(Y)|]}{k}.$$

Notes

Markov's inequality (cont.)

Thus

$$1 - P(|g(Y)| \geq k) \geq 1 - \frac{\mathbb{E}[|g(Y)|]}{k}$$

or equivalently

$$P(|g(Y)| < k) \geq 1 - \frac{\mathbb{E}[|g(Y)|]}{k}.$$

Note that the result is identical for a discrete random variable.

Notes

Chebychev's inequality

aka

- Chebyshev's inequality
- Bienaymé-Chebyshev inequality
- Tchebysheff's Theorem
- Chebysheff, Chebychov, Chebyshev, Tchebychev ...

Notes

Chebychev’s inequality (cont.)

Suppose that Y has finite mean μ and variance σ^2 . In Markov’s inequality, consider the function

$$g(t) = (t - \mu)^2.$$

We have for any $k > 0$ that

$$P(|Y - \mu|^2 < k) \geq 1 - \frac{\mathbb{E}[|(Y - \mu)^2|]}{k}.$$

which can be more simply written as

$$P((Y - \mu)^2 < k) \geq 1 - \frac{\mathbb{E}[(Y - \mu)^2]}{k}.$$

Notes

Chebychev’s inequality (cont.)

But

$$\mathbb{E}[(Y - \mu)^2] = \sigma^2$$

and if we choose to write $k = c^2\sigma^2$, the inequality becomes

$$P((Y - \mu)^2 < c^2\sigma^2) \geq 1 - \frac{\mathbb{E}[(Y - \mu)^2]}{c^2\sigma^2} = 1 - \frac{1}{c^2}$$

Notes

Chebychev’s inequality (cont.)

The left hand side of the inequality is equal to

$$P(\sqrt{(Y - \mu)^2} < \sqrt{c^2\sigma^2}) = P(|Y - \mu| < c\sigma)$$

so the inequality becomes

$$P(|Y - \mu| < c\sigma) \geq 1 - \frac{1}{c^2}.$$

Notes

Chebychev’s inequality (cont.)

An equivalent way to write this is

$$P(|Y - \mu| \geq c\sigma) \leq \frac{1}{c^2}$$

which holds for arbitrary $c > 0$.

Notes

Chebychev’s inequality (cont.)

This states that for **any** distribution with finite mean μ and variance σ^2 , we have that

$$P(\mu - c\sigma < Y < \mu + c\sigma) \geq 1 - \frac{1}{c^2}$$

which bounds how much probability can be present in the tails of the distribution.

Notes

The sample mean

Suppose that $Y_1, \dots, Y_n$ are independent random variables that have the same distribution, and have expectation μ and variance σ^2 . Recall that if

$$U = \sum_{i=1}^n Y_i$$

then

$$\mathbb{E}[U] = n\mu \qquad \mathbb{V}[U] = n\sigma^2.$$

Notes

The sample mean (cont.)

Let

$$\overline{Y}_n = \frac{1}{n} \sum_{i=1}^n Y_i$$

be the *sample mean* random variable. Then

$$\mathbb{E}[\overline{Y}_n] = \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n Y_i\right] = \frac{1}{n} \mathbb{E}\left[\sum_{i=1}^n Y_i\right] = \frac{1}{n} n\mu = \mu$$

and

$$\mathbb{V}[\overline{Y}_n] = \mathbb{V}\left[\frac{1}{n} \sum_{i=1}^n Y_i\right] = \frac{1}{n^2} \mathbb{V}\left[\sum_{i=1}^n Y_i\right] = \frac{1}{n^2} n\sigma^2 = \frac{\sigma^2}{n}.$$

Notes

The sample mean (cont.)

Now recall Chebychev's inequality:

$$P(|Y - \mu| \geq c\sigma) \leq \frac{1}{c^2}.$$

Applying this to $\overline{Y}_n$, we have that

$$P\left(|\overline{Y}_n - \mu| \geq \frac{c\sigma}{\sqrt{n}}\right) \leq \frac{1}{c^2}.$$

Notes

The sample mean (cont.)

We can rewrite this for any n by setting

$$\epsilon = \frac{c\sigma}{\sqrt{n}} \quad \text{so that} \quad c = \frac{\sqrt{n}\epsilon}{\sigma}.$$

Then the inequality becomes

$$P(|\overline{Y}_n - \mu| \geq \epsilon) \leq \frac{\sigma^2}{n\epsilon^2}.$$

Notes

Convergence in probability

Consider a sequence of random variables $Y_1, Y_2, \dots, Y_n, \dots$, and real constant c . We say that the sequence *converges in probability* to c as $n \rightarrow \infty$ if, for every $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} P(|Y_n - c| < \epsilon) = 1$$

that is, the probability

$$P(c - \epsilon < Y_n < c + \epsilon)$$

converges to 1 as $n \rightarrow \infty$. We write

$$Y_n \xrightarrow{p} c$$

as $n \rightarrow \infty$ to indicate convergence in probability.

Notes

Convergence in probability (cont.)

Note

- If, for a fixed $\epsilon > 0$,

$$p_n = P(|Y_n - c| < \epsilon)$$

the definition requires that the real sequence $\{p_n\}$ converges to 1 in the usual sense.

- There is no restriction on the joint distribution of the Y_i s in the sequence (that is, they can be dependent).
- We can equivalently write the condition

$$\lim_{n \rightarrow \infty} P(|Y_n - c| > \epsilon) = 0.$$

Notes

The weak law of large numbers

Suppose that $Y_1, Y_2, \dots, Y_n$, are independent random variables that have the same distribution, and have expectation μ and variance σ^2 . For each n , let

$$\bar{Y}_n = \frac{1}{n} \sum_{i=1}^n Y_i$$

be the sample mean random variable. Then

$$\bar{Y}_n \xrightarrow{p} \mu$$

as $n \rightarrow \infty$.

Notes

The weak law of large numbers (cont.)

This follows from Chebychev's inequality: recall that for the sample mean

$$P(|\bar{Y}_n - \mu| \geq \epsilon) \leq \frac{\sigma^2}{n\epsilon^2}.$$

or equivalently

$$P(|\bar{Y}_n - \mu| < \epsilon) \geq 1 - \frac{\sigma^2}{n\epsilon^2}.$$

for any $\epsilon > 0$.

Notes

The weak law of large numbers (cont.)

But

$$1 - \frac{\sigma^2}{n\epsilon^2} \longrightarrow 1$$

as $n \longrightarrow \infty$, as σ and ϵ are fixed. Thus

$$\lim_{n \longrightarrow \infty} P(|\bar{Y}_n - \mu| < \epsilon) \geq 1$$

and hence $\bar{Y}_n \xrightarrow{p} \mu$.

Notes

The weak law of large numbers (cont.)

What this result means: under the stated conditions, the sample mean random variable converges to the expectation of the distribution of the Y s.

For example, if you repeatedly and independently toss a fair coin, the proportion of heads will converge to $1/2$ as the number of tosses increases.

Notes

The weak law of large numbers (cont.)

Note

The statement

$$P(|\bar{Y}_n - \mu| < \epsilon) \geq 1 - \frac{\sigma^2}{n\epsilon^2}.$$

tells us that the *rate of convergence* is (at least) n ;

- that is, the probability on the left hand side converges to 1 at rate (at least) n .

Notes

Convergence in distribution

Consider the sequence of cdfs, $F_1(y), F_2(y), \dots, F_n(y), \dots$, and suppose that

$$\lim_{n \rightarrow \infty} F_n(y) = F(y)$$

for some other cdf $F(y)$ at all points where $F(y)$ is **continuous**.

Then $F(y)$ is the *limiting distribution* for the sequence.

Notes

Convergence in distribution (cont.)

If $Y_1, Y_2, \dots, Y_n, \dots$ are a sequence of random variables having cdfs $F_1(y), F_2(y), \dots, F_n(y), \dots$ respectively, and Y has cdf $F(y)$, then we say that the sequence $Y_1, Y_2, \dots, Y_n, \dots$ *converges in distribution* to Y ,

$$Y_n \xrightarrow{d} Y$$

as $n \rightarrow \infty$.

Notes

The central limit theorem

Suppose that $Y_1, Y_2, \dots, Y_n$, are independent random variables that have the same distribution, and have finite expectation μ and variance σ^2 . For each n , let

$$\overline{Y}_n = \frac{1}{n} \sum_{i=1}^n Y_i$$

be the sample mean random variable.

Notes

The central limit theorem (cont.)

Then the random variable

$$U_n = \frac{\sqrt{n}(\overline{Y}_n - \mu)}{\sigma}$$

converges in distribution to a standard normal distribution.

That is,

$$\lim_{n \rightarrow \infty} F_{U_n}(u) = \lim_{n \rightarrow \infty} P(U_n \leq u) = \int_{-\infty}^u \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt$$

This result is the *Central Limit Theorem* (CLT).

Notes

The central limit theorem (cont.)

Note that we could write

$$U_n = \frac{(\overline{Y}_n - \mu)}{\sigma/\sqrt{n}}$$

or, multiplying numerator and denominator by n

$$U_n = \frac{\left(\sum_{i=1}^n Y_i - n\mu\right)}{\sqrt{n}\sigma}$$

Notes

The central limit theorem (cont.)

The result says that, provided the conditions on expectation and variance are met, when n is large, we can approximate the distribution of $\bar{Y}_n$, or of

$$S_n = \sum_{i=1}^n Y_i$$

using a Normal distribution, **irrespective** of the actual distribution of $Y_1, \dots, Y_n$.

Notes

The central limit theorem (cont.)

That is,

- $\bar{Y}_n$ is approximately distributed as

$$Normal(\mu, \sigma^2/n)$$

- S_n is approximately distributed as

$$Normal(n\mu, n\sigma^2)$$

Notes

The central limit theorem (cont.)

Note

If $Y_1, \dots, Y_n$ are independent and identically distributed Normal random variables with expectation μ and variance σ^2 , then

- $\bar{Y}_n$ is **exactly** distributed as

$$Normal(\mu, \sigma^2/n)$$

- S_n is **exactly** distributed as

$$Normal(n\mu, n\sigma^2)$$

Notes

The central limit theorem (cont.)

The result is straightforward to prove using moment-generating functions. For $i = 1, 2, \dots, n$, let

$$Z_i = \frac{Y_i - \mu}{\sigma}.$$

Then from properties of expectations and variances, we have that

$$\mathbb{E}[Z_i] = 0 \qquad \mathbb{V}[Z_i] = 1.$$

and also

$$U_n = \frac{1}{\sqrt{n}} \sum_{i=1}^n Z_i$$

Notes

The central limit theorem (cont.)

If we assume that $m(t)$ is the mgf of each Y_i , and $m_Z(t)$ is the mgf for each Z_i , then by previous mgf results

$$m_{U_n}(t) = \left\{ m_Z \left(\frac{t}{\sqrt{n}} \right) \right\}^n$$

Now, using a **Taylor expansion** of $m_Z(t)$ at $t = 0$

$$m_Z(t) = m_Z(0) + t m_Z^{(1)}(0) + \frac{t^2}{2} m_Z^{(2)}(t_0)$$

where $0 < t_0 < t$ yields the remainder term.

Notes

The central limit theorem (cont.)

Now as $\mathbb{E}[Z_i] = 0$, and $\mathbb{V}[Z_i] = 1$ we can deduce that

$$m_Z(0) = 1 \qquad m_Z^{(1)}(0) = \mathbb{E}[Z_i] = 0$$

and hence

$$m_Z(t) = 1 + \frac{t^2}{2} m_Z^{(2)}(t_0).$$

Notes

The central limit theorem (cont.)

Therefore

$$m_{U_n}(t) = \left\{ 1 + \frac{t^2}{2n} m_Z^{(2)}(t_0) \right\}^n \quad 0 < t_0 < \frac{t}{\sqrt{n}}$$

Now, as $n \rightarrow \infty$

$$\frac{t}{\sqrt{n}} \rightarrow 0$$

so $t_0 \rightarrow 0$.

Notes

The central limit theorem (cont.)

Hence

$$\begin{aligned} \lim_{n \rightarrow \infty} m_{U_n}(t) &= \lim_{n \rightarrow \infty} \left\{ 1 + \frac{t^2}{2n} m_Z^{(2)}(t_0) \right\}^n \\ &= \lim_{n \rightarrow \infty} \left\{ 1 + \frac{t^2}{2n} m_Z^{(2)}(0) \right\}^n \end{aligned}$$

Notes

The central limit theorem (cont.)

But $m_Z^{(2)}(0) = \mathbb{E}[Z_1^2] = 1$, so therefore

$$\lim_{n \rightarrow \infty} m_{U_n}(t) = \lim_{n \rightarrow \infty} \left\{ 1 + \frac{t^2}{2n} \right\}^n = e^{t^2/2}$$

by definition of the exponential function. Hence the limiting mgf is that of a **standard Normal distribution**, and so

$$U_n \xrightarrow{d} Z$$

where Z has a standard Normal distribution.

Notes

Applications of the central limit theorem

We can use the central limit theorem to approximate some standard distributions for which it is otherwise awkward to compute probabilities.

Suppose we wish to approximate the distribution of random variable Y . We must first attempt to decompose Y as

$$Y = \sum_{i=1}^n Y_i$$

where $Y_1, \dots, Y_n$ are independent and identically distributed, and have expectation μ and variance σ^2 that can be easily computed.

Notes

Applications of the central limit theorem (cont.)

We then can say that Y is approximately distributed as

$$Normal(n\mu, n\sigma^2).$$

and compute, for example

$$F_Y(y) = \Pr(Y \leq y) = P\left(\frac{Y - n\mu}{\sqrt{n}\sigma} \leq \frac{y - n\mu}{\sqrt{n}\sigma}\right) \simeq \Phi\left(\frac{y - n\mu}{\sqrt{n}\sigma}\right)$$

where $\Phi(\cdot)$ is the standard Normal cdf.

Notes

Applications of the central limit theorem (cont.)

Example (Binomial case)

If $Y \sim \text{Binomial}(n, p)$, the mgf of Y takes the form

$$m_Y(t) = (pe^t + q)^n$$

By the uniqueness of mgfs, we recognize that Y has the representation

$$Y = \sum_{i=1}^n Y_i$$

where $Y_i \sim \text{Binomial}(1, p) \equiv \text{Bernoulli}(p)$ are independent. We have

$$\mathbb{E}[Y_i] = p \quad \mathbb{V}[Y_i] = pq.$$

Notes

Applications of the central limit theorem (cont.)

Example (Binomial case)

Therefore for large n , Y is approximately distributed as

$$Normal(np, npq)$$


and

$$F_Y(y) = P(Y \leq y) \simeq \Phi\left(\frac{y - np}{\sqrt{npq}}\right)$$

Notes

Applications of the central limit theorem (cont.)


Binomial: $n = 100$, with $p = 0.6$.



Notes

Applications of the central limit theorem (cont.)

Binomial: $n = 20$, with $p = 0.6$.



Notes

Applications of the central limit theorem (cont.)

Example (Poisson case)

If $Y \sim \text{Poisson}(\lambda)$, the mgf of Y takes the form

$$m_Y(t) = \exp\{\lambda(e^t - 1)\}.$$

By the uniqueness of mgfs, we recognize that Y has the representation

$$Y = \sum_{i=1}^n Y_i$$

where the $Y_i \sim \text{Poisson}(\lambda/n)$ are independent. We have

$$\mathbb{E}[Y_i] = \frac{\lambda}{n} \quad \mathbb{V}[Y_i] = \frac{\lambda}{n}.$$

Notes

Applications of the central limit theorem (cont.)

Example (Poisson case)

Therefore Y is approximately distributed as

$$\text{Normal}(\lambda, \lambda)$$

and

$$F_Y(y) = P(Y \leq y) \simeq \Phi\left(\frac{y - \lambda}{\sqrt{\lambda}}\right)$$

Note that for this approximation to work, we need to have λ large.

Notes

Applications of the central limit theorem (cont.)

Continuity correction

When we use the central limit theorem to approximate a discrete distribution, a *continuity correction* is sometimes applied: that is, if Y is discrete we compute

$$F_Y(y) = P(Y \leq y) \simeq \Phi\left(\frac{y + \frac{1}{2} - n\mu}{\sqrt{n}\sigma}\right)$$

Notes

Applications of the central limit theorem (cont.)

Example (Gamma case)

If $Y \sim \text{Gamma}(\alpha, \beta)$, the mgf of Y takes the form

$$m_Y(t) = \left(\frac{1}{1 - \beta t} \right)^\alpha$$

By the uniqueness of mgfs, we recognize that Y has the representation

$$Y = \sum_{i=1}^n Y_i$$

where the $Y_i \sim \text{Gamma}(\alpha/n, \beta)$ are independent. We have

$$\mathbb{E}[Y_i] = \frac{\alpha\beta}{n} \quad \mathbb{V}[Y_i] = \frac{\alpha\beta^2}{n}.$$

Notes

Applications of the central limit theorem (cont.)

Example (Gamma case)

Therefore Y is approximately distributed as

$$\text{Normal}(\alpha\beta, \alpha\beta^2)$$

and


$$F_Y(y) = P(Y \leq y) \simeq \Phi\left(\frac{y - \alpha\beta}{\sqrt{\alpha\beta^2}}\right)$$

Note that for this approximation to work, we need to have α large.

Notes

Applications of the central limit theorem (cont.)


Gamma: $\alpha = 200$, with $\beta = 0.1$.



Notes

Applications of the central limit theorem (cont.)

Gamma: $\alpha = 50$, with $\beta = 0.4$.



Notes

Notes

Notes
