

## MATH 556 - PRACTICE EXAM QUESTIONS

1. The joint pdf for continuous random variables  $X, Y$  with ranges  $\mathbb{X} \equiv \mathbb{Y} \equiv R^+$  is given by

$$f_{X,Y}(x, y) = c_1 \exp \left\{ -\frac{1}{2}(x + y) \right\} \quad x, y > 0$$

and zero otherwise, for some normalizing constant  $c_1$ .

Consider continuous random variable  $U$  defined by

$$U = \frac{1}{2}(X - Y).$$

Find the pdf of  $U$ ,  $f_U$ .

2. In biology, a (2-D) confocal microscopy image of a cell nucleus is well represented by an ellipse with parameters  $a > b$ . Within the cell nucleus are found localized protein bodies (called PMLs), and a key biological question relates to the spatial distribution of the PMLs in the nucleus.

Suppose that the  $(x, y)$  coordinates of a PML body in the image of a nucleus (suitably rotated and standardized for magnitude) are continuous random variables  $X$  and  $Y$  with joint pdf

$$f_{X,Y}(x, y) = c_2 \quad \frac{x^2}{a^2} + \frac{y^2}{b^2} < 1$$

and zero otherwise for some normalizing constant  $c_2$  (that is, the pdf is constant on interior of the ellipse, and zero otherwise).

- Find the marginal pdfs for  $X$  and  $Y$  implied by this joint model.
- Show that the covariance between  $X$  and  $Y$  is zero, and hence that the two variables are uncorrelated.
- Are  $X$  and  $Y$  independent? Justify your answer.

3. (a) Compute, from first principles, the correlation

$$\text{Corr}_{f_{X,Y}} [X, Y]$$

when

$$X \sim \text{Normal}(0, 1)$$

and  $Y = X^2$ .

Are  $X$  and  $Y$  independent? Justify your answer.

*Hint: If  $Y = X^2$ , the rules of expectation dictate that for a general function  $h$*

$$E_{f_{X,Y}} [h(X, Y)] \equiv E_{f_X} [h(X, X^2)]$$

- (b) Suppose that  $X_1$  and  $X_2$  are independent standard normal random variables. Define random variables  $Y_1$  and  $Y_2$  by the multivariate linear transformation

$$Y = AX + b$$

where  $X = (X_1, X_2)^T$  and  $Y = (Y_1, Y_2)^T$  are the column vector random variables,  $A$  is the  $2 \times 2$  matrix

$$A = \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix}$$

and  $b = (1, 2)^T$  is a constant column vector.

- (i) The marginal distribution of  $Y_1$ .  
(ii) The covariance and correlation between  $Y_1$  and  $Y_2$ .

4. (a) Suppose that  $X_1$  and  $X_2$  are independent and identically distributed continuous random variables with cumulative distribution function

$$F_X(x) = \frac{x}{1+x} \quad x > 0$$

with  $F_X(x) = 0$  for  $x \leq 0$ .

Find  $P[X_1 X_2 < 1]$ .

- (b) Suppose that  $Z_1$  and  $Z_2$  are independent  $\text{Normal}(0, 1)$  random variables. Let

$$Y_1 = \frac{Z_1}{\sqrt{Z_1^2 + Z_2^2}} \quad Y_2 = \sqrt{Z_1^2 + Z_2^2}.$$

Find the marginal probability density function of  $Y_1$ .

Are  $Y_1$  and  $Y_2$  independent? Justify your answer.

5. (a) Suppose  $X_1, \dots, X_n, \dots$  are a sequence of random variables with cumulative distribution functions defined by

$$F_{X_n}(x) = \left( \frac{1}{1 + e^{-x}} \right)^n \quad x \in \mathbb{R}.$$

Find the limiting distributions as  $n \rightarrow \infty$  (if they exist) of the random variables

- (i)  $X_n$ ,  
(ii)  $U_n = X_n - \log n$ .

Using the result in (ii), find an approximation to the probability

$$P[X_n > k]$$

for large  $n$ .

- (b) In a dice rolling game, a fair die (with all six scores having equal probability) is rolled repeatedly and independently under identical conditions. On each roll, the player wins six points if the score is a 6, loses one point if the score is either 2,3,4 or 5, and loses two points if the score is 1.

Let  $T_n$  denote the points total obtained after  $n$  rolls of the die. The player begins the game with a points total equal to zero, that is  $T_0 = 0$ .

- (i) Find the expectation and variance of the points total after 100 rolls of the die.  
(ii) Find an approximation to the distribution of the points total after  $n$  rolls, for large  $n$ .  
(iii) Describe the behaviour of the sample average points total,  $M_n = T_n/n$ , as  $n \rightarrow \infty$ .

6. (a) (i) Suppose that random variable  $X$  has a Poisson distribution with parameter  $\lambda$ . Show that standardized random variable,

$$Z_\lambda = \frac{X - \lambda}{\sqrt{\lambda}} \xrightarrow{d} Z \sim N(0, 1)$$

as  $\lambda \rightarrow \infty$

- (ii) Suppose that  $X_1, \dots, X_n \sim \text{Poisson}(\lambda_X)$  and  $Y_1, \dots, Y_n \sim \text{Poisson}(\lambda_Y)$ , with all variables mutually independent. Find  $\mu$  such that the random variable  $M$  defined by

$$M = \bar{X} + \bar{Y}$$

satisfies

$$M \xrightarrow{p} \mu$$

where

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \quad \bar{Y} = \frac{1}{n} \sum_{i=1}^n Y_i$$

are the sample mean random variables for the two samples respectively.

- (b) Suppose that  $X_1, \dots, X_n \sim \text{Exponential}(\lambda)$ . The cdf of the random variable  $T_n = \max\{X_1, \dots, X_n\}$  is given by

$$F_{T_n}(t) = \{F_X(t)\}^n.$$

where  $F_X$  is the cdf of  $X_1, \dots, X_n$ .

- (i) Find  $F_{T_n}(t)$  explicitly.  
(ii) Discuss the form of the limiting distribution of  $T_n$  as  $n \rightarrow \infty$ .  
(iii) Find the form of the limiting distribution of random variable  $U_n$ , defined by

$$U_n = \lambda T_n - \log n$$

as  $n \rightarrow \infty$ .