

# On sequences of spectral triples associated to triangulations

Damien Tageddine

Department of Mathematics and Statistics, McGill University

Canadian Operator Symposium, May 26th, 2023

The principal motivations

- The following diagram in the category of Banach  $*$ -algebras commutes

$$\begin{array}{ccc} A_1 & \xrightarrow{d} & A_2 \\ \downarrow \pi & & \downarrow \pi' \\ A_1^{\hbar} & \xrightarrow{d_{\hbar}} & A_2^{\hbar} \end{array}$$

- We are interested in the question of convergence in norm  $\| \cdot \|_{\hbar}$  when  $\hbar \rightarrow 0$ .
- Discretized operators do not commute in general i.e.  $f(d_{\hbar}g) \neq (d_{\hbar}g)f$ .
- The topology of discrete spaces (lattices, triangulations,...) is ill-behaved.

## Definition (Spectral triple)

A *spectral triple* is the data  $(\mathcal{A}, \mathcal{H}, D)$  where:

- (i)  $\mathcal{A}$  is a real or complex  $*$ -algebra;
- (ii)  $\mathcal{H}$  is a Hilbert space and a left-representation  $(\pi, \mathcal{H})$  of  $A$  in  $B(\mathcal{H})$ ;
- (iii)  $D$  is a *Dirac operator*, which is a self-adjoint operator on  $\mathcal{H}$ .

We require in addition that the Dirac operator satisfies the following conditions

- a) The resolvent  $(D - \lambda)^{-1}$ ,  $\lambda \notin \mathbb{R}$ , is a compact operator on  $H$ .
- b)  $[D, a] \in B(\mathcal{H})$ , for any  $a \in A$ .

# The 2-points space

Let  $a = (a_1, a_2) \in M_2(\mathbb{C})$  and the Dirac operator:

$$D = \frac{i}{\hbar} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad da = \frac{i}{\hbar} \begin{pmatrix} 0 & a_2 - a_1 \\ a_1 - a_2 & 0 \end{pmatrix}.$$

If we define the following distance:

$$d(x, y) = \sup_{a \in A} \{|a(x) - a(y)| : \|[D, a]\| \leq 1\}$$

then one can show that for  $X = \{x, y\}$

$$d(x, y) = \hbar.$$

Without prior assumption, we see the emergence of a small parameter  $\hbar$  in place of the usual distance  $\Delta x$ .

The centre of approximately finite  $C^*$ -algebras exhaust all possible abelian separable  $C^*$ -algebras.

## Theorem (Bratteli)

*Let  $\mathfrak{Z}$  be an abelian separable  $C^*$ -algebra with unit. Then there exists an approximately finite-dimensional  $C^*$ -algebra  $\mathfrak{A}$  having  $\mathfrak{Z}$  as center.*

One can associate a  $C^*$ -algebra  $A$  to a triangulation.

## Theorem (Behncke and Leptin)

*For any (finite) partially ordered set  $X$ , there exists a  $C^*$ -algebra  $A$  such that the primitive spectrum  $\text{Prim}(A)$  is homeomorphic to  $X$ .*

- Associate a separable Hilbert space  $H(X)$  to the space  $X$  and attach to every point  $x \in X$  a subspace  $H(x) \subseteq H(X)$ :

$$H(x) = H^-(x) \otimes H^+(x).$$

- Associate to each point  $x \in X$  an operator algebra  $A(x)$  acting on  $H(x)$ , extended by zero to the whole space  $H(X)$ :

$$A(x) = 1_{H^-(x)} \otimes \mathcal{K}(H^+(x)).$$

- Build the  $C^*$ -algebra  $A(X)$  associated to  $X$ :

$$A(X) = \bigoplus_{x \in X} A(x) \quad \text{acting on} \quad H(X) = \bigoplus_{x \in X} H(x).$$

# Sequences of spectral triples

We can draw the following commuting diagram:

$$\begin{array}{ccccccc} A_1 & \xrightarrow{\phi_{12}^*} & A_2 & \xrightarrow{\phi_{23}^*} & \dots & \xrightarrow{\phi_{i-1i}^*} & A_i & \xrightarrow{\phi_{ii+1}^*} & \dots & \longrightarrow & A_\infty \\ \downarrow id_1 & & \downarrow id_2 & & & & \downarrow id_i & & & & \downarrow \\ X_1 & \xleftarrow{\phi_{12}} & X_2 & \xleftarrow{\phi_{23}} & \dots & \xleftarrow{\phi_{i-1i}} & X_i & \xleftarrow{\phi_{ii+1}^*} & \dots & \xleftarrow{} & X_\infty \end{array}$$

## Theorem

The spectrum  $Spec(A_\infty)$  equipped with the hull-kernel topology is homeomorphic to the space  $X_\infty$  and

$$\lim_{\leftarrow} Spec(A_i) \simeq Spec(\lim_{\rightarrow} A_i).$$

# Sequences of spectral triples

The algebra of continuous functions on the manifold  $M$  can be obtained as the centre of the limit algebra  $A_\infty$ .

## Theorem (T.)

*The limit  $C^*$ -algebra  $A_\infty$  is isometrically  $*$ -isomorphic to  $C^*$ -algebra of the complex valued continuous sections  $\Gamma(M, A_\infty)$  over the manifold  $M$ . The centre  $Z(A_\infty)$  is isomorphic to  $C(M, \mathbb{C})$ .*

A similar result is obtained for the representation space  $L^2(M)$ .

## Theorem (T.)

*The Hilbert space  $L^2(M)$  of square integrable functions over the manifold  $M$  is a subspace of  $H_\infty$ :*

$$H_\infty = L^2(M) \oplus H.$$



## Definition

Let  $D \in M_{2m}(\mathbb{C})$  be an odd and hermitian matrix and let  $\omega_{ij}$  be the coefficients of the block  $D^-$ . We say that  $D$  is an admissible Dirac operator associate to  $X$  if it satisfies the additional condition:

- a) vertices  $i$  and  $j$  do not share an edge  $\Leftrightarrow \omega_{ij} = 0, \forall i, j \in \mathfrak{M}$ ,
- b) the eigenvalues  $\mu_n$  satisfy the asymptotic  $\mu_n(D) = O(\hbar^{-1})$ .

The prototypical example is given by the *combinatorial Dirac operator*, for which:

$$\omega_{ij} := \begin{cases} 1 & \text{if } i \sim j, \\ 0 & \text{otherwise.} \end{cases}$$

# A first example on the lattice

We define the following algebra  $A$  and Dirac operator  $D$ :

$$A = M_{2m}(\mathbb{C}), \quad H = \mathbb{C}^{2m}, \quad D = \frac{i}{\hbar} \begin{pmatrix} 0 & D^- \\ D^+ & 0 \end{pmatrix}$$

with  $(D^+)^* = -D^-$  and where  $D^-$  is given by

$$D^- = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & 0 \\ \vdots & & & \ddots & 1 \\ 0 & \dots & \dots & \dots & 0 \end{pmatrix}.$$

# A first example on the lattice

We consider a sequence of the block matrix block matrices  $D_i$

$$D_i = \frac{i}{\hbar} \begin{pmatrix} 0 & D_i^- \\ D_i^+ & 0 \end{pmatrix}$$

Then the limit operator  $D_\infty$  acts on  $A_\infty$  by the commutator:

$$[D_\infty, a] = ([D_0, a_0], [D_1, a_1], \dots, [D_i, a_i], \dots) \in \prod_{i \in I} M_{2m_i}^-(\mathbb{C}).$$

We can compute the spectrum of the commutator  $[D_\infty, a]$ :

$$\text{i) } \sigma_{A_\infty}([D_\infty, a]) = \overline{\cup_i \sigma_{A_i}([D_i, a_i])}$$

$$\text{ii) } \|[D_\infty, a]\| = \|d_c a\|_\infty$$

# A first example on the lattice

## Proposition (Spectral convergence)

There exists a finite measure  $\mu$  and a unitary operator

$$U : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}, d\mu) \quad (1)$$

such that,

$$U[D, a]U^{-1}\phi = \frac{da}{dx}\phi, \quad \forall \phi \in L^2(\mathbb{R}), \quad (2)$$

Moreover, the norm of  $[D, a]$  is given by  $\|[D, a]\| = \|d_c a\|_\infty$ .

This result can be generalized to the  $d$ -dimensional lattice  $\Lambda$ . The  $C^*$ -algebra  $A(\Lambda)$  and the Dirac operator  $D$  are obtained through tensor products:

$$A(\Lambda) = A(L) \otimes \cdots \otimes A(L), \quad D_n = \sum_{k=1}^d 1 \otimes \cdots \otimes D_n^{(k)} \otimes \cdots \otimes 1.$$

- It is known that the canonical spectral triple  $(C^\infty(M), L^2(S), D)$  on a spin manifold  $M$  encodes the metric. The geodesic distance between any two points  $p$  and  $q$  on  $M$  is given by

$$\inf_{\gamma} \int_0^1 \sqrt{g_{\gamma}(\dot{\gamma}(t), \dot{\gamma}(t))} dt = \sup_{f \in \mathcal{A}} \{|f(p) - f(q)| : \|[D, f]\| \leq 1\}$$

- As it defined the combinatorial Dirac operator does not depend on the metric  $g$  of the manifold  $M$ .
- Beyond the case of the lattice, the eigenvalues of the commutator  $[D, a]$  are not immediately accessible.

If we consider the more general definition of  $D$  given by

$$(D)_{ij} := \begin{cases} \omega_{ij} \neq 0 & \text{if } i \sim j, \\ 0 & \text{otherwise.} \end{cases}$$

where the coefficients  $\omega_{ij}$  are obtained from a density distribution, a first approach would be to study the convergence in average:

$$S_n^{\hbar_n}(a) := \frac{1}{n} \sum_{k=1}^n e_k \left[ D_X^k, a_k \right] e_k^*$$

with  $(e_k)$  a family of projectors.

# F-P Equation and the Von-Mises Fisher distribution

Consider the one-parameter family of measures  $(\mu_{x,t})_t$  satisfy the parabolic equation:

$$\left. \frac{\partial \mu_{x,t}}{\partial t} \right|_{t=0} = L_{A,b}(\mu_{x,t}) \quad (3)$$

in the weak sense, with the operator  $L_{A,b}$

$$L_{A,b}f = \text{tr}(AD^2f) + \langle b, \nabla f \rangle, \quad f \in C_c^\infty(M) \quad (4)$$

We consider the von Mises-Fisher distribution on the unit sphere  $\mathbb{S}^d$  given by:

$$\rho_d(x; s, \beta) = C_d(\beta) \exp(-\beta \langle s, x \rangle) \quad (5)$$

where  $\beta \geq 0$ ,  $\|s\| = 1$  and  $C_d(\beta)$  is a normalization constant.

# The Von-Mises Fisher distribution

We show that the von Mises-Fisher distribution satisfies the Fokker-Planck equation:

$$\left. \frac{\partial \rho_{s,t}}{\partial t} \right|_{t=0} = \partial_s(\rho_{s,t}).$$

The distribution can be defined on a normal neighbourhood  $U_p$  of the manifold  $M$  and satisfies a Fokker-Planck equation.

## Proposition

*The following limit holds at a point  $p \in M$*

$$\left. \frac{\partial}{\partial t} \left( C_d(\beta_t) \int_{U_p} e^{\hat{\Phi}_{\beta}(s_i, x)} f(x) \mu(x) \right) \right|_{t=0} = \partial_i(f)(p).$$



# A first convergence result

We defined the family of projectors  $e_k$  such that:

$$e_k D_{X_k} e_k^* = \left( \begin{array}{c|ccccc} & & & & & \\ & 0 & & & & \\ & & * & * & \omega_{i_0 j}^k & * & * \\ \hline & * & & & & & \\ & * & & & & & \\ & \omega_{i_0 j}^k & & & & & \\ & * & & & & & \\ & * & & & & & \\ & & & & & 0 & \\ & & & & & & \end{array} \right)$$

and the coefficients  $\omega_{ij}$  are defined  $\omega_{ij}^k(\hbar) = C_d(\beta_{\hbar}) \exp\left(-\frac{\langle x_i^k, s_j \rangle}{\hbar}\right)$ .

# A first convergence result

## Theorem (T.)

Let  $\{x_{i_0}^k\}_{k=1}^n$  be a sequence of i.i.d. sampled points from a uniform distribution on an open normal neighbourhood  $U_p$  of a point  $p$  in a compact Riemannian manifold  $M$  of dimension  $d$ . Let  $\widehat{S}_n^{\hbar_n}$  be the associated operator given by:

$$\widehat{S}_n^{\hbar_n}(a) := \frac{1}{n} \sum_{k=1}^n e_k \left[ D_X^k, a_k \right] e_k^*.$$

Put  $\hbar_n = n^{-\alpha}$ , where  $\alpha > 0$ , then in probability:

$$\lim_{n \rightarrow \infty} \sup_{a \in F} \left| \Psi \circ \widehat{S}_n^{\hbar_n}(a)(p) - [\mathcal{D}, a](p) \right| = 0.$$

## Theorem (T.)

Let  $\{x_i\}_{i=1}^n$  be a sequence of i.i.d. sampled points from a uniform distribution on an open normal neighbourhood  $U_p$  of a point  $p$  in a compact Riemannian manifold  $M$  of dimension  $d$ .  $\Omega_n^{\hbar_n}$  be the associated operator given by:

$$\Omega_n^{\hbar_n}(a)(p) = \frac{C_d(\beta_{\hbar})}{n\hbar^2} \sum_{k=1}^n \sum_{j=1}^{d+1} \lambda_j^2 \exp\left(-\frac{\langle x_i^k, s_j \rangle}{\hbar}\right) \alpha_{ij}(a_k).$$

Put  $\hbar_n = n^{-\alpha}$ , where  $\alpha > 0$ , then in probability:




$$\lim_{n \rightarrow \infty} \sup_{a \in F} \left| \Omega_n^{\hbar_n}(a)(p) - \Delta_M(a)(p) \right| = 0 \quad (6)$$





Given a compact spin manifold  $(M, g)$ , we have the following:

- associate to each  $K_i$  a  $C^*$ -algebra  $A_i$  with limit  $C(M)$ ,
- define a differential structure  $da = [D_i, a]$  on each  $A_i$ ,
- for the lattice,  $(D_i)$  converges to the usual Dirac operator  $\partial_M$ .
- Using the same tools than the continuous case  $(C^\infty, L^2(M), \partial_M)$ .

Future works:

- convergence results of the  $(D_i)$  to the classical Dirac operator,
- provide a unifying framework in the language of spectral triples.

-  **D. Tageddine, J-C. Nave**  
Noncommutative geometry on Infinitesimal Spaces  
submitted, [arXiv:2209.12929](#).
-  **D. Tageddine, J-C. Nave**  
Statistical Fluctuation of Infinitesimal Spaces  
submitted, [arXiv:2304.10617](#).
-  **Balachandran, A. P. and Bimonte, G. and Ercolessi, E. and Landi, G. and Lizzi, F. and Sparano, G. and Teotonio-Sobrinho, P.**  
Noncommutative Lattices as Finite Approximations and Their Noncommutative Geometries  
[Journal of Geometry and Physics \(1996\)](#), pp. 163-194.

-  **M. Khalkhali and N. Pagliaroli**  
Spectral Statistics of Dirac Ensembles  
*Journal of Mathematical Physics* (2022).
-  **A. Connes**  
Noncommutative geometry  
*Academic Press* (1994).
-  **O. Bratteli**  
The center of approximately finite-dimensional  $C^*$ -algebras  
*Journal of Functional Analysis* (1975), pp. 195-202.
-  **H. Behncke and H. Leptin**  
 $C^*$ -algebras with finite duals  
*Journal of Functional Analysis* (1973), pp. 253-262