

## MATH 599 PROBLEM SET 4

DUE THURSDAY MARCH 28

In this problem set, we will study when geodesics maximize the proper time among timelike curves connecting a hypersurface and a point. Let  $M$  be a Lorentzian manifold (of dimension  $n$ ), and let  $\phi : H \rightarrow M$  be a spacelike hypersurface (of dimension  $n - 1$ ) embedded into  $M$ . We will identify  $\phi(H) \subset M$  with  $H$ . Then  $\mathfrak{X}(\phi)^\top$  can be identified with  $\mathfrak{X}(H)$ , and we have the (pointwise) orthogonal decomposition

$$\mathfrak{X}(\phi) = \mathfrak{X}(\phi)^\top \oplus \mathfrak{X}(\phi)^\perp = \mathfrak{X}(H) \oplus \mathfrak{X}(\phi)^\perp,$$

where  $\mathfrak{X}(\phi)^\perp$  is the space of vector fields along  $\phi$  that are pointwise orthogonal to  $H$ . We denote by  $P^\perp : \mathfrak{X}(\phi) \rightarrow \mathfrak{X}(\phi)^\perp$  the orthogonal projection onto  $\mathfrak{X}(\phi)^\perp$ . Thus, if  $N \in T_p M$  is a nonzero vector at  $p \in H$  satisfying  $N \perp H$ , then

$$(P^\perp X)_p = \frac{\langle X, N \rangle_p}{\langle N, N \rangle_p} N \quad \text{for } X \in \mathfrak{X}(\phi).$$

Suppose that  $q \in M \setminus H$ , and consider the set  $C(H, q)$  of all timelike curves joining  $H$  and  $q$ . Given a curve  $\gamma \in C(H, q)$ , a smooth map  $\omega : (-\varepsilon, \varepsilon) \times [a, b] \rightarrow M$  is called a *deformation of  $\gamma$* , if  $\omega(0, \cdot) = \gamma$ ,  $\omega(\cdot, a) \in H$ ,  $\omega(\cdot, b) = q$ , and each “longitudinal” curve  $\gamma_s = \omega(s, \cdot)$  is timelike. We set  $X = \omega_* \partial_s \in \mathfrak{X}(\omega)$  and  $T = \omega_* \partial_t \in \mathfrak{X}(\omega)$ , where  $s$  and  $t$  are the Cartesian coordinates in the rectangle  $(-\varepsilon, \varepsilon) \times [a, b]$ . Note that  $X(s, b) = 0$  and  $X(s, a) \in T_{\omega(s, a)} H$  for all  $s$ . We will also use  $X$  and  $T$  to denote the *variation field*  $X|_{s=0} \in \mathfrak{X}(\gamma)$  and the *velocity field*  $T|_{s=0} \in \mathfrak{X}(\gamma)$ , respectively. Recall that the proper time of  $\gamma_s$  is given by

$$\tau(s) = \tau(\gamma_s) = \int_a^b |T| dt = \int_a^b \sqrt{-\langle T, T \rangle} dt.$$

- 1) Compute the first variation  $\tau'(s)$ , and show that it depends only on the variation field  $X$  and the velocity field  $T$  along  $\gamma$ . Assuming that  $\gamma$  is parameterized by proper time as  $|T| = 1$ , show that  $\tau'(0) = 0$  for all variation fields  $X$  with  $X(b) = 0$  if and only if  $\gamma$  is a geodesic and  $T(a) \perp H$ .
- 2) We define the *second fundamental form*  $\mathbb{II} : \mathfrak{X}(H) \times \mathfrak{X}(H) \rightarrow \mathfrak{X}(\phi)^\perp$  of  $H$  by

$$\mathbb{II}(X, Y) = P^\perp \nabla_X Y.$$

It is clear that  $\mathbb{II}$  is bilinear. Prove the following, and conclude in particular that the second fundamental form is tensorial in each of its arguments.

- (a)  $\langle \mathbb{II}(X, Y), N \rangle = -\langle \nabla_X N, Y \rangle$  for  $N \in \mathfrak{X}(\phi)^\perp$ .
- (b)  $\mathbb{II}(X, Y) = \mathbb{II}(Y, X)$ .
- 3) Let  $\gamma : [a, b] \rightarrow M$  be a timelike geodesic with  $\gamma(a) = p \in H$ ,  $\gamma(b) = q \in M \setminus H$ , and  $T(a) \perp H$ . Such a geodesic will be called a timelike geodesic *normal to  $H$* . For any deformation of  $\gamma$ , show that

$$\tau''(0) = I(X, X) - \langle \nabla_T X, X \rangle_p,$$

where  $I$  is the index form as defined in class. Show also that the following hold.

- (a)  $X(b) = 0$ .
- (b)  $X(a) \perp T(a)$ , that is,  $X(a) \in T_p H$ .
- (c)  $\langle \nabla_T X, V \rangle_p = -\langle \mathbb{I}(X, V), T \rangle_p$  for all  $V \in T_p H$ .

We call any  $X \in \mathfrak{X}(\gamma)$  satisfying these 3 conditions an  $H$ -proper variation. Moreover, we say that  $q$  is *conjugate to  $H$  along  $\gamma$*  if there exists a nontrivial Jacobi field along  $\gamma$  that is also an  $H$ -proper variation.

- 4) Consider a timelike geodesic  $\gamma$  normal to  $H$ . Let

$$I_H(X, Y) = I(X, Y) - \langle \nabla_T X, Y \rangle_p \quad \text{for } X, Y \in \mathfrak{X}(\gamma),$$

and prove the following.

- (a) For  $H$ -proper variations  $X$  and  $Y$ , we have

$$I_H(X^\perp, Y^\perp) = I_H(X, Y) = I(X, Y) + \langle \mathbb{I}(X, Y), T \rangle_p,$$

where  $X \mapsto X^\perp$  is the orthogonal projection onto  $\mathfrak{X}(\gamma)^\perp$ .

- (b) If  $H$  has no conjugate point along  $\gamma$  on  $(a, b]$ , then  $I_H(X, X) < 0$  for any nonzero  $H$ -proper variation  $X$  in  $\mathfrak{X}(\gamma)^\perp$ .
- 5) In the same setting, prove that if  $H$  has a conjugate point along  $\gamma$  at the parameter value  $c \in (a, b)$ , there exists an  $H$ -proper variation  $X$  in  $\mathfrak{X}(\gamma)^\perp$  such that  $I_H(X, X) > 0$ .
- 6) Show that if  $\gamma$  is a timelike geodesic normal to  $H$ , and if  $X$  is an  $H$ -proper variation, then there is a deformation of  $\gamma$  with its variation field equal to  $X$ .