

The Real Jordan Canonical Form and the Rational Canonical Form

Not all matrices over a given field have a Jordan canonical form as not all polynomials split completely into linear factors. For example, over the reals one can have irreducible quadratic factors. A field in which every polynomial splits completely into a product of linear factors over that field is said to be algebraically closed. An example, of such a field is the field of complex numbers.

Let A be an $n \times n$ real matrix and let $T = T_A : \mathbb{R}^{n \times 1} \rightarrow \mathbb{R}^{n \times 1}$ be the associated linear operator. The minimal polynomial $m_A(X)$ of A has the form

$$(X - c_1)^{m_1} \cdots (X - c_k)^{m_k} ((X - a_1)^2 + b_1^2)^{n_1} \cdots ((X - a_\ell)^2 + b_\ell^2)^{n_\ell}$$

which has the factorization

$$(X - c_1)^{m_1} \cdots (X - c_k)^{m_k} (X - \alpha_1)^{n_1} (X - \bar{\alpha}_1)^{n_1} \cdots (X - \alpha_\ell)^{n_\ell} (X - \bar{\alpha}_\ell)^{n_\ell}$$

over \mathbb{C} with $\alpha_j = a_j + b_j i$, $\bar{\alpha}_j = a_j - b_j i$. Since the nullity of a real matrix is the same as the nullity of the matrix viewed as a complex matrix, the number of Jordan blocks $J_r(\lambda)$ with λ real can be computed as before. The complex Jordan blocks come in conjugate pairs $J = J_r(a + ib)$, $\bar{J} = J_r(a - ib)$ and their number of such pairs is $s_r/2$ where

$$s = (s_0, s_1, \dots, s_j, \dots) = -R(L - 1)^2(d),$$

where $d = (d_0, d_1, \dots, d_j, \dots)$ with $d_j = \text{nullity}((A - a)^2 + b^2)^j$. If $u \in \mathbb{C}^{n \times 1}$ is the cyclic vector associated to $J_r(a + bi)$ then \bar{u} is a cyclic vector for $J_r(a - ib)$. If

$$g = \frac{u - \bar{u}}{2i}, \quad h = \frac{u + \bar{u}}{2},$$

then g, h are real vectors such that, over \mathbb{C} ,

$$W = \text{Span}(g, h, T(g), T(h), \dots, T^{r-1}(g), T^{r-1}(h)) = \text{Span}(u, \bar{u}, T(u), T(\bar{u}), \dots, T^{r-1}(u), T^{r-1}(\bar{u}))$$

If $u_1 = (T - \alpha)^{r-1}(u), \dots, u_{r-1} = (T - \alpha)(u), u_r = u$ and $\alpha = a + bi$ then

$$T(u_1) = \alpha u_1, \quad T(u_j) = \alpha u_j + u_{j-1} \text{ for } 2 \leq j \leq r.$$

Setting $f_{2j-1} = \frac{u_j - \bar{u}_j}{2i}$, $f_{2j} = \frac{u_j + \bar{u}_j}{2}$ for $1 \leq j \leq r$, we get

$$T(f_{2j-1}) = a f_{2j-1} + b f_{2j} + f_{2j-3}, \quad T(f_{2j}) = -b f_{2j-1} + a f_{2j} + f_{2j-2}$$

for $2 \leq j \leq r$ and $T(f_1) = a f_1 + b f_2$, $T(f_2) = -b f_1 + a f_2$. The real vectors f_1, \dots, f_{2r} are linearly independent and span W . The matrix of the restriction of T to the subspace of $\mathbb{R}^{n \times 1}$ spanned by f_1, \dots, f_{2r} with respect to $f = (f_1, \dots, f_{2r})$ is the $2r \times 2r$ block matrix $RJ_r(a, b) = [B_{ij}]$ with

$$B_{ii} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}, \quad B_{i(i+1)} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

and B_{ij} the zero 2×2 matrix for all other i, j . We thus obtain the following theorem:

Theorem. If T is a linear operator on a finite-dimensional real vector space then there is a basis for V such that the matrix of T with respect to this basis has a block diagonal form with blocks of the form $J_k(a)$ or $RJ_k(a, b)$. Moreover, if s_k is the number of blocks of the form $J_k(a)$, then

$$s = (s_0, s_1, \dots, s_i, \dots) = -R(L-1)^2(d)$$

where $d = (d_0, d_1, \dots, d_i, \dots)$ with $d_i = \dim \text{Ker}((T - a)^i)$. If s_k is the number of blocks of the form $RJ_k(a, b)$, then

$$s = (s_0, s_1, \dots, s_i, \dots) = -R(L-1)^2(d)$$

where $d = (d_0, d_1, \dots, d_i, \dots)$ with $2d_i = \dim \text{Ker}(((T - a)^2 + b^2)^i)$.

Corollary 1. If A, B are real matrices that are similar over \mathbb{C} then they are similar over \mathbb{R} .

Corollary 2. Every real or complex square matrix is similar to its transpose.

Proof. Because of Corollary 1, we can assume that the matrix A is complex. We can also assume that A is in Jordan canonical form in which case we are reduced to proving the corollary in the case A is a Jordan matrix. This is left as an exercise for the reader.

If $f(X)$ is a polynomial over a field F , we can always construct a field K containing F as a subfield in which $f(X)$ splits completely into linear factors. One can even construct such a field which is algebraically closed. as a result, we obtain the Cayley-Hamilton Theorem. Namely, if $\Delta_T(X)$ is the characteristic polynomial of T , then $\Delta_T(T) = 0$.

There is a canonical form for the matrix of a linear operator T on a finite-dimensional vector space V over F , called the rational canonical form. The vector space V is a direct sum of cyclic subspaces, ie, subspaces having a basis of the form

$$v, T(v), \dots, T^{p-1}(v)$$

with $T^p(v) = c_1 T^{p-1}(v) + \dots + c_{p-1} T(v) + c_p v$. Such a vector v is called a cyclic vector for T . This subspace, called the cyclic subspace for T generated by v , is T -invariant and the matrix of the restriction of T to this subspace is the $p \times p$ matrix

$$\begin{bmatrix} 0 & 0 & 0 & \cdots & c_p \\ 1 & 0 & 0 & \cdots & c_{p-1} \\ 0 & 1 & 0 & \cdots & c_{p-2} \\ 0 & 0 & 1 & \cdots & c_{p-3} \\ \vdots & & & & \\ 0 & 0 & 0 & \cdots & c_1 \end{bmatrix}.$$

This matrix is called the companion matrix of the cyclic vector v . Since the polynomial $f(X)$ is uniquely determined by v , this matrix is also called the companion matrix of $f(X)$ and is denoted by C_f . The polynomial $f(X)$ is the minimal polynomial of C_f and therefore of the restriction of T to the cyclic subspace generated by v .

If the minimal polynomial $m_T(X)$ of T has the primary decomposition

$$m_T(X) = p_1(X)^{m_1} \cdots p_k(X)^{m_k}$$

with $p_i(X)$ irreducible then V is the direct sum of cyclic subspaces with companion matrices C_f where $f(X) = p_i(X)^j$, $1 \leq i \leq k$, $1 \leq j \leq m_i$. Moreover, if s_j is the number of summands with $f(X) = p_i(X)^j$, then

$$s = (s_0, s_1, \dots, s_j, \dots) = -R(L-1)^2(d),$$

where $d = (d_0, d_1, \dots, d_j, \dots)$ with $d_j = \dim \text{Ker}(p_i(T)^j) / \text{degree}(p_i(X))$. The proof of this result is exactly the same as the proof of the Jordan canonical form except that $p_i(T)$ replaces $T - a_i$ and the vector spaces $W_{ij} = \text{Ker}(p_i(T)^j) / \text{Ker}(p_i(T)^{j-1})$ are viewed as vector spaces over the field $K_i = F[X]/(p_i(X))$, where

$$\overline{f(X)} \cdot \overline{g(T)} = \overline{f(T)g(T)}.$$

This field is a vector space over F of dimension $\ell_i = \text{degree}(p_i(X))$ with basis $1, \overline{X}, \dots, \overline{X}^{\ell_i-1}$. Thus

$$\dim_F(W_{ij}) = \ell_i \dim_{K_i}(W_{ij}).$$

This explains the formula for d_j . The cyclic vectors giving the direct sum decomposition are representatives of basis vectors for the quotient spaces $W_{ij}/\text{Im}(S_{ij})$ where S_{ij} is the linear mapping from $W_{i(j+1)}$ to W_{ij} defined by

$$S_{ij}(\overline{u}) = \overline{p_i(T)(u)}.$$

The details are left to the reader.

It follows that two $n \times n$ matrices over F which are similar over a field containing F as a subfield are similar over F . This implies, for example, that a square matrix is similar to its transpose.

Example. Let us find the rational canonical form of the matrix

$$A = \begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & -1 & 1 & 1 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

over the field \mathbb{Q} of rational numbers. This matrix can be viewed as an upper triangular 2×2 block matrix with blocks that are 2×2 matrices. Since the diagonal blocks have minimum polynomial $X^2 - 2$, it follows that $(A^2 - 2)^2 = 0$. We leave it to the reader to formulate a more general result. We have

$$A^2 - 2 = \begin{bmatrix} 2 & 0 & 2 & 3 \\ 0 & 2 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad (A^2 - 2)^2 = 0$$

so that $\text{Null}(A^2 - 2) = \text{Span}(e_1, e_2)$, $\text{Null}((A - 2)^2) = (e_1, e_2, e_3, e_4)$. Hence

$$\text{Null}((A^2 - 2)^2) / \text{Null}(A^2 - 2) = \text{Span}(\overline{e}_3, \overline{e}_4) = \text{Span}_K(\overline{e}_3)$$

where $K = \mathbb{Q}[X]/(X^2 - 2) \cong \mathbb{Q}[\sqrt{2}]$. Note that $\overline{X}\overline{e}_3 = \overline{Ae}_3 = 2\overline{e}_4$. We also have

$$\text{Null}(A^2 - 2) = \text{Span}(e_1, e_2) = \text{Span}_K(e_1)$$

since $\overline{X}e_1 = Ae_1 = 2e_2$. Since $d = (0, 1, 2, 2, \dots)$, we have $s = -R(L-1)^2(d) = (0, 0, 1, 0, 0, \dots)$ so that we have only one cyclic subspace in the direct sum decomposition of \mathbb{Q}^6 with cyclic vector

e_3 . Since $(X^2 - 2)^2 = X^4 - 4x^2 + 4$ is the companion polynomial of this cyclic vector, there is an invertible matrix over \mathbb{Q} such that

$$P^{-1}AP = \begin{bmatrix} 0 & 0 & 0 & -4 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 4 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

Moreover, the columns of P are the vectors $e_3, Ae_3, A^2e_3, A^3e_3$ so that

$$P = \begin{bmatrix} 0 & 1 & 2 & 4 \\ 0 & 1 & 0 & 4 \\ 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 2 \end{bmatrix}.$$

The minimum of A is $(X^2 - 2)^2$. This is also the characteristic polynomial of A . We leave it to the reader to show that the Jordan canonical form of A over the field $\mathbb{Q}[\sqrt{2}]$ is

$$\begin{bmatrix} \sqrt{2} & 1 & 0 & 0 \\ 0 & \sqrt{2} & 0 & 0 \\ 0 & 0 & -\sqrt{2} & 1 \\ 0 & 0 & 0 & -\sqrt{2} \end{bmatrix}.$$