

MATH 255: Assignment 2

(due Friday, January 24)

1. Let $(a_n)_{n \geq 1}$ be a sequence of real numbers. For $x \geq 0$, define

$$A(x) = \sum_{n \leq x} a_n = \sum_{n=1}^{[x]} a_n,$$

where $[x]$ is the largest integer $\leq x$ and empty sums are equal to zero. If f has a continuous derivative on $[1, a]$, show that

$$\sum_{n \leq a} a_n f(n) = - \int_1^a A(x) f'(x) dx + A(a) f(a)$$

using Stieltjes integrals.

2. Show that the following integrals exist and compute their value:

(a) $\int_0^5 x^2 d[x];$

(b) $\int_{-2}^3 [|x|] d|x|;$

(c) $\int_0^6 (x^2 + [x]) d|3 - x|;$

(d) $\int_0^2 x d([x] - x).$

3. Using integration by parts, derive the following identities:

(a) $\sum_{k=1}^n \frac{1}{n^s} = \frac{1}{n^{s-1}} + s \int_1^n \frac{[x]}{x^{s+1}} dx$ if $s \neq 1$.

(b) $\sum_{k=1}^n \frac{1}{k} = \log n - \int_1^n \frac{x - [x]}{x^2} dx + 1.$

(Last modified 8:30 pm January 20, 2003)