

MATH 255: Assignment 3

(due Monday, February 3)

1. If  $f \in \mathcal{R}(a, b)$  and  $g$  is a function on  $[a, b]$  such that  $g(x) = f(x)$ , except possibly for a finite set of points, show that  $g \in \mathcal{R}(a, b)$  and  $\int_a^b g(x) dx = \int_a^b f(x) dx$ .
2. If  $f \in \mathcal{R}$  on  $[a, b]$ , prove that

$$\lim_{n \rightarrow \infty} \frac{b-a}{n} \sum_{k=1}^n f\left(a + k \frac{b-a}{n}\right) = \int_a^b f(x) dx.$$

Deduce that

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{n}{n^2 + k^2} = \frac{\pi}{4}, \quad \lim_{n \rightarrow \infty} \sum_{k=1}^n (n^2 + k^2)^{-1/2} = \log(1 + \sqrt{2}).$$

3. (a) If  $a < b$ , give an example of a function  $f$  on  $[a, b]$  such that  $|f| \in \mathcal{R}(a, b)$  but  $f \notin \mathcal{R}(a, b)$ .  
(b) Give an example of a function  $f$  on  $[0, 1]$  such that  $f'$  exists but  $f' \notin \mathcal{R}(0, 1)$ .

(Last modified 7:45 am, January 29, 2003)