

McGill University
Math 270A: Applied Linear Algebra
Assignment 3: due Tuesday October 26 , 1999

1. If V is the real vector space \mathbb{R}^X (X any set), show that its complexification $V_{\mathbb{C}}$ is isomorphic to \mathbb{C}^X . **Hint:** Show that any $f \in \mathbb{C}^X$ can be written in the form $f_1 + if_2$ with $f_1, f_2 \in \mathbb{R}^X$.
2. (a) Let S be a subset of \mathbb{R}^2 and let W_S be the subset of \mathbb{R}^6 consisting of those 6-tuples (a, b, c, d, e, f) such that $ax^2 + bxy + cy^2 + dx + ey + f = 0$ for all (x, y) in S . Show that W_S is a subspace of \mathbb{R}^6 ;
(b) If $S = \{(1, 1), (2, 5), (3, 0), (4, 6)\}$, show that $\dim W_S = 2$;
(c) Find a basis for W_S by taking products of suitable pairs of equations of lines passing through points of S ;
(d) Use the above to find the equation of the conic $ax^2 + bxy + cy^2 + dx + ey + f = 0$ which passes through the points of S and the point $(-1, 1)$.
3. Let V be the real inner product space of real-valued functions on the interval $[0, 1]$ with the inner product

$$\langle f, g \rangle = \int_0^1 f(x)g(x)dx,$$

let W be the subspace of polynomial functions of the form $f(x) = a + bx + cx^2$ with $a, b, c \in \mathbb{R}$ and let $T : \mathbb{R}^3 \rightarrow W$ is the function defined by $T((a, b, c)) = f$, where $f(x) = a + bx + cx^2$.

- (a) Show that T is an isomorphism of vector spaces.
 - (b) Show that $\langle u, v \rangle = \langle T(u), T(v) \rangle$ is an inner product on \mathbb{R}^3 and find a formula for $\|(a, b, c)\|$ using this inner product.
4. Let W be the subspace of \mathbb{R}^4 spanned by the vectors

$$u_1 = (1, 2, 2, 1), \quad u_2 = (1, 1, -1, -1), \quad u_3 = (2, -1, -1, 2).$$

Find the orthogonal projection of the vector $v = (a, b, c, d)$ on W and use this to find the least squares solution of the system

$$\begin{aligned}x + y + 2z &= a \\2x + y - z &= b \\2x - y - z &= c \\x - y + 2z &= d.\end{aligned}$$

5. Using the notation of question 3, find the orthogonal projection of the function $f(x) = e^x$ on W . In what sense is this orthogonal projection the best approximation to $f(x) = e^x$ by a polynomial function of degree ≤ 3 ?