McGill University Math 270A: Applied Linear Algebra Solution Sheet for Test 1

- 1. (a) The zero vector of V is the zero function 0. Since 0(0) = 0(1) = 0, we have  $0 \in W$ . If  $f, g \in W$  and  $a, b \in \mathbb{R}$ , we have (af + bg)(0) = af(0) + bg(0) = 0, (af + bg)(1) = af(1) + bg(1) = 0 which implies  $af + bg \in W$ . Hence W is a subspace of V.
  - (b) If f(x) = x, g(x) = 1 x then f(0) = 0, g(1) = 0 which imply  $f, g \in W$ . But (f + g)(x) = 1 which implies that  $f + g \notin W$ .
- 2. (a) We have  $x_1A_1 + x_2A_2 + x_3A_3 + x_4A_4 = 0$  if and only if
  - $x_1 + 2x_2 + x_4 = 0$   $x_1 + x_2 + x_3 + x_4 = 0$   $-x_1 + x_2 - 3x_3 - 2x_4 = 0$  $x_1 - 2x_2 + 4x_4 + 3x_3 = 0$

which has as solutions  $(x_1, x_2, x_3, x_4) = a(2, -1, -1, 0)$ . Thus, up to a scalar multiple, the only dependence relation is  $2A_1 - A_2 - A_3 = 0$ . From this is follows that  $A_3 \in \text{span}(A_1, A_2)$  and  $A_1, A_2, A_4$  are linearly independent and span W. Thus  $\dim(W) = 3 < \dim(\mathbb{R}^{2\times 2} = 4)$  which implies  $W \neq \mathbb{R}^{2\times 2}$ .

- (b) In (a) we found that  $A_1, A_2, A_4$  was a basis of W.
- 3. (a) Since elmentary row operations don't change the row space of a matrix, and since the non-zero rows of a matrix in echelon form are a basis for the row space of this matrix, we see that [1, 0, 2, 0, 1], [0, 1, 1, 1] is a basis for the row space of A. Moreover, since the pivot columns of A are a basis for the column space of A we see that  $[7, 8, 6, 2]^T, [8, 7, 9, -2]^T$  is a basis for the column space of A. Since the null space of  $A = \{X \mid AX = 0\}$  and  $AX = 0 \iff X = a[2, -1, 1, 0, 0]^T + b[0, -1, 0, 1, 0]^T + c[-1, -1, 0, 0, 1]^T$ , the vectors  $[-2, -1, 1, 0, 0]^T, [0, -1, 0, 1, 0]^T, [-1, -1, 0, 0, 1]^T$  span the null space and are a basis since they are linearly independent.
  - (b) The rank of a matrix is the dimension of the row space and the nullity of a matrix is the dimension of its null space. Hence, by (a), we have  $\operatorname{rank}(A) = \operatorname{rank}(A^T) = 2$ . Since  $\operatorname{rank} + \operatorname{nullity} = \#$  of cols, we have  $\operatorname{nullity}(A) = 5 - 2 = 3$ ,  $\operatorname{nullity}(A^T) = 4 - 2 = 2$ .
- 4. (a) We have  $u, v \in W$  since  $5u_{n+1} 6u_n = 5 \cdot 2^{n+1} 6 \cdot 2^n = (10-6)2^n = 2^{n+2} = u_{n+2}$  and  $5v_{n+1} 6v_n = 5 \cdot 3^{n+1} 6 \cdot 3^n = (15-6)3^n = 3^{n+2} = v_{n+2}$ . Now au + bv = 0 implies a + b = 0, 2a + 3b = 0 which implies a = b = 0 and hence that u, v are linearly independent. Since dim(W) = 2 the vectors also span W since any n linearly independent vectors in an n-dimensional vector space V are a basis for V.
  - (b) If  $w \in W$  we have w = au + bv by (a). But then a + b = 1, 2a + 3b = 1 which has the only solution a = 2, b = -1. Hence the coordinate vector of w with respect to the basis u, v is  $[2, -1]^T$ .