

McGill University  
Math 371B: Algebra IV  
Assignment 2: due Monday, February 1, 1999

A ring  $A$  is called a valuation ring if  $A$  is an integral domain such that for any non-zero element  $x$  of its quotient field we have  $x \in A$  or  $1/x \in A$ . A valuation ring which is a field is called a trivial valuation ring. A discrete valuation ring is an example of a non-trivial valuation ring. Any subring  $A$  of a field  $K$  such that for any  $x \in K^* = K - \{0\}$  we have  $x \in A$  or  $1/x \in A$  is a valuation ring with quotient field  $K$ . Such a subring of  $K$  is called a valuation ring of  $K$ .

1. Show that a valuation ring is a local ring. If  $F$  is a field, show that the localization of the polynomial ring  $F[x, y]$  at the maximal ideal  $(x, y)$  is not a valuation ring.
2. Show that a ring is a discrete valuation ring if and only if it is a PID with a single non-zero maximal ideal. Show also that a discrete valuation ring is a maximal element in set of non-trivial valuation rings of its quotient field.
3. Show that the non-trivial valuation rings of  $\mathbb{Q}$  are the discrete valuation rings  $\mathbb{Z}_{(p)}$  where  $p$  is a prime of  $\mathbb{Z}$ . **Hint:** If  $M$  is the maximal ideal of the valuation ring, show that  $M \cap \mathbb{Z} = p\mathbb{Z}$  for some prime  $p$  of  $\mathbb{Z}$ .
4. If  $F$  is a field, find the non-trivial valuation rings of  $F(x)$ , the quotient field of the polynomial ring  $F[x]$ , which contain  $F$ . **Hint:** Use the fact that any valuation ring of  $F(x)$  contains  $x$  or  $1/x$  and that  $F[x] \cong F[1/x]$  is a PID.
5. (a) If  $F$  is a field, show that the polynomial ring  $F[x]$  has infinitely many primes (even if  $K$  is finite).  
(b) Give an example of a PID with finitely many primes.
6. Let  $A$  be a principal ideal domain, let  $Q$  be a non-zero prime ideal of the polynomial ring  $B = A[X]$  and let  $P = Q \cap A$ .
  - (a) If  $P = pA$  with  $p$  a prime of  $A$ , show that either  $Q = pB$  or  $Q$  is the maximal ideal  $pB + fB$  with  $f \in B$  of degree  $\geq 1$  and irreducible mod  $p$ .
  - (b) If  $P = 0$ , show that  $Q = gB$  with  $g \in B$  irreducible of degree  $\geq 1$ . **Hint:** Localize  $A$  and  $B$  with respect to  $S = A - \{0\}$ .
  - (c) If  $f \in B$  is of degree  $\geq 1$  and irreducible modulo a prime  $p$  of  $A$  and  $g \in B$  is irreducible of degree  $\geq 1$ , show that  $gB \subseteq pB + fB$  iff  $f$  divides  $g$  mod  $p$ . Deduce that  $gB$  is maximal iff  $A$  has finitely many primes and  $g = a_0 + a_1x + \cdots + a_nx^n$  with  $a_0$  a unit of  $A$  and  $a_1, \dots, a_n$  divisible by all the primes of  $A$ .
  - (d) If  $f$  is algebraically closed, show that the non-zero prime ideals of the polynomial ring  $F[x, y]$  are the ideals  $(x - a, y - b)$  and  $(f)$  with  $a, b \in F$  and  $f$  irreducible. Deduce that the maximal ideals of  $F[x, y]$  are the point ideals  $(x - a, y - b)$ .