Invariance of $(\Box, \psi)$ under renaming transformations

Although it is trivial to prove, perhaps it's worth pointing out that the definition of the $(\Box, \psi)$-structure is well-defined in the sense that the items in (11.1) are independent of the choice of the particular copy of $\boldsymbol{C} \xrightarrow{\mathsf{can}} \boldsymbol{C}[r]$ (the latter is defined only up to isomorphism, whereas (11.1) is to be uniquely determined from (α, β, p) and the HAA-structure).

A useful generalization of the last remark is the invariance of the $([\beta/r]^{\theta}, \psi)$-structure under renaming transformations, generalizing the corresponding fact for the HAA-structure. Here is how this can be stated.

Suppose we have the renaming morphism $F:\boldsymbol{C} \longrightarrow \bar{\boldsymbol{C}}$, and we have $[\beta/r]^{\theta}:\boldsymbol{C}[r] \to \boldsymbol{C}$ as above. Then we can introduce a new indeterminate $\bar{r}$ over $\bar{\boldsymbol{C}}$ by requiring that $\bar{r}$ be parallel to $\bar{f}_{\mathrm{def}} Ff$. This gives us $\bar{F}:\boldsymbol{C}[r] \longrightarrow \bar{\boldsymbol{C}}[\bar{r}]$ that extends F, takes r to $\bar{r}$, and has the action on places in r given by

$$\bar{F}_r = F_f : F \circ \mathsf{s}r = F \circ \mathsf{s}f \xrightarrow{\cong} \mathsf{s}(Ff) = \mathsf{s}\bar{r} .$$

Any map $F:A \to \bar{A}$ of sets induces a functor $F:A^{\#} \longrightarrow \bar{A}^{\#}$ in an obvious way for which the action on objects is $s \longmapsto F \circ s$. Notice that for any $\mu \in A = \mathsf{A}(\boldsymbol{C}[r])$, we have that $\langle \mu \rangle \backslash r$ is an object in $A^{\#}$, and $\psi:\langle \mu \rangle \backslash r \to \langle \mu[\beta/r]^{\theta} \rangle$ is an arrow of $A^{\#}$. Therefore, with $F:\boldsymbol{C} \longrightarrow \bar{\boldsymbol{C}}$ as given, it makes sense to write $F\psi : F \circ \langle \mu \rangle \backslash r \to F \circ \langle \mu[\beta/r]^{\theta} \rangle$.

Let us indicate the effect of $\bar{F}$, and also that of F, by $\overline{(\text{-})}$; thus, for instance, we have $\bar{\beta} = F\beta$, $\bar{f} = Ff$. The following determines the isomorphism $\bar{\theta}:\mathsf{s}\bar{r} \xrightarrow{\cong} \mathsf{s}\bar{\beta}$:

$$\begin{array}{ccc} F \circ \mathsf{s}r & \xrightarrow{\quad F\theta \quad} & F \circ \mathsf{s}\beta \\ \bar{F} \downarrow \cong & \circlearrowright & \cong \downarrow \mathsf{F} \\ \mathsf{s}\bar{r} & \xrightarrow[\quad \bar{\theta} \quad]{} & \mathsf{s}\bar{\beta} \end{array} \qquad (14)$$

With this, we have $[\bar{\beta}/\bar{r}]^{\bar{\theta}}:\bar{\boldsymbol{C}}[\bar{r}]\longrightarrow\bar{\boldsymbol{C}}$, $\langle\bar{\beta}/\bar{r}\rangle^{\bar{\theta}}:\bar{\boldsymbol{C}}[\bar{r}]\longrightarrow\bar{\boldsymbol{C}}^{@}$.

We define the morphism $F^{@}:\boldsymbol{C}^{@}\longrightarrow\bar{\boldsymbol{C}}^{@}$. It is defined so that, among others, the following (with Γ, $\bar{\Gamma}$ the forgetful maps) commutes:

$$\begin{array}{ccc} \boldsymbol{C}^{@} & \xrightarrow{\quad\Gamma\quad} & \boldsymbol{C} \\ {\scriptstyle F^{@}}\downarrow & \circ & \downarrow{\scriptstyle F} \\ \bar{\boldsymbol{C}}^{@} & \xrightarrow[\quad\bar{\Gamma}\quad]{} & \bar{\boldsymbol{C}} \end{array} .$$

In particular, $F^{@}$ maps the arrow $\Lambda=(\gamma,\ \mu,\ \tau,\ t,\ \dot{\psi},\ \iota)$ to $(\bar{\mu},\bar{\gamma},\bar{\tau},\bar{\mathrm{t}},\dot{\bar{\psi}},\bar{\iota})$ where the last four items are determined by the commutative diagrams

$$\begin{array}{ccc} \mathrm{s}(\bar{\mu}) & \xrightarrow{\quad\bar{\tau}\quad} & \mathrm{s}(\bar{\gamma}) \\ \bar{F}\uparrow\cong & \circ & F\uparrow\cong \\ F\circ\mathrm{s}\mu & \xrightarrow[\quad F\tau\quad]{} & F\circ\mathrm{s}\gamma \end{array} \qquad \begin{array}{ccccc} \langle\bar{\mu}\rangle & \xleftarrow{\quad\bar{\iota}\quad} & \bar{t} & \xrightarrow{\quad\dot{\bar{\psi}}\quad} & \langle\bar{\gamma}\rangle \\ \bar{F}\uparrow\cong & \circ & \uparrow & \circ & \uparrow F \\ \bar{F}\circ\langle\mu\rangle & \xleftarrow[\quad\iota\quad]{} & t & \xrightarrow[\quad\dot{\psi}\quad]{} & F\circ\langle\gamma\rangle \end{array} \qquad (16)$$

and the fact that $\bar{\iota}$ is to be an inclusion. It is left to the reader to check that $F^{@}$ is thereby determined, and it is a morphism of multicategories.

We claim that the diagram

$$\begin{array}{ccc} \boldsymbol{C}[r] & \xrightarrow{\quad\langle\beta/r\rangle^{\theta}\quad} & \boldsymbol{C}^{@} \\ {\scriptstyle\bar{F}}\downarrow & \circ & \downarrow{\scriptstyle F^{@}} \\ \bar{\boldsymbol{C}}[\bar{r}] & \xrightarrow[\quad\langle\bar{\beta}/\bar{r}\rangle^{\bar{\theta}}\quad]{} & (\bar{\boldsymbol{C}})^{@} \end{array} \qquad (17)$$

commutes. The commutativity of the diagram on the generators of $\boldsymbol{C}$ is clear. The

commutativity on the generator r, including the action on places, is ensured by the definition of $\bar{\theta}$. The conclusions follow by the universal property of $\boldsymbol{C}[r]$.

The commutativity

$$\begin{array}{ccc} \boldsymbol{C}[r] & \xrightarrow{\quad [\beta/r]^{\theta} \quad} & \boldsymbol{C} \\ {\scriptstyle \bar{F}}\downarrow & \circ & \downarrow{\scriptstyle F} \\ \bar{\boldsymbol{C}}[\bar{r}] & \xrightarrow[\quad [\bar{\beta}/\bar{r}]^{\bar{\theta}} \quad]{} & \bar{\boldsymbol{C}} \end{array} \tag{18}$$

is a consequence.

Applying (18) to an arbitrary $\mu \in \mathsf{A}(\boldsymbol{C}[r])$, and any $\beta \in A$ parallel to r, we have

$$\overline{(\mu[\beta/r]^{\theta})} = \bar{\mu}[\bar{\beta}/\bar{r}]^{\bar{\theta}}; \tag{19}$$

whereas the application of (17) gives

$$\begin{array}{ccc} \langle \bar{\mu} \rangle \backslash \bar{r} & \xrightarrow{\quad \dot{\psi} \quad} & \langle \bar{\mu}[\bar{\beta}/\bar{r}]^{\bar{\theta}} \rangle \\ {\scriptstyle \bar{F}}\uparrow & \circ & \uparrow{\scriptstyle F} \\ F \circ (\langle \mu \rangle \backslash r) & \xrightarrow[\quad F\dot{\psi} \quad]{} & F \circ \langle \mu[\beta/r] \rangle^{\theta} \end{array} \tag{20}$$

Next, let $\alpha \in A$, $p \in |\langle \alpha \rangle|$, $f = \langle \alpha \rangle(p)$, and let the indeterminate r be parallel to f. Continuing with our "bar" notation, we have

$$\overline{\alpha(r/p)} = \bar{\alpha}(\bar{r}/\bar{p}) \tag{21}$$

$$\overline{\alpha \square_p^{\theta} \beta} = \bar{\alpha} \square_{\bar{p}}^{\bar{\theta}} \bar{\beta} \tag{22}$$

and

$$\begin{array}{ccc} F\circ\langle\alpha\rangle\backslash p & \xrightarrow{\ \psi\ } & F\circ\langle\alpha\,\square\,{}^{\theta}_{p}\beta\rangle \\ \Big\downarrow{\scriptstyle \bar{F}} & \circ & \Big\downarrow{\scriptstyle F} \\ \langle\bar{\alpha}\rangle\backslash\bar{p} & \xrightarrow{\ \bar{\psi}\ } & \langle\bar{\alpha}\,\square\,{}^{\bar{\theta}}_{\bar{p}}\bar{\beta}\rangle \end{array} \qquad (23)$$

For (21), let $\mu=\alpha(r/p)$; we have to see that $\bar{\mu}$ is mapped by $[\bar{f}/\bar{r}]$ to $\bar{\alpha}$, and that $\bar{\mu}$ is $\bar{p}'$-special, for $\bar{p}'=[\bar{f}/\bar{r}]^{-1}(p)$. The first fact follows from (18) for $\beta=f$ and $\theta=\mathrm{id}$. The same special case of (18), now applied to the actions on places, gives

$$\begin{array}{ccc} F\circ\langle\mu\rangle & \xrightarrow{\ [f/r]\ } & \langle\alpha\rangle \\ \Big\downarrow{\scriptstyle \bar{F}} & \circ & \Big\downarrow{\scriptstyle F} \\ \langle\bar{\mu}\rangle & \xrightarrow[\ [\bar{f}/\bar{r}]\ ]{} & \langle\bar{\alpha}\rangle \end{array} \quad . \qquad (24)$$

It follows that the isomorphism $\bar{F}:\bar{F}\circ\langle\mu\rangle \xrightarrow{\ \cong\ } \langle\bar{\mu}\rangle$ takes p' to $\bar{p}'$; now, the desired second fact follows from the presence of the same isomorphism.

(22) is a special case of (19).

(23) follows from (20) and (24).

The commutative law for the $\square$-composition

Before we complete defining amalgamation for $\square$ with the φ-item, let us see the commutativity law for $\square$: for α , β , γ in $\mathsf{A}(\boldsymbol{C})$,

$$(\alpha\,\square\,{}^{\theta}_{p}\beta)\,\square\,{}^{\tau}_{\bar{q}}\gamma \;=\; (\alpha\,\square\,{}^{\tau}_{q}\gamma)\,\square\,{}^{\theta}_{\bar{p}}\beta \qquad (25)$$

provided

$p, q\in |\langle\alpha\rangle|$, $p\neq q$,

$$\theta : s(\langle\alpha\rangle(p)) \xrightarrow{\cong} s\beta\,,\ \ t(\langle\alpha\rangle(p)) = t\beta\,,$$
$$\tau : s(\langle\alpha\rangle(q)) \xrightarrow{\cong} s\gamma\,,\ \ t(\langle\alpha\rangle(q)) = t\gamma\,,$$
$$\bar{q} = \psi^{\Box}[\alpha, \beta, p, \theta](q)\,,\ \ \bar{p} = \psi^{\Box}[\alpha, \gamma, q, \tau](p) \qquad (25')$$

Note that $\langle\alpha\rangle(p) = \langle\alpha \Box_{q}^{\tau} \gamma\rangle(\bar{p})$; thus the $\Box_{\bar{p}}^{\theta}$-composition makes sense; similarly for $\Box_{\bar{q}}^{\tau}$.

We introduce two different indeterminates r and s over $\boldsymbol{C}$, with $r \| f$ $_{\mathrm{d}\bar{\mathrm{e}}\mathrm{f}}\alpha(p)$, $s \| g$ $_{\mathrm{d}\bar{\mathrm{e}}\mathrm{f}}\alpha(q)$, and consider the multicategory $\boldsymbol{C}[r, s]$ obtained by simultaneously adjoining r and s to $\boldsymbol{C}$. We have the inclusion $\boldsymbol{C} \longrightarrow \boldsymbol{C}[r, s]$, with the universal property that for any $F : \boldsymbol{C} \longrightarrow \boldsymbol{E}$, any r', s' in $\boldsymbol{E}$ and any θ, τ such that $r' \|^{\theta} Ff$, $s' \|^{\tau} Fg$, there is a unique arrow $\bar{F} : \boldsymbol{C}[r, s] \longrightarrow \boldsymbol{E}$ extending F, mapping r to r', s to s', and having the actions $\bar{F} \circ sr \longrightarrow sr'$, $\bar{F} \circ ss \longrightarrow ss'$ equal to $\theta : F \circ sf \longrightarrow sr'$, $\tau : F \circ sg \longrightarrow ss'$, respectively.

The multicategories $\boldsymbol{C}[r]$, $\boldsymbol{C}[s]$ can be regarded as included in $\boldsymbol{C}[r, s]$: the canonical morphisms $\boldsymbol{C}[r] \longrightarrow \boldsymbol{C}[r, s]$, $\boldsymbol{C}[s] \longrightarrow \boldsymbol{C}[r, s]$, given by the universal properties of $\boldsymbol{C}[r]$ and $\boldsymbol{C}[s]$, respectively, are inclusions. In fact, the inclusion $\boldsymbol{C}[r] \longrightarrow \boldsymbol{C}[r, s]$ endows $\boldsymbol{C}[r, s]$ with the the universal property of $\boldsymbol{C}[r][s]$. Similarly, we can regard $\boldsymbol{C}[r, s]$ as being $\boldsymbol{C}[s][r]$.

We fix an HAA structure on $\boldsymbol{C}[r, s]$, which may be, and is, assumed to extend those on $\boldsymbol{C}$, $\boldsymbol{C}[r]$ and $\boldsymbol{C}[s]$.

For any $\beta, \gamma \in \boldsymbol{C}$ and θ, τ such that $\beta \|^{\theta} f$, $\gamma \|^{\tau} g$, we have the morphism

$$[\beta/r,\ \gamma/s]^{\theta,\ \tau} : \boldsymbol{C}[r, s] \longrightarrow \boldsymbol{C}$$

determined by the properties that it extends the identity $\boldsymbol{C} \longrightarrow \boldsymbol{C}$, takes r to β, s to γ, and has the actions $\theta : sr \longrightarrow s\beta$, $\gamma : ss \longrightarrow s\gamma$ on places.

Note that, besides $[\beta/r]^{\theta}:\mathbf{C}[r]\longrightarrow\mathbf{C}$ and $[\gamma/s]^{\tau}:\mathbf{C}[s]\longrightarrow\mathbf{C}$, we also have $[\beta/r]^{\theta}:\mathbf{C}[r,s]\longrightarrow\mathbf{C}[s]$ and $[\gamma/s]^{\tau}:\mathbf{C}[r,s]\longrightarrow\mathbf{C}[r]$, by our identifications of $\mathbf{C}[r,s]$ with $\mathbf{C}[s][r]$ and $\mathbf{C}[r][s]$.

We have the following commutative diagram:



(26)

The commutativity is checked, as usual, on the generators. A special case is



(27)

$\alpha(r/p, s/q)$ is defined to be the unique $\mu\in\mathsf{A}(\mathbf{C}[r,s])$ for which $\mu[f/r, g/s]=\alpha$, and, for $p^{\#}, q^{\#}\in|\mathsf{s}\mu|$ defined by $[f/r, g/s](p^{\#})=p$, $[f/r, g/s](q^{\#})=q$, we have that μ is $(r/p^{\#}, s/q^{\#})$-special, meaning that $(\mathsf{s}\mu)(p^{\#})=r$, $(\mathsf{s}\mu)(q^{\#})=s$, and for all $n\in|\mathsf{s}\mu|-\{p^{\#}, q^{\#}\}$, $(\mathsf{s}\mu)(n)\in L$. The proof of the existence and uniqueness of $\alpha(r/p, s/q)$ is similar to that of (5).

With the actions $[f/r]:|\langle\alpha(r/p)\rangle|\xrightarrow{\cong}|\langle\alpha\rangle|$, $[g/s]:|\langle\alpha(s/q)\rangle|\xrightarrow{\cong}|\langle\alpha\rangle|$, we write, as before, $p'=[f/r]^{-1}(p)$, $q'=[g/s]^{-1}(q)$; but also, we put $q^{*}=[f/r]^{-1}(q)$, $p^{*}=[g/s]^{-1}(p)$.

Let $\mu=\alpha(r/p, s/q)$. By (27), we get

$$\mu[f/r][g/s]=\alpha. \qquad (28)$$

With $[g/s]: |\langle \mu[f/r] \rangle| \longrightarrow |\langle \alpha \rangle|$, let

$$\tilde{q} = [g/s]^{-1}(q) \, . \tag{29}$$

Consider

$$\begin{array}{ccc} [f/r] \circ \langle \mu \rangle & \xrightarrow[\cong]{[f/r]} & \langle \mu[f/r] \rangle \\ q^{\#} & \longmapsto & \tilde{q} \end{array} \, . \tag{30}$$

The fact that $q^{\#} \mapsto \tilde{q}$ follows from (27). The fact that μ is $(r/p^{\#}, s/q^{\#})$-special, the existence of the isomorphism (30), and the fact $f \in L$ together immediately imply that $\mu[f/r]$ is $s/\tilde{q}$-special. This, with (28), (29), tells us that

$$\mu[f/r] = \alpha(s/q) \tag{31}$$

and $\tilde{q} = q'$. Of course, similarly,

$$\mu[g/s] = \alpha(r/p) \, . \tag{32}$$

Notice that (31) is the first step towards proving the fact that $\mu = \alpha(s/q)(r/p^{*})$. (27) implies that $p^{\#} \xmapsto{[f/r]} p^{*}$. Thus, what remains is to show

μ is $(r/p^{\#})$-special over $\boldsymbol{C}[s]$.

But this is obvious from μ being $(r/p^{\#}, s/q^{\#})$-special. We conclude that

$$\mu = \alpha(r/p, s/q) = \alpha(s/q)(r/p^{*}) = \alpha(r/p)(s/q^{*}) \tag{33}$$

We repeat that we have the following:

$$p^{\#} \xmapsto{[f/r]} p^{*} \, , \qquad q^{\#} \xmapsto{[f/r]} q'$$

$$p^{\#} \overset{[g/s]}{\longmapsto} p' , \qquad q^{\#} \overset{[g/r]}{\longmapsto} q^{*} \ .$$

The commutativity

$$\begin{array}{ccc} \boldsymbol{C}[r,s] & \xrightarrow{\ [\beta/r]^{\theta}\ } & \boldsymbol{C}[s] \\ {\scriptstyle [g/s]}\downarrow & \circ & \downarrow{\scriptstyle [g/s]} \\ \boldsymbol{C}[r] & \xrightarrow[\ [\beta/r]^{\theta}\ ]{} & \boldsymbol{C} \end{array}$$

is a(nother) special case of (26). When applied to μ, we obtain that

$$\mu[\beta/r]^{\theta}[g/s] = \mu[g/s][\beta/r]^{\theta} = \alpha(r/p)[\beta/r]^{\theta} \ \bar{\mathrm{def}} \ \alpha \square_{p}^{\theta} \beta \ .$$

This is the first step towards showing the fact that $(\alpha \square_{p}^{\theta} \beta)(s/\bar{q}) = \mu[\beta/r]^{\theta}$; here, $\bar{q}$ is as in (25). It remains to see that, for $\hat{q} \overset{[g/s]}{\longmapsto} \bar{q}$, we have that $\mu[\beta/r]^{\theta}$ is $s/\hat{q}$-special. Because of (33), we may write $\mu[\beta/r]^{\theta}$ as $\alpha(s/q) \square_{p}^{\theta} {}_{*}\beta$.

We apply (23) with $[g/s]:\boldsymbol{C}[s] \longrightarrow \boldsymbol{C}$ as F, and $[g/s]:\boldsymbol{C}[s][r] \longrightarrow \boldsymbol{C}[r]$ as $\bar{F}$. The $\bar{\theta}$ in (23) is determined by (14). Now, $\bar{r}=r$, and by the definition of $[g/s]$, the actions $[g/s]:\mathrm{s}r \to \mathrm{s}r$, $[g/s]:\mathrm{s}\beta \to \mathrm{s}\beta$ are identities. Therefore, $\bar{\theta}=\theta$. We obtain

$$\begin{array}{ccc} [g/s] \circ \langle \alpha(s/q) \rangle \backslash p^{*} & \xrightarrow[\cong]{\ [g/s]\ } & \langle \alpha \rangle \backslash p \\ \downarrow{\scriptstyle \psi^{\square}} & \circ & \downarrow{\scriptstyle \psi^{\square}} \\ [g/s] \circ \langle \alpha(s/q) \square_{p}^{\theta} {}_{*}\beta \rangle & \xrightarrow[\cong]{\ [g/s]\ } & \langle \alpha \square_{p}^{\theta} \beta \rangle \end{array}$$

From the last commutativity, it follows that the left vertical $\psi^{\square}$ maps q' to $\hat{q}$. Now, looking at (12), and knowing that $\alpha(s/q)$ is q'-special, we see that $\alpha(s/q) \square_{p}^{\theta} {}_{*}\beta$ is $s/\hat{q}$-special.

We conclude, as was foreseen, that

$$(\alpha \square^{\theta}_{p} \beta)(s/\bar{q}) = \mu[\beta/r]^{\theta} \quad . \tag{34}$$

But then

$$(\alpha \square^{\theta}_{p} \beta) \square^{\tau}_{\bar{q}} \gamma = (\alpha \square^{\theta}_{p} \beta)(s/\bar{q})[\gamma/s]^{\tau} = \mu[\beta/r]^{\theta}[\gamma/s]^{\tau} \overset{*}{=} \mu[\beta/r,\ \gamma/s]^{\theta,\ \tau} ,$$

where the equality marked with $*$ is by (26). Of course, the equality

$$(\alpha \square^{\tau}_{q} \gamma) \square^{\theta}_{\bar{p}} \beta = \mu[\beta/r,\ \gamma/s]^{\theta,\ \tau}$$

is true also by reasons of symmetry. We have proved (25).

By the way, $\mu[\beta/r,\ \gamma/s]^{\theta,\ \tau}$ may be reasonably denoted as $\alpha(\square^{\theta}_{p}\beta,\ \square^{\tau}_{q}\gamma)$, and it amounts to having defined an instance of simultaneous $\square$-composition directly. We have

$$\alpha(\square^{\theta}_{p}\beta,\ \square^{\tau}_{q}\gamma) = (\alpha \square^{\theta}_{p} \beta) \square^{\tau}_{\bar{q}} \gamma = (\alpha \square^{\tau}_{q} \gamma) \square^{\theta}_{\bar{p}} \beta \quad .$$

We can extend the just-given arguments to prove the part of the coherence conditions relating to commutativity that refers to the ψ-maps alone (the quadrangle in 1.(xi) in [H/M/P]), by first defining a new "simultaneous" ψ-map $\psi\colon \langle \alpha \rangle \setminus \{p, q\} \longrightarrow \alpha(\square^{\theta}_{p}\beta,\ \square^{\tau}_{q}\gamma)$, and then showing that both composites of ψ's in the quadrangle are equal to it. However, it is not really necessary to do this, because the arguments used in [H/M/P] for the coherence conditions via invariance remain available and easy.

The $\varphi^{\square}$–structure

Next, we deal with the φ-part of the amalgamation structure for the composition $\square$.

The definition is again a "multicategory recursion", except that the domain on which we define the assignment will be a proper subclass of the arrows in $\boldsymbol{C}[r]$. (We simply do not see how to define the φ-structure meaningfully on the whole of $\boldsymbol{C}[r]$; this is, of course, in contrast to the ψ-structure which was defined on the whole of $\boldsymbol{C}[r]$.)

The reader may see already now that every time we want to define recursively a partial function F on the arrows, with F having the domain X, we can do it as if we were defining a total function, provided X is a *convex* class: whenever $\alpha \circ_i \beta \in X$, both α and β belong to X as well: we extend F by assigning an arbitrary value to each argument not in X.

Let us call an arrow μ of $\boldsymbol{C}[r]$ *normal* if there is *at most one* $p \in |\langle \mu \rangle|$ for which $\langle \mu \rangle(p)=r$. Thus, the normal arrows are those that are in $\boldsymbol{C}$, and those that are r-*special*: r/p-special for some p. Clearly, the normal arrows form a convex class.

Let us fix $\beta \in A$ and θ so that θ-parallel to r.

Define the multicategory $\boldsymbol{C}^{(\beta)}$ as having the same objects as $\boldsymbol{C}$, and having arrows of the form

$$\Xi = (\gamma, \mu, \tau : s\mu \xrightarrow{\cong} s\gamma, m, \langle \gamma \rangle \xleftarrow{\varphi} m \cdot \langle \beta \rangle)$$

where $\mu \in A(\boldsymbol{C}[r])$, $\gamma \in A(\boldsymbol{C})$, $m \in \{0, 1\}$; in fact, $m=0$ iff $\mu \in \boldsymbol{C}$ or μ is not normal, thus $m=1$ iff μ is r-special. We mean: $0 \cdot \langle \beta \rangle = \emptyset$, the empty tuple, $1 \cdot \langle \beta \rangle = \langle \beta \rangle$.

As before in the case of $\boldsymbol{C}^{@}$, we have a forgetful map $\Gamma : \boldsymbol{C}^{(\beta)} \longrightarrow \boldsymbol{C}$ that is the identity on objects, takes $\Xi = (\gamma, \mu, ...)$ to γ, and is strict. This implies, for instance, that $s\Xi = s\gamma$, $t\Xi = t\gamma$.

Composition in $\boldsymbol{C}^{(\beta)}$: given $\Xi_j = (\gamma_j, \mu_j, \tau_j, m_j, \varphi_j)$ $(j=1, 2)$, $j \in |s\gamma_1|$ such that $\gamma = \gamma_1 \circ_i \gamma_2$ is well-defined, to define $\Xi = \Xi_1 \circ_i \Xi_2 = (\gamma, \mu, \tau, m, \varphi)$. It follows that for $i' = \tau_1^{-1}(i)$, $\mu = \mu_1 \circ_{i'} \mu_2$ is well-defined. We calculate $\tau : s\mu \xrightarrow{\cong} s\gamma$ by the diagram (11'). We form the commutative diagram



(35)

with the right-hand vertical side being a coproduct; of course, this means $m=m_1+m_2$. Case 1: μ is normal. In this case $m\leq 1$, since both μ_1 , μ_2 have to be normal, and at most one can be r-special. We complete the definition by requiring that ι_j be the identity if $m=m_j=1$ ($j=1, 2$); when $m_j=0$, ι_j must be the initial map $\emptyset\to m\cdot\langle\beta\rangle$; this determines φ as well. Now, we define $\Xi_1\circ_i\Xi_2$ to be $(\gamma, \mu, \tau, m, \varphi)$. Case 2: μ is not normal. Now, we let $\Xi_1\circ_i\Xi_2$ be $(\gamma, \mu, \tau, 0, !)$ (somewhat arbitrarily).

When verifying either of the two multicategory laws for $\boldsymbol{c}^{(\beta)}$: associativity or commutativity, we have two triple composites, call them $\Xi_{(1)}=(\mu_{(1)}, \ldots)$, $\Xi_{(2)}=(\mu_{(2)}, \ldots)$, and we want to show they are equal. Only the φ map is in question. First note that $\mu_{(1)}$ is normal iff $\mu_{(2)}$ is normal iff all three factors are normal and at most one is r-special. When $\mu_{(1)}$, $\mu_{(2)}$ are not normal, there is nothing to show. When they are normal, all the m's and φ's involved are defined "normally" according to the picture (35), with additional specifications on the ι's as stated above; using the coherence laws for the σ's and ρ's involved, the task of showing the φ's to be equal is now easy.

We use $\boldsymbol{c}^{(\beta)}$ for defining a morphism

$$\begin{aligned}\langle\beta/r\rangle^{\theta} : \boldsymbol{c}[r] &\longrightarrow \boldsymbol{c}^{(\beta)}\\ g &\longmapsto (g, g, \mathrm{id}:sg\to sg, \langle g\rangle \xleftarrow{!} 0\cdot\langle\beta\rangle)\\ r &\longmapsto (r, \beta, \theta:s\beta\to sr, \langle\beta\rangle \xleftarrow{\mathrm{id}} 1\cdot\langle\beta\rangle)\end{aligned}$$

such that, also, $\langle\beta/r\rangle^{\theta}\circ\Gamma = [\beta/r]^{\theta}$ (as before), and let Φ act on $\alpha(r/p)$ (with α p-special as usual), to get

$$\alpha(r/p)\langle\beta/r\rangle^{\theta} = (\alpha\Box_p\beta,\ \alpha(r/p),\ \tau,\ 1,\ \langle\alpha\Box_p\beta\rangle \xleftarrow{\varphi[\alpha,\ \beta,\ p,\ \theta]} \langle\beta\rangle)$$

(it is obvious, by induction, that the first two components are as shown; that the codomain of φ is $\langle\text{-}\rangle$ applied to the first item; and finally, that the domain of φ is $\langle\beta\rangle$ and not $\emptyset$). This is our definition of the φ-component of the amalgamation structure for $\Box$.

(36) Proposition In

$$\langle\alpha\rangle\backslash p \xrightarrow{\psi} \langle\alpha\Box_p^{\theta}\beta\rangle \xleftarrow{\varphi} \langle\beta\rangle \qquad (\psi=\psi[\alpha,\ \beta,\ p,\theta],\ \varphi=\varphi[\alpha,\ \beta,\ p,\theta])$$

φ and ψ are the coprojections of a coproduct diagram in the category $L^{\#}$.

Proof: induction.