

(9) 1. Name and sketch the following surfaces:

(a) $x^2 = 4(y - z^2)$

(b) $r^2 = z^2 + 4$

(c) $\rho \tan \phi (\cos \theta + \sin \theta) = 2 \sec \phi - \rho$

(8) 2. The motion of an object is given by $\mathbf{r}(t) = \langle \ln t^2, \sqrt{8t}, t^2 \rangle$ for $t > 0$.

(a) Find parametric equations of the tangent line to the trajectory at time $t = 1$.

(b) Find an expression for the speed v of the object in terms of t .

(c) Find the curvature κ of the trajectory at time $t = 1$.

(d) Find the tangential and normal components of acceleration at time $t = 1$.

(4) 3. Find the limit if it exists or show that it does not exist.

(a) $\lim_{(x,y) \rightarrow (0,0)} \frac{(-9x + y)^2}{81x^2 + y^2}$

(b) $\lim_{(x,y) \rightarrow (0,0)} \frac{3x^2 + 5y^2}{\sqrt{3x^2 + 5y^2 + 1} - 1}$

(4) 4. Consider $z = f(x, y) = \sqrt{x^2 + y^2}$.

(a) Find the differential dz .

(b) Use dz to find an approximation for $f(3.06, 3.92)$.

(7) 5. Consider the surface $F(x, y, z) = xz + 2x^2y + y^2z^3 = 11$ and the point $P(2, 1, 1)$.

(a) Find the directional derivative of F at P in the direction of $\mathbf{v} = \langle -1, 1, 1 \rangle$.

(b) Find the maximum rate of increase of F at P ?

(c) In what direction (unit vector) does F increase the fastest at P ?

(d) Find the equation (in $ax + by + cz = d$ form) of the tangent plane to the surface at P .

(e) Assume that $Q \neq P$ is a point on this tangent plane. What is the directional derivative of F at P in the direction \overrightarrow{PQ} ?

(f) Find $\frac{\partial z}{\partial y}$.

(3) 6. Assume f is a differentiable function and $z = yf(x^2 - y^2)$. Show that $y \frac{\partial z}{\partial x} + x \frac{\partial z}{\partial y} = \frac{xz}{y}$

(5) 7. Find and classify the critical points of $f(x, y) = x^4 + 2y^2 - 4xy$.

(5) 8. Use the method of Lagrange multipliers to find the point on the sphere $x^2 + y^2 + z^2 = 4$ that is farthest from the point $P(1, -1, 1)$.

(12) 9. Evaluate the integrals.

(a) $\int_0^8 \int_{\sqrt[3]{x}}^2 \sin(y^4) dy dx$

(b) $\int_0^1 \int_0^{\sqrt{1-x^2}} \cos(x^2 + y^2 + 4) dy dx$

$$(c) \int_0^3 \int_0^{\sqrt{9-y^2}} \int_{\sqrt{x^2+y^2}}^{\sqrt{18-x^2-y^2}} (x^2 + y^2 + z^2) dz dx dy.$$

- (6) 10. **Set up, but do not evaluate**, triple integrals to find the volume of the region between the sphere $x^2 + y^2 + z^2 = 19$ and the upper sheet of the hyperboloid $z^2 - x^2 - y^2 = 1$, $z > 0$ in
- Cartesian coordinates
 - cylindrical coordinates
- (4) 11. Using a suitable change of variables, find the following double integral over T where T is the triangle enclosed by the lines $y - x = 0$, $y + x = 2$ and the x -axis.

$$\iint_T (x + y)^3 dx dy$$

- (2) 12. Let $f(x) = \sum_{n=1}^{\infty} \frac{n(x+6)^{3n}}{(3n+1)!}$; evaluate $f^{(27)}(-6)$.
- (5) 13. Find the Maclaurin series for the following functions and state the radius of convergence.

$$(a) f(x) = \frac{x^3}{5+x^2}$$

$$(b) g(x) = \frac{\arctan(3x^2)}{x}$$

- (5) 14. Approximate $\int_0^{0.1} x e^{-x^3} dx$ to six decimal places of accuracy.

$$(7) 15. \text{ Let } f(x) = \frac{1}{\sqrt{x}}$$

- Use the binomial series to expand $f(x)$ as a power series centered at $x = 9$ and state the radius of convergence.
- If $T_2(x)$ is used to approximate $f(9.5)$, give an upper bound on the error using the Lagrange form of the remainder.

- (8) 16. Consider the curve \mathcal{C} having parametric equations: $\begin{cases} x = 2 \cos t + 1 \\ y = 3 \sin t \end{cases}$ where $t \in \mathbb{R}$.

- Find dy/dx and d^2y/dx^2 .
 - Find all the points on \mathcal{C} where the tangent line is vertical or horizontal.
 - Eliminate the parameter t to express the curve in the form $f(x, y) = d$. Using this equation, identify and sketch \mathcal{C} .
 - Set up, but do not evaluate**, an integral expression that gives the area bounded by the curve.
- (6) 17. Consider the polar curves $r = \cos(3\theta)$ and $r = \frac{1}{2}$.
- Sketch the two curves on the same axes.
 - Set up, but do not evaluate**, an integral expression for the area of the region common to both curves.
 - Set up, but do not evaluate**, the integral needed to find the length of $r = \cos(3\theta)$.