

DROPLET PHASE IN A NONLOCAL ISOPERIMETRIC PROBLEM UNDER CONFINEMENT

STAN ALAMA AND LIA BRONSARD

Department of Mathematics and Statistics
McMaster University, Hamilton, ON, Canada

RUSTUM CHOKSI

Department of Mathematics and Statistics
McGill University, Montréal, QC, Canada

IHSAN TOPALOGLU

Department of Mathematics and Applied Mathematics
Virginia Commonwealth University, Richmond, VA, USA

(Communicated by Juncheng Wei)

ABSTRACT. We address small volume-fraction asymptotic properties of a nonlocal isoperimetric functional with a confinement term, derived as the sharp interface limit of a variational model for self-assembly of diblock copolymers under confinement by nanoparticle inclusion. We introduce a small parameter η to represent the size of the domains of the minority phase, and study the resulting droplet regime as $\eta \rightarrow 0$. By considering confinement densities which are spatially variable and attain a unique nondegenerate maximum, we present a two-scale asymptotic analysis wherein a separation of length scales is captured due to competition between the nonlocal repulsive and confining attractive effects in the energy. A key role is played by a parameter M which gives the total volume of the droplets at order η^3 and its relation to existence and non-existence of Gamow's Liquid Drop model on \mathbb{R}^3 . For large values of M , the minority phase splits into several droplets at an intermediate scale $\eta^{1/3}$, while for small M minimizers form a single droplet converging to the maximum of the confinement density.

1. Introduction. In this paper we study the asymptotic properties of a nonlocal isoperimetric functional with a confinement term. This functional appears as the sharp interface limit of a model of diblock copolymer/nanoparticle blend where a large number static nanoparticles serve as a confinement term, penalizing the energy outside of a fixed region. We choose a scaling regime in which the mass fraction between the two phases tends to zero, but with very strong nonlocal interactions, wherein the minimizing phases resemble small spherical inclusions of one phase in a large sea of the second phase. This is often called the *droplet* regime, in which the sparse A -phase can be described effectively by the droplet centers, which we think of as *particles* [8]. The advantage of the droplet regime is to permit the decomposition

2000 *Mathematics Subject Classification.* 35Q70, 49Q20, 49S05, 74N15, 82D60.

Key words and phrases. Nonlocal isoperimetric problem, Γ -convergence, self-assembly of diblock copolymers, confinement, phase separation, uniformly charged liquid.

of the nonlocal effects into *self-effects* on the shape of a single droplet and *interaction effects* between different particles.

Let $\eta > 0$ be a small parameter, which will represent the scale of the radius of the droplets. We consider periodic configurations $v \in BV(\mathbb{T}^3; \{0, \eta^{-3}\})$, defined on the standard unit torus \mathbb{T}^3 , which we model as the unit cube $[-\frac{1}{2}, \frac{1}{2}]$ with periodic boundary conditions, and subject to a mass constraint,

$$\int_{\mathbb{T}^3} v dx = M,$$

for a constant $M > 0$. For a fixed function $\rho \in C(\mathbb{T}^3)$, representing the local density of nanoparticles we analyze the energy functional,

$$\mathbb{E}_\eta(v) := \eta \int_{\mathbb{T}^3} |\nabla v| + \eta \|v - M\|_{H^{-1}(\mathbb{T}^3)}^2 - \int_{\mathbb{T}^3} v(x)\rho(x) dx \tag{1.1}$$

in the limit as $\eta \rightarrow 0$. The physical background and justification for the choices of the parameters appearing in \mathbb{E}_η will be discussed in the Appendix. Here, the first term in the energy denotes the *total variation* of the function v and is defined as

$$\int_{\mathbb{T}^3} |\nabla v| := \sup \left\{ \int_{\mathbb{T}^3} v \operatorname{div} \varphi dx : \varphi \in C_0^1(\mathbb{T}^3; \mathbb{R}^3), |\varphi(x)| \leq 1 \right\}.$$

For characteristic functions $\chi_A \in BV(\mathbb{T}^3; \{0, 1\})$, this term computes the perimeter of the interface ∂A . The second term, on the other hand, is nonlocal and is defined via

$$\|v - M\|_{H^{-1}(\mathbb{T}^3)}^2 = \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} G(x, y)v(x)v(y) dx dy = \int_{\mathbb{T}^3} |\nabla w_v|^2 dx,$$

where $G(x, y)$ is the mean-zero Laplace Green's function on \mathbb{T}^3 , and w_v is the solution of $-\Delta w_v = v - M$ on \mathbb{T}^3 .

The last term reflects the presence of nanoparticles of density ρ in the sample, which attract the $\{\eta^3 v = 1\}$ phase of the copolymer. Thus we expect that droplets accumulate near maxima of the density ρ . For most of the paper we will assume that the density has a unique non-degenerate global maximum at the origin.

- (H1) $\rho \in C(\mathbb{T}^3)$ with $\rho \geq 0$.
- (H2) $\rho_{\max} := \rho(0) > \rho(x)$ for all $x \in \mathbb{T}^3 \setminus \{0\}$.
- (H3) $\rho \in C^2(B_r)$ for some $r > 0$ and

$$\begin{aligned} \rho(x) &= \rho_{\max} - q(x) + o(|x|^2) \quad \text{as } |x| \rightarrow 0 \quad \text{where} \\ q(x) &:= \frac{1}{2} \sum_{i,j=1}^3 H_{ij} x_i x_j \quad \text{for } x = (x_1, x_2, x_3) \in \mathbb{T}^3 \end{aligned} \tag{1.2}$$

and H_{ij} are the entries of the Hessian matrix of $-\rho$ given by

$$H_{ij} = -\frac{\partial^2 \rho}{\partial x_i \partial x_j}(0) \quad \text{with } H_{ij} x_i x_j \geq C|x|^2 \quad \text{for some constant } C > 0.$$

In section 5 we will examine how to treat ρ which have degenerate but homogeneous behavior $\rho_{\max} - \rho(x) \sim |x|^q$, $q > 2$, at the maximum point.

We also note that it is not necessary to impose periodic boundary conditions to observe the droplet splitting and confinement described in this paper; for instance, it is both physically and mathematically reasonable to replace the torus \mathbb{T}^3 by a smooth bounded domain $\mathcal{D} \subset \mathbb{R}^3$ with Neumann boundary conditions.

Droplet splitting. In their study of the NLIP under droplet scaling without confinement, (that is, with $\rho(x) \equiv 0$.) Choksi and Peletier [8] show that a minimizing sequence v_η will decompose into droplets of radius on the order of η ; that is, v_η is approximated by a finite sum of weighted Dirac measures, with centers $x_\eta^i \in \mathbb{T}^3$. The shape of the droplets is determined by the leading order term in an expansion of the energy: blowing up at scale η , they arrive at a nonlocal isoperimetric problem (NLIP) on all of \mathbb{R}^3 . In the unconfined setting of [8], the location of the centers x_η^i is determined by the next order term in the energy expansion, a Coulomb-like repulsion arising from the nonlocal term. The result is convergence of minimizers to a discrete pattern of droplets spread out on \mathbb{T}^3 , where the interdroplet distances are bounded below.

For our energy (1.1) with nonconstant density ρ we observe a different qualitative picture. The confinement term in E_η will be felt at the level of particle interactions, drawing the centers x_η^i towards the global maximum of the nanoparticle density $\rho(x)$. This is the essential difference between our problem and that of [8]: as the droplet centers will be drawn towards a single point we must isolate individual droplets at a much smaller scale. The interaction terms in the energy expansion of [8] are of order one, and their method suffices to discern droplets at arbitrarily small distance of order one. This is not enough for our setting, where the second-order term in the energy expansion is of much smaller order than the order of approximation of the first-order term in [8]. Thus, we must develop a more refined method of isolating the droplets and calculating the interaction distance on a much smaller scale.

To motivate our main result, let us consider an ansatz for v_η , which will form the basis of the upper bound on the energy (see Lemma 2.4.) We will suppose that minimizers form n droplets, centered at points $x_\eta^i = \delta p_i$ with fixed p_1, \dots, p_n and $\delta = \delta(\eta) \rightarrow 0$, reflecting the tendency of the droplets to approach the maximizer of $\rho(x)$ at a particular rate (to be determined.) We thus define an admissible test configuration,

$$\nu_\eta := \sum_{i=1}^n \eta^{-3} w_i \left(\frac{x - \delta p_i}{\eta} \right), \tag{1.3}$$

where $w_i \in BV(\mathbb{R}^3; \{0, 1\})$ are characteristic functions representing the droplet sets, blown up by scale η to \mathbb{R}^3 . We define $m^i := \int_{\mathbb{R}^3} w_i dx$, and note that the constraint $\int_{\mathbb{T}^3} \nu_\eta dx = M$ forces $\sum_{i=1}^n m^i = M$.

We may then evaluate $E_\eta(\nu_\eta)$ asymptotically, expressing the H^{-1} -norm in terms of the Green's function, $G(x, y) \sim 1/4\pi|x - y|$, as the points coalesce. Assuming the distance between droplet centers $\delta(\eta) \gg \eta$, a back-of-the-envelope calculation yields,

$$\begin{aligned} E_\eta(\nu_\eta) \simeq & \sum_{i=1}^n \left[\int_{\mathbb{R}^3} |\nabla w_i| + \|w_i\|_{H^{-1}(\mathbb{R}^3)}^2 \right] - M\rho_{\max} \\ & + \left[\frac{\eta}{\delta} \sum_{\substack{i,j=1 \\ i \neq j}}^n \frac{m^i m^j}{4\pi|p_i - p_j|} + \delta^2 \sum_{i=1}^n m^i q(p_i) \right]. \end{aligned} \tag{1.4}$$

Optimizing the right-hand expression over $\delta > 0$ (holding all other quantities fixed) we predict the droplet separation scale $\delta = \mathcal{O}(\eta^{1/3})$ (see Figure 1). Thus, we expect droplet separation to occur on **two different scales** in the length and in energy.

The droplets themselves have characteristic length η , and contribute to the energy at order one, while the pattern they form is observed on the much larger length scale $\eta^{1/3}$, which affects the energy to order $O(\eta^{2/3})$.

The first sum in the expansion is the energy for Gamow’s Liquid Drop model, and this indicates how droplet shape is determined via minimization. Droplet profiles z_i should be minimizers for the nonlocal isoperimetric problem on \mathbb{R}^3 :

$$e_0(m) := \inf \left\{ \int_{\mathbb{R}^3} |\nabla z| + \|z\|_{H^{-1}(\mathbb{R}^3)}^2 : z \in BV(\mathbb{R}^3, \{0, 1\}), \int_{\mathbb{R}^3} z dx = m \right\}, \quad (1.5)$$

where

$$\|z\|_{H^{-1}(\mathbb{R}^3)}^2 = \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{z(x)z(y)}{4\pi|x-y|} dx dy.$$

This problem has been extensively studied, in the form above and in various generalizations; see [4, 10, 12, 16–18, 21, 22] as well as [7] for a review on the liquid drop model. In [8] it was conjectured that there exists a critical mass m_c such that minimizers are balls for $m \leq m_c$ and fail to exist otherwise. The results will play an important role in this article and we summarize them in the next theorem, which may be found in [18] (see also [21]):

Theorem 1.1 (Knüpfer-Muratov). *There exist three constants $0 < m_{c_0} \leq m_{c_1} \leq m_{c_2}$ such that the following hold.*

- (i) *If $m \leq m_{c_1}$ then $e_0(m)$ admits a minimizer. If $m \leq m_{c_0}$ the minimizer is unique (up to translation), and is given by the ball of volume m . The ball of volume m ceases to be the minimizer of $e_0(m)$ for $m > m_{c_0}$.*
- (ii) *If $m > m_{c_2}$ then $e_0(m)$ does not admit a minimizer.*

To date, it remains an open problem to prove or disprove whether any (or all) of the constants m_{c_i} , $i = 0, 1, 2$, above are pairwise equal.

Returning to the energy expansion of the ansatz (1.4), we may now complete the heuristic picture of minimizers of E_η in terms of the Gamow functional. Defining the “good” set of m for which the nonlocal isoperimetric problem attains a minimizer,

$$\mathcal{M}_0 := \{m > 0: e_0(m) \text{ admits a minimizer}\},$$

we observe that when the total mass $M \in \mathcal{M}_0$ (for instance, when $M \leq m_{c_1}$ is small enough,) there is no need for splitting, and we anticipate that minimizers v_η remain connected as $\eta \rightarrow 0$. However, if $M \notin \mathcal{M}_0$ then we infer that according to (1.4) minimizers will split into droplets with mass $m^i \in \mathcal{M}_0$, for which there is an optimal droplet blowup function $z_i \in BV(\mathbb{R}^3; \{0, 1\})$, and with $\sum_{i=1}^n m^i = M$. That is, the collection of droplet masses must lie in the special set,

$$\mathcal{M} := \left\{ \{m^i\}_{i=1}^n : n \in \mathbb{N}, m^i \geq 0, \sum_{i=1}^n m^i = M, \right.$$

and $e_0(m^i)$ admits a minimizer for each i $\left. \right\}$.

These droplets will then arrange themselves in a pattern with separation distance $\delta(\eta) = O(\eta^{1/3})$, so as to minimize the interaction energy formed by combining Coulomb repulsion and confinement,

$$F_{m^1, \dots, m^n}(p_1, \dots, p_n) := \sum_{\substack{i, j=1 \\ i \neq j}}^n \frac{m^i m^j}{4\pi|p_i - p_j|} + \sum_{i=1}^n m^i q(p_i),$$

with $q(x)$ the Hessian of $\rho(x)$ near $x = 0$, defined in (H3). (See Figure 1).




FIGURE 1. The attraction to the origin and scaling at the rate $\delta = \eta^{1/3}$.

Our main result is to confirm this expected behavior by means of a precise asymptotic expansion of the energy of minimizers. We prove:

Theorem 1.2. *Let v_η be minimizers of E_η in $BV(\mathbb{T}^3, \{0, \eta^{-3}\})$ with $\int_{\mathbb{T}^3} v_\eta dx = M$.*

- (i) *For any $r > 0$, $\text{supp } v_\eta \subset B_r(0) \subset \mathbb{T}^3$ for all sufficiently small $\eta > 0$.*
- (ii) *If $M \in \mathcal{M}_0$, then there exists a subsequence of $\eta \rightarrow 0$ and points $y_\eta \in \mathbb{T}^3$ with $|y_\eta| \leq C\eta^{1/2}$ such that*

$$v_\eta - \eta^{-3} z_M \left(\frac{x - y_\eta}{\eta} \right) \rightarrow 0 \quad \text{in } L^1(\mathbb{T}^3),$$

where z_M attains the minimum $e_0(M)$.

- (iii) *If $M \notin \mathcal{M}_0$, then, there exists a subsequence of $\eta \rightarrow 0$, $n \in \mathbb{N}$, $\{m^i\}_{i=1}^n \in \mathcal{M}$, and distinct points $\{x_\eta^1, \dots, x_\eta^n\}$ such that:*

$$v_\eta - \sum_{i=1}^n \eta^{-3} w_i \left(\frac{x - x_\eta^i}{\eta} \right) \rightarrow 0, \quad \text{in } L^1(\mathbb{T}^3), \quad (1.6)$$

where $w_i \in BV(\mathbb{R}^3, \{0, 1\})$ attains the minimum in $e_0(m^i)$, $i = 1, \dots, n$. In addition,

$$\eta^{-\frac{1}{3}} x_\eta^i \rightarrow x_i \in \mathbb{R}^3, \quad \text{and}$$

$$E_\eta(v_\eta) = e_0(M) - M\rho_{\max} + \eta^{2/3} \left[\sum_{i=1}^n m^i q(x_i) + \frac{1}{4\pi} \sum_{\substack{i,j=1 \\ i \neq j}}^n \frac{m^i m^j}{|x_i - x_j|} \right] + o\left(\eta^{2/3}\right).$$

Moreover, the expression in brackets above is minimized by the choice of points $\{x_1, \dots, x_n\}$ given the values $\{m^i\}_{i=1}^n \in \mathcal{M}$.

Remark 1.3. As noted above, we may also consider ρ which attain a unique degenerate maximum, assuming a different homogeneity, i.e., $\rho(x) - \rho_0 \sim |x|^q$, $q > 2$, around the maximum point. The analysis will be essentially the same, except for the scale of droplet separation and the form of the interaction energy; see Section 5, and Figure 2 in which minimizers of the resulting interaction energy $F_{m^1, \dots, m^n}(p_1, \dots, p_n)$

are illustrated for different values of q . In case ρ attains its global maximum at a finite number of points we expect the same general behavior near one or several of the maximizers, but the distribution of mass around each one depends subtly on the shape of the function at each point.

Part (ii) of Theorem 1.2 implies that when $M \in \mathcal{M}_0$ there is no splitting of the droplet: apart from a possible drift of $\mathcal{O}(\eta^{1/2})$ away from the origin, it is essentially a rescaling of the minimizer of the nonlocal isoperimetric problem at scale η . It is known [18, 21] that the $e_0(M)$ minimizers (when they exist) have connected compact support. Thus, the theorem asserts that splitting can only occur if the $e_0(M)$ problem fails to achieve a minimum and that the separation must be $\mathcal{O}(\eta^{1/3})$ and at no other scale. For clarity, we note that for η sufficiently small, the sum in (1.6) is disjointly supported in $B_{\frac{1}{4}}(0) \subset \mathbb{R}^3$, and may thus be treated as a function in $BV(\mathbb{T}^3; \{0, \eta^{-3}\})$.

The interaction energy F_{m^1, \dots, m^n} , which determines the location of the droplets at $\eta^{1/3}$ -scale, is (for fixed mass distribution $\{m^i\}_{i=1}^n$) coercive and clearly attains a minimum among all possible configurations $\{x_i\}_{i=1}^n$ in \mathbb{R}^3 . The two-dimensional analogue of F_{m^1, \dots, m^n} , with logarithmic repulsion, was studied in [15], but in the case of equal masses. (The motivation behind [15] came from Ginzburg–Landau vortices, which have quantized flux playing the role of the mass in F_{m^1, \dots, m^n} .) We know of no studies of F_{m^1, \dots, m^n} which address the fundamental question of how to optimally choose both $\{m^i\}_{i=1}^n$ and $\{x_i\}_{i=1}^n$ to minimize F_{m^1, \dots, m^n} .

We also note that similar concentration or coalescence phenomena appear in various examples of singularly perturbed variational problems, notably in low density phases in a uniformly charged liquid [19] and for vortices in the 2D Ginzburg–Landau model of superconductivity [27]. See, for example, Chapter 7 of [27], in which finitely many vortices concentrate at specific points in a sample subjected to an external magnetic field h_{ex} of order $\ln \kappa$, where κ is the Ginzburg–Landau parameter. As above, there are two length scales: the radii of the vortex cores are given by κ^{-1} , and the distance between coalescing vortices is of order $h_{ex}^{-1/2}$. Note in particular the similarity of their renormalized energy (11.2) and our interaction energy F_{m^1, \dots, m^n} .

Finally, attractive-repulsive nonlocal interaction energies of the form F_{m^1, \dots, m^n} have attracted much interest lately in connection with models of pattern formation in biological aggregations such as swarming or collective behavior of multi-agent systems [6, 9, 24, 29]. In these models the particles can be approximated by a density distribution leading to a continuum-level energy, and minimization with respect to these densities yields challenging variational and geometric problems (see e.g. [5, 11]).

Our paper is organized as follows: In the appendix, we describe the physical origins of the model as well as the derivation of the droplet scaling regime which we will use in the paper. Section 2 describes the concentration structure of minimizers. The most important result there is a key concentration-compactness lemma (Lemma 2.5) which gives the framework for our result, and is based on a recent compactness result for sets of bounded perimeter due to Frank and Lieb [12] which can conveniently be used to decompose the support of a minimizing sequence into sets of diameter $O(\eta)$. In Section 3 we use the regularity of minimizers to refine the concentration lemma and obtain a lower bound with a precise error estimate of order $o(\eta^{2/3})$, needed to isolate the interaction terms in the energy. In Section 4

we prove Theorem 1.2 by matching upper and lower bounds on the energy. Finally in Section 5 we discuss second-order approximations for degenerate confining penalization densities ρ which have degenerate maxima at a single point.




FIGURE 2. Minimizing configurations of the second-order energy F_{m^1, \dots, m^n} with equal mass $m^i = 1/100$ for 100 particles with varying powers q of degenerate penalization $\rho(x) - \rho_0 \sim |x|^q$. Minimizing configurations are obtained as steady-states of the gradient flow of the energy F_{m^1, \dots, m^n} .

2. The concentration structure of minimizers. We begin by defining the first-order limit functional,

$$E_0(v) := \begin{cases} \sum_{i=1}^{\infty} e_0(m^i) - m^i \rho(x_i) & \text{if } v = \sum_{i=1}^{\infty} m^i \delta_{x_i}, \{x_i\} \text{ distinct, and } m^i \geq 0, \\ +\infty & \text{otherwise,} \end{cases}$$

where e_0 is defined by (1.5). The following first-order Gamma convergence result follows directly from [8], since the confinement term is a continuous perturbation of the functional in [8, Theorem 4.3].

Theorem 2.1 (First-order limit). *The sequence of functionals E_η Γ -converge to E_0 as $\eta \rightarrow 0$ in the space of Radon measures with respect to the weak- $*$ convergence.*

Remark 2.2. Theorem 2.1 holds true for any confinement term defined via a penalizing measure $\rho \in C(\mathbb{R}^3)$ and is not special to those which satisfy (H1)–(H3).

Remark 2.3. A standard consequence of Γ -convergence, combined with the compactness of a sequence of minimizers, is that minimizers of E_η converge weakly in the space of Radon measures to a minimizer of the limiting energy E_0 . Hence, using

the simple fact that $e_0(M) \leq \sum_{i=1}^{\infty} e_0(m^i)$ (cf. [8]) we may conclude that for any family of mass-constrained minimizers of E_η ,

$$v_\eta \rightharpoonup M\delta_0 \quad \text{and} \quad \lim_{\eta \rightarrow 0} E_\eta(v_\eta) = e_0(M) - M\rho_{\max} \quad (2.1)$$

where $\rho_{\max} := \rho(0)$.

As Theorem 2.1 shows the first-order approximation does not include the nonlocal non-self interaction effects of the energy E_η . To capture these effects we will look at the second-order approximation. This limit will depend on the specific form of the penalizing measure ρ as we will normalize the energy E_η by subtracting the ground state of the first-order limit E_0 . We just noted that as $\eta \rightarrow 0$, mass constrained minimizers of E_η concentrate at the origin. How the sequence collapses to the origin depends on the size of the mass constraint M . When M is sufficiently large so that a minimizer of e_0 fails to exist, we show that minimizers of E_η split at a scale larger than η as $\eta \rightarrow 0$, while for small enough M we show that no splitting may occur.

To this end, recall the notation,

$$\mathcal{M}_0 := \{M > 0: e_0(M) \text{ admits a minimizer}\}.$$

It is clear from Theorem 1.1 that \mathcal{M}_0 is not empty. Moreover, $M < m_{c_1}$ is a sufficient condition for $M \in \mathcal{M}_0$ and $M > m_{c_2}$ is sufficient for $M \notin \mathcal{M}_0$; however, the necessity of either condition is an open problem.

As a first illustration of the role played by \mathcal{M}_0 , we state an ensemble of upper bounds on the minimum energy of E_η :

Lemma 2.4. *For any $n \in \mathbb{N}$, $\{p_i\}_{i=1}^n$ distinct fixed points in \mathbb{R}^3 , and $\{m^i\}_{i=1}^n \in \mathcal{M}$, we have*

$$\min_{\int_{\mathbb{T}^3} v = M} E_\eta(v) \leq (e_0(M) - M\rho_{\max}) + \eta^{2/3} \left[\sum_{i=1}^n m^i q(p_i) + \frac{1}{4\pi} \sum_{\substack{i,j=1 \\ i \neq j}}^n \frac{m^i m^j}{|p_i - p_j|} \right] + o(\eta^{2/3}). \quad (2.2)$$

Moreover, if $M \in \mathcal{M}_0$, then

$$\min_{\int_{\mathbb{T}^3} v dx = M} E_\eta(v) \leq (e_0(M) - M\rho_{\max}) + \mathcal{O}(\eta). \quad (2.3)$$

Thus, if $M \in \mathcal{M}_0$ the upper bound on the energy is vastly improved. In fact, we expect that fewer droplets (smaller n) generally should yield a smaller upper bound, and so energy minimization should split the mass M into the smallest number of pieces which permit $\{m^i\}_{i=1}^n \in \mathcal{M}$. However, we are not aware of any such results concerning the minimization of F_{m^1, \dots, m^n} with respect to the masses m^i .

The proof of the upper bound follows along the lines of the heuristic argument presented in the Introduction, and we defer it to Section 4.

As usual, the difficulty in verifying the desired structure of energy minimizers lies mostly in developing a matching lower bound on the energy. The first and crucial step is obtaining a compactness result which establishes the existence of points in \mathbb{T}^3 which are separated by a scale much larger than η apart, so that weighted Dirac-delta measures at these points approximate v_η . This result also gives the existence of components of $\text{supp } v_\eta$ whose supports are η -rescalings of minimizers of the nonlocal isoperimetric problem, and thus achieve the minimum of the first-order energy E_0 . In addition, we establish that there can only be finitely many distinct

components for minimizers of E_η , and in case there is a unique component we obtain a stronger convergence result:

Lemma 2.5. *For each $\eta > 0$ let v_η be a minimizer of E_η with $\int_{\mathbb{T}^3} v_\eta dx = M$. Then there exists a subsequence $\eta \rightarrow 0$, $n \in \mathbb{N}$, $\{m^i\}_{i=1}^n \subset (0, \infty)$, $\{x_\eta^i\}_{i=1}^n \subset \mathbb{T}^3$ and functions $w_\eta^i \in BV(\mathbb{T}^3; \{0, 1/\eta^3\})$ with $\|w_\eta^i\|_{L^1(\mathbb{T}^3)} = m^i + o(1)$ as $\eta \rightarrow 0$ such that for $n \geq 2$*

$$\frac{|x_\eta^i - x_\eta^j|}{\eta} \rightarrow \infty \text{ for every } i \neq j, \tag{2.4}$$

$$\left\| v_\eta - \sum_{i=1}^n w_\eta^i \right\|_{L^1(\mathbb{T}^3)} \rightarrow 0 \tag{2.5}$$

$$e_0(m^i) \text{ is attained for each } i = 1, 2, \dots, n, \quad \text{and} \quad e_0(M) = \sum_{i=1}^n e_0(m^i), \tag{2.6}$$

$$\liminf_{\eta \rightarrow 0} E_\eta(v_\eta) \geq \liminf_{\eta \rightarrow 0} \sum_{i=1}^n E_\eta(w_\eta^i) \geq \sum_{i=1}^n e_0(m^i) - M\rho_{\max}. \tag{2.7}$$

Moreover, if $n = 1$, then $M \in \mathcal{M}_0$, and there exist points $x_\eta \in \mathbb{T}^3$ such that

$$v_\eta - \eta^{-3} z_M \left(\frac{x - x_\eta}{\eta} \right) \rightarrow 0 \text{ in } L^1(\mathbb{T}^3) \text{ as } \eta \rightarrow 0, \tag{2.8}$$

where z_M attains the minimum $e_0(M)$.

We remark that the last part of the lemma already indicates that splitting is only to be expected when the $e_0(M)$ problem has no minimizer. Indeed, the second-order behavior is very different depending on whether $M \in \mathcal{M}_0$ or not. However, it will be more convenient to use the dichotomy suggested by Lemma 2.5 and separate the cases $n = 1$ and $n \geq 2$ determined above; the full connection to \mathcal{M}_0 will only be apparent later on.

The proof of Lemma 2.5 is based on blowing up the minimizing sets $A_\eta = \text{supp } v_\eta$ at order η . When we blow up the minimizing set A_η we are effectively thinking of it as a subset of \mathbb{R}^3 , and are able to utilize a technical concentration-compactness result for sets of finite perimeter by Frank and Lieb [12], but this can only work locally. As the balls $B_r(x_0) \subset \mathbb{T}^3$ with radius $0 < r < \frac{1}{2}$ lift isometrically to \mathbb{R}^3 , blowup makes sense for sets defined in these balls. For $x_0 \in \mathbb{T}^3$ and $0 < r < \frac{1}{2}$, we may thus define a family of diffeomorphisms,

$$\begin{cases} \Phi_{\eta,r}(\cdot; x_0) : B_r(x_0) \subset \mathbb{T}^3 \rightarrow B_{\frac{r}{\eta}}(0) \subset \mathbb{R}^3, \\ x \mapsto \Phi_{\eta,r}(x; x_0) = \frac{x - x_0}{\eta}, \end{cases}$$

where the last quantity is lifted to the covering space \mathbb{R}^3 . Thus, the map $\Phi_{\eta,r}(x; x_0)$ performs a blowup of $B_r(x_0)$, viewed as a subset of \mathbb{R}^3 . For simplicity, we will often abuse notation and write $\Phi_{\eta,r}(S; x_0) = \eta^{-1}(S - x_0)$ for the blowup to \mathbb{R}^3 of a set $S \subset B_r(x_0) \subset \mathbb{T}^3$, and for $U \in B_{\frac{r}{\eta}}(0) \subset \mathbb{R}^3$, we write $\eta U + x_0 = \Phi_{\eta,r}^{-1}(U; x_0) \subset \mathbb{T}^3$.

Proof of Lemma 2.5. For any $\eta > 0$ let $v_\eta \in BV(\mathbb{T}^3; \{0, 1/\eta^3\})$ with $\int_{\mathbb{T}^3} v_\eta dx = M$ be a minimizer of E_η . Such a minimizer exists by the direct method in the calculus of variations. By the first-order limit, Theorem 2.1, the energy is bounded above, and we have that $E_\eta(v_\eta) \leq e_0(M) - M\rho_{\max} + o(1)$.

Step 1. We restrict to a neighborhood of zero. Write $v_\eta = \eta^{-3}\chi_{A_\eta}$, with $A_\eta \subset \mathbb{T}^3$. Since by (H3) zero is a strict local maximum of ρ , and thus we may find $\epsilon > 0$ and $\delta \in (0, \frac{1}{8})$ so that

$$\rho(x) < \rho_{\max} - \epsilon \quad \text{for all } x \in \mathbb{T}^3 \setminus B_\delta(0).$$

We now claim that for this δ , $\eta^{-3}|A_\eta \setminus B_\delta(0)| \rightarrow 0$. Indeed, assume the contrary, so there is a value > 0 such that along some subsequence $\eta \rightarrow 0$, $\eta^{-3}|A_\eta \setminus B_\delta(0)| \gg 0$. However, in that case,

$$\begin{aligned} \int_{\mathbb{T}^3} \rho v_\eta dx &= \eta^{-3} \int_{A_\eta \cap B_\delta(0)} \rho dx + \eta^{-3} \int_{A_\eta \setminus B_\delta(0)} \rho dx \\ &\leq \rho_{\max} \eta^{-3} |A_\eta \cap B_\delta(0)| + (\rho_{\max} - \epsilon) \eta^{-3} |A_\eta \setminus B_\delta(0)| \\ &\leq \rho_{\max} M - \epsilon < \rho_{\max} M. \end{aligned}$$

Using the lower bound from [8] on the first two terms in the energy together with the upper bound from Theorem 2.1,

$$e_0(M) - M\rho_{\max} \geq \lim_{\eta \rightarrow 0} E_\eta(v_\eta) \geq e_0(M) - \rho_{\max}M + \epsilon,$$

which gives the desired contradiction.

We next show we make only a very small error in the energy of minimizers by restricting to a ball around the origin. Indeed, following the proof of Lemma 2.2 of [11], for each η there exists a radius $\delta_\eta \in (2\delta, 3\delta)$ so that the surface area of the intersection,

$$\mathcal{H}^2(\partial B_{\delta_\eta}(0) \cap A_\eta) \rightarrow 0.$$

(The argument involves noting that $|A_\eta \cap (B_{3\delta} \setminus B_{2\delta})| \rightarrow 0$ and using Fubini's theorem in spherical coordinates.) Define the sets

$$\Omega_\eta := A_\eta \cap B_{\delta_\eta}(0), \quad \Xi_\eta := A_\eta \setminus B_{\delta_\eta}(0),$$

and functions

$$v_\eta^0 := \eta^{-3}\chi_{\Omega_\eta}.$$

Then, we may conclude that

$$\begin{aligned} \int_{\mathbb{T}^3} |\nabla \chi_{A_\eta}| &\leq \int_{\mathbb{T}^3} |\nabla \chi_{\Omega_\eta}| + \int_{\mathbb{T}^3} |\nabla \chi_{\Xi_\eta}| \\ &\leq \int_{\mathbb{T}^3} |\nabla \chi_{A_\eta}| + 2\mathcal{H}^2(\partial B_{\delta_\eta}(0) \cap A_\eta) + o(1) \leq \int_{\mathbb{T}^3} |\nabla \chi_{A_\eta}| + o(1), \end{aligned}$$

and therefore

$$\liminf_{\eta \rightarrow 0} \int_{\mathbb{T}^3} |\nabla \chi_{A_\eta}| \geq \liminf_{\eta \rightarrow 0} \int_{\mathbb{T}^3} |\nabla \chi_{\Omega_\eta}|,$$

and

$$\lim_{\eta \rightarrow 0} \left[\int_{\mathbb{T}^3} |\nabla \chi_{A_\eta}| - \int_{\mathbb{T}^3} |\nabla \chi_{\Omega_\eta}| - \int_{\mathbb{T}^3} |\nabla \chi_{\Xi_\eta}| \right] = 0.$$

Furthermore, $\eta^{-3}|\Xi_\eta| \rightarrow 0$ (globally on \mathbb{T}^3) implies that $\|v_\eta - v_\eta^0\|_{L^1(\mathbb{T}^3)} \rightarrow 0$, and

$$\begin{aligned} \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} G(x, y) v_\eta(x) v_\eta(y) dx dy &= \eta^{-6} \int_{A_\eta} \int_{A_\eta} G(x, y) dx dy \\ &= \eta^{-6} \int_{\Omega_\eta} \int_{\Omega_\eta} G(x, y) dx dy + o(1), \end{aligned}$$

and

$$\int_{\mathbb{T}^3} v_\eta \rho dx = \eta^{-3} \int_{A_\eta} \rho dx = \eta^{-3} \int_{\Omega_\eta} \rho dx + o(1) = \int_{\mathbb{T}^3} v_\eta^0 \rho dx + o(1),$$

and hence we have

$$E_\eta(v_\eta) \geq E_\eta(v_\eta^0) + o(1), \quad \text{and} \quad \int_{\mathbb{T}^3} v_\eta^0 dx = \eta^{-3} |\Omega_\eta| = M - o(1). \quad (2.9)$$

Effectively, we may consider v_η^0 to be our minimizing sequence, with support in $B_{3\delta}(0) \subset \mathbb{T}^3$ (although it only satisfies the mass constraint to within $o(1)$.) However, it will often be convenient to treat the truncated support $\Omega_\eta \subset B_{3\delta}(0) \subset \mathbb{R}^3$ as belonging to the Euclidean space \mathbb{R}^3 .

Step 2. Finding a concentrating set. We blow up on the scale of η , to obtain

$$\widehat{\Omega}_\eta := \eta^{-1} \Omega_\eta = \Phi_{\eta, 3\delta}(\Omega_\eta; 0) \subset \mathbb{R}^3.$$

We note that in fact $\widehat{\Omega}_\eta \subset B_{\frac{3\delta}{\eta}}(0)$, and by (2.9), $|\widehat{\Omega}_\eta| = \eta^{-3} |\Omega_\eta| = M - o(1)$. For a set $\widehat{\Omega} \subset \mathbb{R}^3$, we define the energy

$$\mathcal{G}(\widehat{\Omega}) := \int_{\mathbb{R}^3} |\nabla \chi_{\widehat{\Omega}}| + \int_{\widehat{\Omega}} \int_{\widehat{\Omega}} \frac{1}{4\pi|x-y|} dx dy,$$

corresponding to the Liquid Drop model (see cf. [12, 18] or (1.5)). For $x, y \in B_{3\delta}(0) \subset \mathbb{T}^3$ we may express the Green's function on the torus locally,

$$G(x, y) = \frac{1}{4\pi|x-y|} + g(x-y), \quad (2.10)$$

in terms of the Newtonian potential and the regular part $g \in C^\infty(B_{\frac{3}{8}}(0))$, and observe that the contribution to $E_\eta(v_\eta^0)$ due to the regular part is of the order $O(\eta)$:

$$\begin{aligned} & \eta \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} v_\eta^0(x) v_\eta^0(y) G(x, y) dx dy \\ &= \eta \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} \left[\frac{1}{4\pi|x-y|} + g(x-y) \right] v_\eta^0(x) v_\eta^0(y) dx dy \\ &= \eta^{-5} \int_{\Omega_\eta} \int_{\Omega_\eta} \frac{1}{4\pi|x-y|} dx dy + \mathcal{O}(\eta) = \int_{\widehat{\Omega}_\eta} \int_{\widehat{\Omega}_\eta} \frac{1}{4\pi|x-y|} dx dy + \mathcal{O}(\eta) \end{aligned}$$

Consequently, we may relate the energies before and after blowup via

$$E_\eta(v_\eta^0) = \mathcal{G}(\widehat{\Omega}_\eta) - \int_{\mathbb{T}^3} v_\eta^0 \rho dx + \mathcal{O}(\eta). \quad (2.11)$$

To prove the lemma we employ concentration-compactness methods to show that, if $\widehat{\Omega}_\eta$ is not tight, then it must split into a finite number of components, each of which minimizes the blowup energy \mathcal{G} . To locate a first component, we observe that $|\widehat{\Omega}_\eta| = M - o(1)$ for all $\eta > 0$, and apply Proposition 2.1 of [12] to obtain a set $\Omega_1 \subset \mathbb{R}^3$ of positive measure and finite perimeter, and a sequence of points $\{y_\eta^1\}_{\eta>0} \subset \mathbb{R}^3$ such that

$$\begin{cases} \chi_{\widehat{\Omega}_\eta}(\cdot + y_\eta^1) \rightarrow \chi_{\Omega_1} \text{ in } L^1_{\text{loc}} \text{ as } \eta \rightarrow 0, \quad 0 < |\Omega_1| \leq \liminf_{\eta \rightarrow 0} |\widehat{\Omega}_\eta|, \text{ and} \\ \int_{\mathbb{R}^3} |\nabla \chi_{\Omega_1}| \leq \liminf_{\eta \rightarrow 0} \int_{\mathbb{R}^3} |\nabla \chi_{\widehat{\Omega}_\eta}|. \end{cases} \quad (2.12)$$

If it turns out that $|\Omega_1| = M$, then we need go no further, and the lemma holds with $n = 1$. Indeed, let

$$\tilde{\Omega}_\eta := \widehat{\Omega}_\eta - y_\eta^1.$$

Since $\chi_{\tilde{\Omega}_\eta} \rightarrow \chi_{\Omega_1}$ in $L^1_{loc}(\mathbb{R}^3)$ and $\{\chi_{\tilde{\Omega}_\eta}\}_{\eta>0}$ is a tight sequence, in fact we have convergence globally in L^1 norm. Then using the fact that $\tilde{\Omega}_\eta + y_\eta^1 = \widehat{\Omega}_\eta$ and the translation invariance of the total variation and H^{-1} -norms, and Step 1, we conclude that

$$\begin{aligned} \mathcal{G}(\Omega_1) &\leq \liminf_{\eta \rightarrow 0} \mathcal{G}(\widehat{\Omega}_\eta) = \liminf_{\eta \rightarrow 0} \left[\mathbf{E}_\eta(v_\eta^0) + \int_{\mathbb{T}^3} v_\eta^0 \rho dx \right] \\ &\leq \liminf_{\eta \rightarrow 0} \left[\mathbf{E}_\eta(v_\eta) + \int_{\mathbb{T}^3} v_\eta \rho dx \right] \leq e_0(M), \end{aligned}$$

using the upper bound from the first order limit. Thus $e_0(M)$ is attained by χ_{Ω_1} , and we may conclude that (with $r = 3\delta$ and $x_\eta = \eta y_\eta^1 \in \mathbb{T}^3$),

$$\eta^3 v_\eta^0(\eta x + x_\eta) \chi_{B_{r/\eta}(0)} = \chi_{\tilde{\Omega}_\eta}(x) \rightarrow \chi_{\Omega_1} \quad \text{in } L^1_{loc}(\mathbb{R}^3).$$

The desired conclusion (2.8) then follows by a change of variables from \mathbb{R}^3 to $B_r(0) \subset \mathbb{T}^3$, and $\eta^{-3}|\Xi_\eta| \rightarrow 0$ from Step 1.

In the remainder of the proof we may therefore assume that $0 < |\Omega_1| < M$.

Step 3. When $0 < |\Omega_1| < M$ we expect there is no unique choice for the set Ω_1 and translations y_η^1 for which we obtain local convergence. Following Lions [20] (see also Nam and van den Bosch [25]) we define

$$\begin{aligned} \mathcal{M}(\{\widehat{\Omega}_\eta\}) &:= \sup\{|\Omega| : \text{there exist } y_\eta \in \mathbb{R}^3 \text{ and } \Omega \subset \mathbb{R}^3 \text{ such that} \\ &\quad \chi_{\widehat{\Omega}_\eta - y_\eta} \rightarrow \chi_\Omega \text{ in } L^1_{loc}(\mathbb{R}^3)\}. \end{aligned}$$

We may then choose Ω_1 and $y_\eta^1 \in \mathbb{R}^3$ such that $|\Omega_1| > \frac{1}{2} \mathcal{M}(\{\widehat{\Omega}_\eta\}) > 0$, for which (2.12) holds. Define

$$m^1 := |\Omega_1| \quad \text{and} \quad x_\eta^1 := \eta y_\eta^1.$$

By Lemma 2.2 of [12] there exist $\{r_{1,\eta}\}_{\eta>0} \subset (0, \infty)$ such that the sets

$$U_\eta^1 := \tilde{\Omega}_\eta \cap B_{r_{1,\eta}}, \quad V_\eta^1 := \tilde{\Omega}_\eta \cap (\mathbb{R}^3 \setminus \bar{B}_{r_{1,\eta}})$$

satisfy

$$\chi_{U_\eta^1} \rightarrow \chi_{\Omega_1} \text{ in } L^1(\mathbb{R}^3), \quad \chi_{V_\eta^1} \rightarrow 0 \text{ in } L^1_{loc}(\mathbb{R}^3), \quad \text{and} \quad m_\eta^1 := |U_\eta^1| \rightarrow |\Omega_1| = m^1.$$

Let $\widehat{\Omega}_\eta^1 := V_\eta^1 + y_\eta^1$, with $|\widehat{\Omega}_\eta^1| = |\widehat{\Omega}_\eta| - m_\eta^1 = M - m^1 + o(1) \in (0, M)$. Moreover,

$$\lim_{k \rightarrow \infty} \left(\int_{\mathbb{R}^3} |\nabla \chi_{\tilde{\Omega}_\eta}| - \int_{\mathbb{R}^3} |\nabla \chi_{U_\eta^1}| - \int_{\mathbb{R}^3} |\nabla \chi_{V_\eta^1}| \right) = 0,$$

and

$$\liminf_{\eta \rightarrow 0} \int_{\mathbb{R}^3} |\nabla \chi_{U_\eta^1}| \geq \int_{\mathbb{R}^3} |\nabla \chi_{\Omega_1}|.$$

Using Lemma 2.3 in [12], the splitting also extends to the nonlocal energy,

$$\int_{\widehat{\Omega}_\eta} \int_{\widehat{\Omega}_\eta} \frac{1}{|x-y|} dx dy = \int_{U_\eta^1} \int_{U_\eta^1} \frac{1}{|x-y|} dx dy + \int_{V_\eta^1} \int_{V_\eta^1} \frac{1}{|x-y|} dx dy + o(1)$$

and

$$\int_{U_\eta^1} \int_{U_\eta^1} \frac{1}{|x-y|} dx dy = \int_{\Omega_1} \int_{\Omega_1} \frac{1}{|x-y|} dx dy + o(1).$$

Thus, the blowup energy splits up to an error of order $o(1)$,

$$\mathcal{G}(\widehat{\Omega}_\eta) = \mathcal{G}(U_\eta^1) + \mathcal{G}(V_\eta^1) + o(1) = \mathcal{G}(\Omega_1) + \mathcal{G}(\widehat{\Omega}_\eta^1) + o(1). \quad (2.13)$$

To return to the original scale, define the functions

$$u_\eta^1(x) := \frac{1}{\eta^3} \chi_{U_\eta^1} \left(\frac{x - x_\eta^1}{\eta} \right) = \frac{1}{\eta^3} \chi_{U_\eta^1} (\Phi_{\eta, \delta_\eta}(x; x_\eta^1)), \quad w_\eta^1(x) := \frac{1}{\eta^3} \chi_{\Omega_1} \left(\frac{x - x_\eta^1}{\eta} \right), \quad (2.14)$$

$$v_\eta^1(x) := \frac{1}{\eta^3} \chi_{V_\eta^1} \left(\frac{x - x_\eta^1}{\eta} \right) = \frac{1}{\eta^3} \chi_{\widehat{\Omega}_\eta^1} \left(\frac{x}{\eta} \right).$$

Then, $v_\eta^0 = u_\eta^1 + v_\eta^1$, each supported in $B_{3\delta}(0) \subset \mathbb{T}^3$, and applying (2.11) to (2.13) we obtain the following splitting,

$$\mathbb{E}_\eta(v_\eta^0) = \mathbb{E}_\eta(u_\eta^1) + \mathbb{E}_\eta(v_\eta^1) + o(1) \geq \mathbb{E}_\eta(w_\eta^1) + \mathbb{E}_\eta(v_\eta^1) + o(1).$$

Step 4. We now repeat Step 3 for the remainder term $\{v_\eta^1\}_{\eta>0}$, with associated blowup set $\widehat{\Omega}_\eta^1$, of mass $|\widehat{\Omega}_\eta^1| = M - m_\eta^1 = M - m^1 + o(1) \in (0, M)$. Using the compactness results in [12] as in Step 3, we obtain translations $y_\eta^2 \in \mathbb{R}^3$, a limiting set Ω_2 with $|\Omega_2| \in (0, M - m^1]$ and $|\Omega_2| > \frac{1}{2} \mathcal{M}(\{\widehat{\Omega}_\eta^1\})$, and a new remainder set $\widehat{\Omega}_\eta^2$, for which the energy splits to within an error of $o(1)$. We iterate this procedure to create sequences of translations, limiting sets, and remainder sets, as long as the limit sets do not exhaust the total mass M . More specifically, given the remainder $v_\eta^{i-1} = \eta^{-3} \chi_{\widehat{\Omega}_\eta^{i-1}}(x/\eta)$, we obtain translations $y_\eta^i \in \mathbb{R}^3$ and $\Omega_i \subset \mathbb{R}^3$, with $|\Omega_i| > \frac{1}{2} \mathcal{M}(\{\widehat{\Omega}_\eta^{i-1}\})$, and a disjoint decomposition $(\widehat{\Omega}_\eta^{i-1} - y_\eta^i) = U_\eta^i \cup V_\eta^i$, with $r_{i,\eta} \in (0, \infty)$, $U_\eta^i := (\widehat{\Omega}_\eta^{i-1} - y_\eta^i) \cap B_{r_{i,\eta}}$, $V_\eta^i := (\widehat{\Omega}_\eta^{i-1} - y_\eta^i) \setminus B_{r_{i,\eta}}$, and radii $r_{i,\eta}$ chosen such that $\chi_{U_\eta^i} \rightarrow \chi_{\Omega_i}$ in $L^1(\mathbb{R}^3)$ and $\chi_{V_\eta^i} \rightarrow 0$ in $L^1_{\text{loc}}(\mathbb{R}^3)$. We set

$$m_\eta^i := |U_\eta^i| \rightarrow |\Omega_i| := m^i, \quad \widehat{\Omega}_\eta^i := V_\eta^i + y_\eta^i, \quad x_\eta^i := \eta y_\eta^i, \\ u_\eta^i := \frac{1}{\eta^3} \chi_{U_\eta^i} \left(\frac{x - x_\eta^i}{\eta} \right), \quad w_\eta^i := \frac{1}{\eta^3} \chi_{\Omega_i} \left(\frac{x - x_\eta^i}{\eta} \right), \quad v_\eta^i := \frac{1}{\eta^3} \chi_{\widehat{\Omega}_\eta^i} \left(\frac{x}{\eta} \right).$$

Then, as in Step 3, we may assert that for any $k \geq 1$:

$$|\Omega_k| > \frac{1}{2} \mathcal{M}(\{\widehat{\Omega}_\eta^{k-1}\}); \quad (2.15)$$

$$M = \sum_{i=1}^k m_\eta^i + |\widehat{\Omega}_\eta^k| + o(1) = \sum_{i=1}^k m^i + |\widehat{\Omega}_\eta^k| + o(1); \quad (2.16)$$

$$\lim_{\eta \rightarrow 0} \left\| v_\eta - \sum_{i=1}^k w_\eta^i - v_\eta^k \right\|_{L^1(\mathbb{T}^3)} = 0 = \lim_{\eta \rightarrow 0} \left| \widehat{\Omega}_\eta \Delta \left(\widehat{\Omega}_\eta^k \cup \bigcup_{i=1}^k \Omega_i \right) \right|; \quad (2.17)$$

$$\liminf_{\eta \rightarrow 0} \left[\mathcal{G}(\widehat{\Omega}_\eta) - \sum_{i=1}^k \mathcal{G}(\Omega_i) - \mathcal{G}(\widehat{\Omega}_\eta^k) \right] \geq 0; \quad (2.18)$$

$$\liminf_{\eta \rightarrow 0} \left[\mathbb{E}_\eta(v_\eta) - \sum_{i=1}^k \mathbb{E}_\eta(w_\eta^i) - \mathbb{E}_\eta(v_\eta^k) \right] \geq 0. \tag{2.19}$$

Since for each $i < k$, $(U_\eta^k + (y_\eta^k - y_\eta^i)) \subset V_\eta^i \rightarrow 0$ locally, but $\chi_{U_\eta^k} \rightarrow \chi_{\Omega_k}$ in $L^1(\mathbb{R}^3)$, we must have $|y_\eta^i - y_\eta^k| \rightarrow \infty$, that is,

$$\frac{|x_\eta^i - x_\eta^k|}{\eta} \rightarrow \infty, \quad i \neq k$$

Note that this countable process might stop after finitely many iterations: If $|\Omega_k| = M - \sum_{i=1}^{k-1} m^i$ after k iterations, then as in Step 2 we have $\chi_{\Omega_\eta^k} \rightarrow \chi_{\Omega_k}$ in $L^1(\mathbb{R}^3)$ and we may take $V_\eta^i, \Omega_i = \emptyset$ for all $i > k$. As we shall see in Step 7, this is indeed the case.

Step 5. The above process exhausts all the mass, and yields a lower bound on the energy. Indeed, passing to the $k \rightarrow \infty$ limit in (2.16) we observe that $\sum_{i=1}^\infty m^i < \infty$, and hence from (2.15) we must have $\mathcal{M}(\{\widehat{\Omega}_\eta^k\}) < 2m^k \rightarrow 0$ as $k \rightarrow \infty$. By Proposition 2.1 of [12] we may then conclude that $|\widehat{\Omega}_\eta^k| \rightarrow 0$, and so we have strong convergence,

$$\lim_{\eta \rightarrow 0} \left\| v_\eta - \sum_{i=1}^\infty w_\eta^i \right\|_{L^1(\mathbb{T}^3)} = \lim_{\eta \rightarrow 0} \left\| v_\eta^0 - \sum_{i=1}^\infty w_\eta^i \right\|_{L^1(\mathbb{T}^3)} = 0.$$

Finally, from (2.16) and (2.18) we may conclude

$$\mathcal{G}(\widehat{\Omega}_\eta) \geq \sum_{i=1}^\infty \mathcal{G}(\Omega_i) + o(1), \quad M = \sum_{i=1}^\infty m^i. \tag{2.20}$$

Using (2.11) and $\rho(x) \leq \rho_{\max}$, we may then conclude that

$$\mathbb{E}_\eta(v_\eta) \geq \mathbb{E}_\eta(v_\eta^0) + o(1) = \mathcal{G}(\widehat{\Omega}_\eta) - \int_{\mathbb{T}^3} v_\eta^0 \rho dx + o(1) \geq \sum_{i=1}^\infty \mathcal{G}(\Omega_i) - M\rho_{\max} + o(1). \tag{2.21}$$

Step 6. The next assertion is that $e_0(m^i)$ is attained for each $i = 1, 2, \dots$. Let Ω_i be as above. Then combining the lower bound (2.21) with (2.1) and the fact that $e_0(M) \leq \sum_{i=1}^\infty e_0(m^i)$ we obtain

$$\begin{aligned} \sum_{i=1}^\infty e_0(m^i) - M\rho_{\max} &\geq e_0(M) - M\rho_{\max} \geq \liminf_{\eta \rightarrow 0} \mathbb{E}_\eta(v_\eta) \\ &\geq \sum_{i=1}^\infty \mathcal{G}(\Omega_i) - M\rho_{\max} \geq \sum_{i=1}^\infty e_0(m^i) - M\rho_{\max}. \end{aligned} \tag{2.22}$$

Adding $M\rho_{\max}$ to each term, and noting that every term in these infinite sums are positive, we conclude that $e_0(m^i) = \mathcal{G}(\Omega_i)$; since $|\Omega_i| = m^i$, they attain the minimum value and each $\{m^i\}_{i \in \mathbb{N}} \in \mathcal{M}$. Moreover, we note that

$$e_0(M) = \sum_{i=1}^\infty e_0(m^i). \tag{2.23}$$

Step 7. There are only finitely many components. Let $m_{c_0} > 0$ be the constant given in Theorem 1.1. That is, e_0 is attained uniquely by a ball of volume m for

$m \leq m_{c_0}$. Then an explicit calculation shows that

$$e_0(m) = \left(3(4\pi/3)^{1/3}\right)m^{2/3} + \left((3/4\pi)^{5/3} \int_{B_1(0)} \int_{B_1(0)} \frac{1}{4\pi|x-y|} dx dy\right)m^{5/3}.$$

Computing the Coulomb term of a unit ball we see that $e_0''(m) < 0$ if $m < \widehat{m} := \min\{m_{c_0}, 2\pi\}$. Now suppose the first two terms m^1 and m^2 of the infinite sequence $\{m^i\}_{i=1}^\infty$ are in the interval $(0, \widehat{m})$. Then

$$\frac{d^2}{d\epsilon^2} \Big|_{\epsilon=0} \left(e_0(m^1 + \epsilon) + e_0(m^2 - \epsilon) \right) = e_0''(m^1) + e_0''(m^2) < 0.$$

However this contradicts the fact that the sequence $\{m^1, m^2, \dots\}$ is optimal for (2.6). Therefore there is at most one m^i on the interval $(0, \widehat{m})$. Since $\sum_{i=1}^\infty m^i = M < +\infty$ the number of nonzero m^i has to be finite.

This concludes the proof of Lemma 2.5. □

Remark 2.6. The same compactness result could be obtained within the context of Lions' concentration-compactness [20], without recourse to the results of [12]. Indeed, by defining a Lévy concentration function which is rescaled by η ,

$$Q_{v_\eta}(t) = \eta^3 \sup_{x \in \mathbb{T}^3} \int_{B_{\eta t}(x)} v_\eta(y) dy$$

we may proceed as in Lemma I.1 of [20]. However, Proposition 2.1 of [12] effectively eliminates the vanishing case, and the following Lemmas incorporate dichotomy and compactness in a very convenient form for BV spaces of characteristic functions.

Remark 2.7. We observe that $\{v_\eta\}$ need not be absolute minimizers of E_η : the same conclusions may be drawn for any sequence $v_\eta = \eta^{-3} \chi_{A_\eta}$ with $\lim_{\eta \rightarrow 0} \int_{\mathbb{T}^3} v_\eta = M$, and assuming an upper bound of the form,

$$\limsup_{\eta \rightarrow 0} E_\eta(v_\eta) \leq e_0(M) - M\rho_{\max}.$$

As a last remark, we note that thanks to Steps 6 and 7, we obtain more detailed information on the sets $U_\eta^i, V_\eta^i, i = 1, \dots, n$, which form the basis for the decomposition. This more refined description of these sets will be useful in proving regularity of the minimizing sequences.

Lemma 2.8. *Let U_η^i, V_η^i be as in the proof of Lemma 2.5. There exists $R > 0$, independent of η , such that:*

$$U_\eta^i \subset B_{2R}(0), \quad V_\eta^i \cap B_R = \emptyset, \quad \text{and} \quad \chi_{U_\eta^i \setminus B_{R/2}(0)} \rightarrow 0 \text{ in } L^1(\mathbb{R}^3).$$

Proof. By Step 6, Ω_i attains the minimum of $e_0(m^i)$, and hence is compact and connected. By Step 7, there are only finitely many, so we may choose $R > 0$ with $\Omega_i \subset B_{R/2}(0)$ for each $i = 1, \dots, n$. Returning to Steps 3 and 4, we may now choose the radii $r_{i,\eta} \in (R, 2R)$ (see the proof of [12, Lemma 2.3],) which yields the desired result. □

3. Regularity of minimizers of E_η . In determining the concentration structure of minimizers in Lemma 2.5 we obtain a rough lower bound on the energy with a possible error of $o(1)$, which is enough to identify each component as a minimizer of the liquid drop energy (at length scale η), but it not sharp enough to compute the interaction between droplets. We use the regularity theory of minimizers in order

to refine this decomposition, and in particular the lower bound inequality (2.7), in order to obtain an optimal lower bound with no error term to compete with the interdroplet interaction energy (which we will see later is of order $\mathcal{O}(\eta^{2/3})$.)

We start with a definition which gives a property of perimeter functionals that, as we will show, provides a strong regularity result for minimizers v_η .

Definition 3.1 (ω -minimality). Let $\mathcal{O} \subset \mathbb{R}^3$ be an open set, and $\omega > 0$. A set of finite perimeter $A \subset \mathbb{R}^3$ is an ω -minimizer for the perimeter functional $\int_{\mathbb{R}^3} |\nabla \chi_A|$ in \mathcal{O} if for any ball $B_r(x_0) \subset \mathcal{O}$ and any set of finite perimeter $B \subset \mathbb{R}^3$ such that $A \Delta B \subset \subset B_r(x_0)$ we have

$$\int_{\mathcal{O}} |\nabla \chi_A| \leq \int_{\mathcal{O}} |\nabla \chi_B| + \omega r^3.$$

Our goal in this section is to show that the minimizers v_η of E_η blow up to ω -minimizers of perimeter in \mathbb{R}^3 , with ω independent of η . To state this result we need to blow up the minimizing set $A_\eta = \text{supp } v_\eta$, truncated to a neighborhood of a point $p \in \mathbb{T}^3$, to a set in \mathbb{R}^3 ,

$$\tilde{A}_{p,\eta} := \Phi_{\eta, \frac{1}{4}}(A_\eta; p) = \eta^{-1} \left[(A_\eta - p) \cap B_{\frac{1}{4}}(0) \right],$$

so $\tilde{A}_{p,\eta} \subset \mathbb{R}^3$ and centered at the origin.

Lemma 3.2. *There exists $\omega > 0$, such that for any fixed $p \in \mathbb{T}^3$ and $R > 0$, $\tilde{A}_{p,\eta}$ is an ω -minimizer of perimeter in $B_R(0)$, uniformly for all $\eta \in (0, \frac{1}{2R})$.*

We recall that R is given in Lemma 2.8, such that the limit sets $\Omega_i \subset B_{R/2}$ for each i .

In proving Lemma 3.2 it is convenient to replace the mass constraint $\int_{\mathbb{T}^3} v_\eta = M$ with a penalization in the energy which effectively enforces this constraint, but allows us to choose comparison functions with arbitrary mass. Following [1], for any $\lambda > 0$ we define an unconstrained functional

$$E_\eta^\lambda(u) := E_\eta(u) + \lambda \left| \int_{\mathbb{T}^3} u dx - M \right|$$

over $u \in BV(\mathbb{T}^3; \{0, 1\})$. which penalizes deviations from the usual mass constraint. Then we have:

Lemma 3.3. *There exists constants $\eta_0, \lambda_0 > 0$ such that for every $0 < \eta < \eta_0$*

$$\inf \{ E_\eta^{\lambda_0}(u) : u \in BV(\mathbb{T}^3; \{0, 1\}) \} = E_\eta^{\lambda_0}(v_\eta) = E_\eta(v_\eta).$$

We first prove Lemma 3.2 assuming Lemma 3.3, whose proof follows immediately afterwards.

Proof of Lemma 3.2. Let $R > 0$ be fixed, and take any $r > 0$ with $B_r(x_0) \subset B_R(0)$, and $\tilde{S} \subset \mathbb{R}^3$ be any set satisfying $\tilde{S} \Delta \tilde{A}_{p,\eta} \subset B_r(x_0) \subset B_R(0)$. We now pull back the set \tilde{S} to \mathbb{T}^3 to construct a competitor $S_\eta \subset \mathbb{T}^3$ for the minimization problem for $E_\eta^{\lambda_0}$ as follows: inside $B_{\frac{1}{4}}(p)$, we take S_η with $S_\eta \cap B_{\frac{1}{4}}(p) = \Phi_{p,\eta}^{-1}(\tilde{S})$, and outside that neighborhood, $S_\eta \cap (B_{\frac{1}{4}}(p))^c = A_\eta$. Thus, S_η coincides with A_η except in the preimage $\Phi_{p,\eta}^{-1}(B_r(x_0)) \subset B_{\eta R}(p)$, that is

$$S_\eta \Delta A_\eta \subset \Phi_{p,\eta}^{-1}(B_r(x_0)) \subset B_{\eta R}(p),$$

and by rescaling,

$$|S_\eta \Delta A_\eta| = \eta^3 |\tilde{S} \Delta \tilde{A}_{p,\eta}|.$$

Define $\varphi_\eta := \eta^{-3} \chi_{S_\eta}$. By Lemma 3.3, $E_\eta(v_\eta) = E_\eta^{\lambda_0}(v_\eta) \leq E_\eta^{\lambda_0}(\varphi_\eta)$. Thus,

$$\begin{aligned} \eta \int_{\mathbb{T}^3} |\nabla v_\eta| &\leq \eta \int_{\mathbb{T}^3} |\nabla \varphi_\eta| + \eta \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} [\varphi_\eta(x)\varphi_\eta(y) - v_\eta(x)v_\eta(y)] G(x, y) dx dy \\ &\quad + \int_{\mathbb{T}^3} \rho(x) [v_\eta(x) - \varphi_\eta(x)] dx + \lambda_0 \left| \int_{\mathbb{T}^3} \phi_\eta - M \right|. \end{aligned} \quad (3.1)$$

We now estimate each term involving the difference $(\varphi_\eta - v_\eta)$, which we recall is supported in $B_{\eta R}(p)$. For the nonlocal term, we must split the integral by decomposing \mathbb{T}^3 into $B := B_{2\eta R}(p)$ and $B^c := \mathbb{T}^3 \setminus B_{2\eta R}(p)$. When both $x, y \in B$, we may decompose the Green's function as in (2.10), and so

$$\begin{aligned} &\eta \int_B \int_B [\varphi_\eta(x)\varphi_\eta(y) - v_\eta(x)v_\eta(y)] G(x, y) dx dy \\ &= \eta \int_B \int_B [\varphi_\eta(x)\varphi_\eta(y) - v_\eta(x)v_\eta(y)] \left(\frac{1}{4\pi|x-y|} + g(x, y) \right) dx dy \\ &\leq \int_{\tilde{S}} \int_{\tilde{S}} \frac{1}{4\pi|x-y|} dx dy - \int_{\tilde{A}_{p,\eta}} \int_{\tilde{A}_{p,\eta}} \frac{1}{4\pi|x-y|} dx dy + \eta^{-5} \int_B \int_{S_\eta \triangle A_\eta} |g(x, y)| dx dy \\ &\leq C_1 |\tilde{S} \triangle \tilde{A}_{p,\eta}| + C_2 \eta^{-5} |B_{2\eta R}(p)| |S_\eta \triangle A_\eta| \leq (C_1 + \mathcal{O}(\eta)) |\tilde{S} \triangle \tilde{A}_{p,\eta}|, \end{aligned}$$

where the difference of the Newtonian potential terms is bounded using [4, Proposition 2.3]. For $x \in B^c$ but $y \in B$ (or vice-versa,) we recall that $\varphi_\eta(x) = v_\eta(x)$, and $S_\eta \triangle A_\eta \subset B_{\eta R}(p)$, and thus $\text{dist}_{\mathbb{T}^3}(x, y) \geq R\eta$. Since $G(x, y)$ satisfies a uniform estimate,

$$|G(x, y)| \leq \frac{C}{\text{dist}_{\mathbb{T}^3}(x, y)} + C$$

with constant independent of $x \neq y \in \mathbb{T}^3$, we have

$$\begin{aligned} &\left| \eta \int_B \int_{B^c} [\varphi_\eta(x)\varphi_\eta(y) - v_\eta(x)v_\eta(y)] G(x, y) dx dy \right| \\ &= \left| \eta \int_B \int_{B^c} v_\eta(x) [\varphi_\eta(y) - v_\eta(y)] G(x, y) dx dy \right| \\ &\leq \eta^{-2} \int_{B^c} v_\eta(x) \left[\int_{S_\eta \triangle A_\eta} C \left[\frac{1}{\text{dist}_{\mathbb{T}^3}(x, y)} + 1 \right] dy \right] dx \\ &\leq C_3 \eta^{-3} |S_\eta \triangle A_\eta| = C_3 |\tilde{S} \triangle \tilde{A}_{p,\eta}|. \end{aligned}$$

Finally, if $x, y \in B^c$, then $[\varphi_\eta(x)\varphi_\eta(y) - v_\eta(x)v_\eta(y)] = 0$, so the remaining integral vanishes identically, and we obtain the estimate,

$$\eta \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} [\varphi_\eta(x)\varphi_\eta(y) - v_\eta(x)v_\eta(y)] G(x, y) dx dy \leq C_4 |\tilde{S} \triangle \tilde{A}_{p,\eta}|. \quad (3.2)$$

The remaining terms in the difference of the energies (3.1) are easier to bound. Indeed,

$$\begin{aligned} &\int_{\mathbb{T}^3} \rho(x) [v_\eta(x) - \varphi_\eta(x)] dx \leq \eta^{-3} \rho_{\max} |S_\eta \triangle A_\eta| = \rho_{\max} |\tilde{S} \triangle \tilde{A}_{p,\eta}|, \\ &\text{and } \lambda_0 \left| \int_{\mathbb{T}^3} \phi_\eta - M \right| = \lambda_0 \eta^{-3} ||S_\eta| - |A_\eta|| \leq \lambda_0 |\tilde{S} \triangle \tilde{A}_{p,\eta}|. \end{aligned}$$

Since $\tilde{S}\Delta\tilde{A}_{p,\eta} \subset B_r(x_0) \subset B_R(0) := \mathcal{O}$, from the above estimates applied to (3.1) we have:

$$\begin{aligned} \int_{\mathcal{O}} |\nabla\chi_{\tilde{A}_{p,\eta}}| - \int_{\mathcal{O}} |\nabla\chi_{\tilde{S}}| &= \eta \int_{B_{\eta R}(p)} |\nabla v_\eta| - \eta \int_{B_{\eta R}(p)} |\nabla\varphi_\eta| \\ &= \eta \int_{\mathbb{T}^3} (|\nabla v_\eta| - |\nabla\varphi_\eta|) \\ &\leq C_5 |\tilde{S}\Delta\tilde{A}_{p,\eta}| \leq \omega r^3, \end{aligned}$$

with constant $C_5 = (C_4 + \rho_{\max} + \lambda_0)$ and $\omega = C_5|B_1|$, independent of $\eta > 0$. \square

Proof of Lemma 3.3. We observe that by [1, Proposition 2.7], we may assume the existence of $\lambda = \lambda(\eta)$ for which the minimizer of the penalized problem coincides with the minimizer of the constrained problem, and so it suffices to show that $\lambda = \lambda_0$ may be chosen independently of η .

Suppose that the conclusion of the lemma is not true. That is, there exist sequences $\eta \rightarrow 0$, $\lambda_\eta \rightarrow \infty$, and $\varphi_\eta = \eta^{-3}\chi_{S_\eta}$ such that $\eta^{-3}|S_\eta| \neq M$ and

$$E_\eta^{\lambda_\eta}(\varphi_\eta) < E_\eta^{\lambda_\eta}(v_\eta) = E_\eta(v_\eta).$$

Since $\lambda_\eta \rightarrow \infty$, we have that $\eta^{-3}|S_\eta| \rightarrow M$. By the above upper bound and Remark 2.7, Lemma 2.5 applies to the family $\{\varphi_\eta\}$, and we return to the notation established there. By Step 1, we may decompose $S_\eta = \Omega_\eta \cup \Xi_\eta$ disjointly, with $\Omega_\eta \subset B_{3\delta}(0)$ and $\eta^{-3}|\Xi_\eta| \rightarrow 0$. By Step 3 there exist translations $x_\eta^1 = \eta y_\eta^1$ and disjoint sets U_η^1, V_η^1 so that after blowing up $\hat{\Omega}_\eta := \eta^{-1}(\Omega_\eta - x_\eta^1)$ to \mathbb{R}^3 ,

$$\hat{\Omega}_\eta = U_\eta^1 \cup V_\eta^1, \quad \chi_{U_\eta^1} \rightarrow \chi_{\Omega_1} \text{ in } L^1(\mathbb{R}^3), \quad \chi_{V_\eta^1} \rightarrow 0 \text{ in } L^1_{\text{loc}}(\mathbb{R}^3).$$

Moreover, $|U_\eta^1| \rightarrow m^1 = |\Omega_1|$. By Lemma 2.8 following the proof of Lemma 2.5, there exists $R > 0$ so that $\Omega_1 \subset\subset B_{R/2}(0)$, $U_\eta^1 \subset B_{2R}(0)$, $V_\eta^1 \cap B_R(0) = \emptyset$, and $\chi_{U_\eta^1 \setminus B_R(0)} \rightarrow 0$ in $L^1(\mathbb{R}^3)$.

Let $\mu_\eta := M - |V_\eta^1| - \eta^{-3}|\Xi_\eta|$. Then, $\mu_\eta = m_\eta^1 + o(1) = m^1 + o(1)$. The goal is to deform the sets U_η^1 via a family of dilations such that the resulting set \tilde{U}_η^1 has measure exactly equal to μ_η . Then, we may scale back down to \mathbb{T}^3 , substituting the new set $(\eta\tilde{U}_\eta^1 + x_\eta^1)$ inside $B_{2\eta R}(x_\eta^1)$, and obtain a set with mass exactly equal to $\eta^{-3}M$, and eventually obtain a contradiction. To that end, we define a smooth, non-increasing cut-off function of $r \geq 0$ with $0 \leq \phi(r) \leq 1$, $\phi(r) = 1$ for $0 \leq r \leq R/2$, and $\phi(r) = 0$ when $r \geq R$. Then, we let $\Psi_t : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the flow map associated to the dynamical system,

$$\mathbf{x}'(t) = \mathbf{F}(\mathbf{x}), \quad \mathbf{F}(\mathbf{x}) = \phi(|\mathbf{x}|)\mathbf{x}, \quad \mathbf{x}(0) = x.$$

The right-hand side is smooth with linear growth, and so Ψ_t maps $B_R(0)$ to itself for any $t \in \mathbb{R}$, and $\Psi_t(x) = x$ for all $x \notin B_R(0)$. For any set S of finite perimeter we define $S(t) := \Psi_t(S)$. Then, the measure $|S(t)|$ is a differentiable function of t , and

$$\frac{d}{dt}|S(t)| = \int_{S(t)} \operatorname{div}\mathbf{F}dx = \int_{S(t)} (3\phi(|x|) + \phi'(|x|)|x|) dx.$$

As $\Omega_1 \subset\subset B_{R/2}$, there exist constants $\epsilon, C > 0$ so that

$$\frac{d}{dt}|\Omega_1(t)| \geq 3|\Omega_1(t)| \geq 2C > 0, \quad \text{for all } t \in (-\epsilon, \epsilon).$$

Since $U_\eta^1 \rightarrow \Omega_1$ globally, we have $\frac{d}{dt}|U_\eta^1(t)| \geq C > 0$ for all $t \in (-\epsilon, \epsilon)$, for all $\eta > 0$ sufficiently small. Recalling that $|\mu_\eta - m_\eta^1| \rightarrow 0$, we may conclude that for each sufficiently small $\eta > 0$, there exists a unique $t_\eta \in (-\epsilon, \epsilon)$ so that $\tilde{U}_\eta^1 := U_\eta^1(t_\eta)$ has measure $|\tilde{U}_\eta^1| = \mu_\eta$. Moreover, $t_\eta = \mathcal{O}(\mu_\eta - m_\eta^1)$. Using the expansion of the Jacobian (see [23, Theorems 17.5,17.8],) we also have

$$\int_{\mathbb{R}^3} |\nabla \chi_{\tilde{U}_\eta^1}| \leq \int_{\mathbb{R}^3} |\nabla \chi_{U_\eta^1}| + Ct_\eta, \quad |\tilde{U}_\eta^1 \Delta U_\eta^1| \leq Ct_\eta, \quad (3.3)$$

with constant C depending on the perimeter of U_η^1 , but independent of η . Finally, we observe that since $\Psi_t(x) = x$ for $x \notin B_R(0)$, $\tilde{U}_\eta^1 \cap V_\eta^1 = \emptyset$.

Now we switch \tilde{U}_η^1 for U_η^1 and scale back down to \mathbb{T}^3 . Let \tilde{S}_η be the set which coincides with S_η in $\mathbb{T}^3 \setminus B_{3\delta}(0)$, and with $\Phi^{-1}(\tilde{U}_\eta^1 \cup V_\eta^1, y_\eta^1)$ in $B_{3\delta}(0)$, and $\tilde{\varphi}_\eta := \eta^{-3} \chi_{\tilde{S}_\eta}$. Then, by construction of \tilde{U}_η^1 , $\int_{\mathbb{T}^3} \tilde{\varphi}_\eta dx = M$ exactly. In addition, $(S_\eta \tilde{\Delta} \tilde{S}_\eta) \subset B_{2\eta R}(x_\eta^1)$, and so by the same calculation as (3.2) we have

$$\left| \eta \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} [\tilde{\varphi}_\eta(x) \tilde{\varphi}_\eta(y) - \varphi_\eta(x) \varphi_\eta(y)] G(x, y) dx dy \right| \leq C_4 |\tilde{U}_\eta^1 \Delta U_\eta^1| \leq C' t_\eta,$$

and

$$\left| \int_{\mathbb{T}^3} \rho(\tilde{\varphi}_\eta - \varphi_\eta) dx \right| \leq C' t_\eta,$$

for constant C' independent of η small, by the second estimate in (3.3). Lastly, by the first inequality in (3.3) we have

$$\left| \eta \int_{\mathbb{T}^3} (|\nabla \tilde{\varphi}_\eta| - |\nabla \varphi_\eta|) \right| \leq Ct_\eta.$$

Thus, for a constant C (independent of η ,) we have:

$$\begin{aligned} \mathbf{E}_\eta^{\lambda_\eta}(\tilde{\varphi}_\eta) - \mathbf{E}_\eta^{\lambda_\eta}(\varphi_\eta) &\leq Ct_\eta - \lambda_\eta \left| \int_{\mathbb{T}^3} \varphi_\eta dx - M \right| \\ &= Ct_\eta - \lambda_\eta (|U_\eta^1| - |\tilde{U}_\eta^1|) \\ &= Ct_\eta - \lambda_\eta (m_\eta^1 - \mu_\eta) < 0 \end{aligned}$$

for all sufficiently small $\eta > 0$, as $\lambda_\eta \rightarrow \infty$ and $t_\eta = \mathcal{O}(m_\eta^1 - \mu_\eta)$. This contradicts the choice of φ_η as the minimizer of $\mathbf{E}_\eta^{\lambda_\eta}$, and thus the lemma is proven. \square

The important consequence of ω -minimality is the following regularity result.

Proposition 3.4. *Let $\mathcal{O} \subset \mathbb{R}^3$ be a bounded open set, and $\hat{\Omega}_\eta \subset \mathcal{O}$ be a sequence of ω -minimizers of the perimeter functional such that*

$$\sup_{\eta > 0} \int_{\mathcal{O}} |\nabla \chi_{\hat{\Omega}_\eta}| < +\infty \quad \text{and} \quad \chi_{\hat{\Omega}_\eta} \rightarrow \chi_\Omega \quad \text{in } L^1(\mathcal{O})$$

for $\Omega \Subset \mathcal{O}$ of class C^2 . Then, for η small enough, $\hat{\Omega}_\eta$ is of class $C^{1,1/2}$ and

$$\partial \hat{\Omega}_\eta = \{x + \psi_\eta(x) \nu^\Omega(x) : x \in \partial \Omega\}$$

with $\psi_\eta \rightarrow 0$ in $C^{1,\alpha}(\partial \Omega)$ for all $\alpha \in (0, 1/2)$ where ν^Ω denotes the unit outward normal to $\partial \Omega$.

This result appears, for example, in [1, Theorem 4.2]. There it is stated on the torus; however, when the target set $\Omega \subset \mathbb{R}^3$ is bounded, using [23, Theorem 13.8], we see that the regularity theorem holds also in this context, with $\mathcal{O} = B_{2R}$ with R as in Lemma 2.8.

Applying the regularity theorem to minimizers of E_η we may conclude that minimizers split exactly and disjointly into the sets U_η^i found in Lemma 2.5:

Lemma 3.5. *Let $v_\eta = \eta^{-3}\chi_{A_\eta}$ be minimizers of E_η with $|A_\eta| = M\eta^3$. Using the notation of Lemma 2.5, let $n \in \mathbb{N}$, and u_η^i and $U_\eta^i \subset \mathbb{R}^3$, $i = 1, \dots, n$, as in (2.14). Then there exists $R > 0$ independent of η such that for all sufficiently small $\eta > 0$ we have:*

$$U_\eta^i \subset B_R(0), \quad \sum_{i=1}^n |U_\eta^i| = \sum_{i=1}^n m_\eta^i = M, \quad \text{and } v_\eta(x) = \sum_{i=1}^n u_\eta^i(x).$$

In addition, for sufficiently small $\eta > 0$,

$$E_\eta(v_\eta) \geq \sum_{i=1}^n E_\eta(u_\eta^i) + \sum_{\substack{i=1 \\ i \neq j}}^n \eta \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} \frac{u_\eta^i u_\eta^j}{4\pi|x-y|} dx dy + \mathcal{O}(\eta). \quad (3.4)$$

That is, v_η splits into exactly n well-separated components, the residual sets exactly vanish, that is $\Xi_\eta, V_\eta^n = \emptyset$ for small η , and we have a sharp lower bound on the energy.

Proof. Using the notation from the proof of Lemma 2.5, we have that (after blowing up,)

$$\widehat{\Omega}_\eta = \bigcup_{i=1}^n [U_\eta^i + y_\eta^i] \cup \widehat{\Omega}_\eta^n, \quad (3.5)$$

a disjoint union in \mathbb{R}^3 , with $|y_\eta^i - y_\eta^j| \rightarrow \infty$ for $i \neq j$. Moreover, each $\chi_{U_\eta^i} \rightarrow \chi_{\Omega_i}$ in $L^1(\mathbb{R}^3)$, with each Ω_i a minimizer of $e_0(m^i)$. By [4, Theorem 2.7], Ω_i is precompact and $\partial\Omega_i$ is of class $C^{3,\beta}$ for any $\beta < 1$. Thus, there exists $R > 0$ so that $\Omega_i \subset B_{R/2}(0)$ for all $i = 1, \dots, n$. From the proof of Lemma 2.2 of [12], the radii r_η in the definition of U_η^i, V_η^i may be chosen with $r_\eta \in (R, 2R)$. By Lemma 3.2, $\widehat{\Omega}_\eta$ are ω -minimizers of the perimeter in $B_{2R}(y_\eta^i)$, with ω independent of η , it follows from Proposition 3.4 that $\partial U_\eta^i \cap B_{2R} \subset [\partial\widehat{\Omega}_\eta - y_\eta^i] \cap B_{2R}(0)$ is a $C^{1,\alpha}$ graph converging to $\partial\Omega_i$ in $C^{1,\alpha}$ -norm. In particular, each $U_\eta^i \subset B_R(0)$ for all sufficiently small $\eta > 0$.

We next claim that $V_\eta^n = \emptyset$ for small enough η . Indeed, suppose this is not the case. Then there exists a subsequence $\eta \rightarrow 0$ (not relabelled) and point $z_\eta \in \mathbb{R}^3$ for which $z_\eta \in \partial V_\eta^n$. Consider the sequence of sets $\widetilde{V}_\eta := (V_\eta^n - z_\eta)$, which are uniformly ω -minimizers of perimeter in $B_1(0)$, by Lemma 3.2. By the compactness of ω -minimizing sequences, Proposition 21.13 of [23], there is a further subsequence along which $\widetilde{V}_\eta \rightarrow \widetilde{V}$ in $B_{\frac{1}{2}}(0)$, and by the regularity Theorem 21.14 of [23], \widetilde{V} is itself an ω -minimizer. Therefore $\partial\widetilde{V} \cap B_{\frac{1}{2}}(0)$ is a C^1 smooth surface, and by (i) of the same theorem $0 \in \partial\widetilde{V}$. However, $|\widetilde{V} \cap B_{\frac{1}{2}}(0)| = \lim_{\eta \rightarrow 0} |\widetilde{V}_\eta \cap B_{\frac{1}{2}}(0)| = 0$, which is impossible given the smoothness of $\widetilde{V} \cap B_{\frac{1}{2}}(0)$. Hence we conclude that $V_\eta^n = \emptyset$ for all small η .

Finally, we recall that $v_\eta = \eta^{-3}\chi_{A_\eta}$, with $A_\eta = \Omega_\eta \cup \Xi_\eta$ a disjoint union, $\Xi_\eta \cap B_{2\delta}(0) = \emptyset$, and $\eta^{-3}|\Xi_\eta| \leq \eta^{-3}|A_\eta \setminus B_\delta(0)| \rightarrow 0$. As for V_η^n , we also claim that $\Xi_\eta = \emptyset$ for all sufficiently small $\eta > 0$. Indeed, assume the contrary and so there

exists a subsequence (not relabelled) $\eta \rightarrow 0$ and points $x_\eta \in \partial \Xi_\eta \subset \mathbb{T}^3 \setminus B_{2\delta}(0)$. By compactness of $\mathbb{T}^3 \setminus B_\delta(0)$ we may extract a further subsequence and $x_0 \in \mathbb{T}^3 \setminus B_{2\delta}(0)$ so that $x_\eta \rightarrow x_0$. Consider $\Xi_\eta \cap B_\delta(x_0)$, and the blowup sets

$$\tilde{\Xi}_\eta := \Phi_{\eta,\delta}(A_\eta \setminus B_{\delta_\eta}(0); x_\eta) = \eta^{-1}[(\Xi_\eta - x_\eta) \cap B_\delta(0)] \subset \mathbb{R}^3.$$

As in the preceding paragraph, $\tilde{\Xi}_\eta$ are uniformly ω -minimizers of perimeter in $B_1(0)$, and (by the choice of x_η ,) $0 \in \partial \tilde{\Xi}_\eta$ for each η , and so (by the regularity theorem) the sets converge and zero lies in the boundary of the limiting set. Again, since $|\tilde{\Xi}_\eta| \rightarrow 0$, there is no limiting set, and by this contradiction we may conclude that $\hat{\Xi}_\eta = \emptyset$ for all small η .

Since $v_\eta(x) = \eta^{-3} \chi_{A_\eta}(x)$ and $A_\eta = \Omega_\eta \cup \Xi_\eta$, the final identity $v_\eta = \sum_{i=1}^n u_\eta^i$ is verified. As the supports of u_η^i are well separated, the perimeters split exactly,

$$\int_{\mathbb{T}^3} |\nabla v_\eta| = \sum_{i=1}^n \int_{\mathbb{T}^3} |\nabla u_\eta^i|.$$

The nonlocal term, being quadratic, decomposes into diagonal terms, which are included in $E_\eta(u_\eta^i)$, and off-diagonal terms,

$$\begin{aligned} & \eta \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} G(x, y) u_\eta^i(x) u_\eta^j(y) dx dy \\ &= \eta \int_{B_{3\delta}(0)} \int_{B_{3\delta}(0)} \left[\frac{1}{4\pi|x-y|} + g(x, y) \right] u_\eta^i(x) u_\eta^j(y) dx dy \\ &= \eta \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} \frac{u_\eta^i u_\eta^j}{4\pi|x-y|} dx dy + \mathcal{O}(\eta), \end{aligned}$$

for $i \neq j$. Finally, the confinement term is linear in the disjoint components, and so the lemma is proven. \square

4. Proof of the lower and upper bounds. Finally, in this section we use the sharp form of the decomposition to prove our main result by providing sharp upper and lower bounds on $E_\eta(v_\eta)$.

We begin by verifying the upper bound construction, which follows the heuristic argument given in the Introduction.

Proof of Lemma 2.4. Let $n \in \mathbb{N}$, points $p_1, \dots, p_n \in \mathbb{R}^3$, and $\{m^i\}_{i=1}^n \in \mathcal{M}$ be given. Since $e_0(m^i)$ is attained for each i , there exists z^i which attains the minimum in (1.5). Since the support of z^i are bounded, and we only consider a finite number of them, there exists a constant $r > 0$ for which $Z^i := \text{supp } z^i \subset B_r(0)$ for all $i = 1, \dots, n$. Let

$$\nu_\eta^i := \frac{1}{\eta^3} z^i \left(\frac{x - \eta^{1/3} p_i}{\eta} \right), \quad \text{and} \quad \nu_\eta := \sum_{i=1}^n \nu_\eta^i.$$

Since $\text{supp } \nu_\eta \subset B_{\frac{1}{4}}(0)$ for all sufficiently small $\eta > 0$, we also view ν_η as a function on \mathbb{T}^3 .

Then plugging ν_η into E_η we get that

$$\begin{aligned} & E_\eta(\nu_\eta) \\ &= \sum_{i=1}^n \eta \int_{\mathbb{T}^3} |\nabla \nu_\eta| + \eta \sum_{i,j=1}^n \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} G(x-y) \nu_\eta^i(x) \nu_\eta^j(y) dx dy - \sum_{i=1}^n \int_{\mathbb{T}^3} \nu_\eta^i(x) \rho(x) dx \end{aligned}$$

$$\begin{aligned}
 &= \sum_{i=1}^n e_0(m_i) - M\rho_{\max} + \eta \sum_{\substack{i,j=1 \\ i \neq j}}^n \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} \frac{\nu_\eta^i(x)\nu_\eta^j(y)}{4\pi|x-y|} dx dy \\
 &+ \sum_{i=1}^n \int_{\mathbb{T}^3} \nu_\eta^i(x)(\rho_{\max} - \rho(x)) dx + O(\eta), \tag{4.1}
 \end{aligned}$$

where we have used the representation of the Green’s function in terms of the Newtonian kernel and its regular part in $B_{\frac{1}{4}}(0)$, and the choice of z^i as minimizers of $e_0(m^i)$.

Taking each term separately, we evaluate for $i \neq j$, via the change of variables $\eta\xi = x - \eta^{1/3}p_i$, $\eta\zeta = y - \eta^{1/3}p_j$,

$$\begin{aligned}
 \eta \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} \frac{\nu_\eta^i(x)\nu_\eta^j(y)}{4\pi|x-y|} dx dy &= \eta^{2/3} \int_{Z^i} \int_{Z^j} \frac{1}{4\pi|\eta^{2/3}(\xi - \zeta) - (p_i - p_j)|} d\xi d\zeta \\
 &= \eta^{2/3} \frac{m^i m^j}{4\pi|p_i - p_j|} + o(\eta^{2/3}), \tag{4.2}
 \end{aligned}$$

by Dominated Convergence applied to the integral as $\eta \rightarrow 0$. And, in a similar way, using hypothesis (H3) on $\rho(x)$,

$$\begin{aligned}
 \int_{\mathbb{T}^3} \nu_\eta^i(x)(\rho_{\max} - \rho(x)) dx &= \int_{Z^i} q\left(\eta^{1/3}[p_i + \eta^{2/3}\xi]\right) d\xi + o(\eta^{2/3}) \\
 &= \eta^{2/3} \int_{Z^i} q(p_i + \eta^{2/3}\xi) d\xi + o(\eta^{2/3}) \\
 &= \eta^{2/3} m^i q(p_i) + o(\eta^{2/3}). \tag{4.3}
 \end{aligned}$$

Inserting (4.2), (4.3) into (4.1) yields the desired upper bound for $n \geq 2$.

For $M \in \mathcal{M}_0$, we take $n = 1$, $p_1 = 0$ and $z = \chi_Z$ which attains $e_0(M)$, and define $\nu_\eta(x) = z(x/\eta)$. Then, as in (4.1) we have

$$\begin{aligned}
 E_\eta(\nu_\eta) &= e_0(M) - M\rho_{\max} + \int_Z [\rho(\eta\xi) - \rho_{\max}] d\xi + O(\eta) \\
 &\leq e_0(M) - M\rho_{\max} + O(\eta),
 \end{aligned}$$

as the last integral is $O(\eta^2)$ (by hypothesis (H3),) but the principal error comes from the regular part of the Green’s function. \square

It remains to derive a matching sharp lower bound in order to complete the proof of the main result. We begin by showing that the droplet centers $\{x_\eta^i\}$ for a sequence of minimizers v_η converge to zero at a precise rate.

Lemma 4.1. *Let $n \in \mathbb{N}$ be given as in Lemma 2.5. If $n \geq 2$, then the points $\{x_\eta^i\}_{i=1}^n$ found in Lemma 2.5 satisfy*

$$x_\eta^i = \mathcal{O}(\eta^{1/3})$$

for each $i = 1, \dots, n$. If $n = 1$, then the points y_η from Step 1 in the proof of Lemma 2.5 satisfy

$$y_\eta = \mathcal{O}(\eta^{1/2}).$$

Proof. Let v_η be a minimizer of E_η with $\int_{\mathbb{T}^3} v_\eta dx = M$. Define

$$\lambda_\eta := \min_{i \neq j} |x_\eta^i - x_\eta^j| \quad \text{and} \quad \beta_\eta := \max_i |x_\eta^i| \tag{4.4}$$

over finite number of indices $i, j = 1, \dots, n$, and let $\{u_\eta^i\}_{i=1}^n$ with $v_\eta = \sum_{i=1}^n u_\eta^i$ be the functions found in Lemma 2.5. Since by their construction, $|x_\eta^i - x_\eta^j| \gg \eta$, it follows that $\beta_\eta \gg \eta$.

Step 1. Evaluating the nonlocal and confinement terms. We will require the following simple estimates for the lower and upper bounds on the energy as well. Assume $n \geq 2$. Then, since $U_\eta^i \subset B_R(0)$, for sufficiently small η we have

$$\left| \frac{1}{|x-y|} - \frac{1}{|x_\eta^i - x_\eta^j|} \right| \leq \frac{4R\eta}{|x_\eta^i - x_\eta^j|^2},$$

for all $x \in \text{supp } u_\eta^i = (\eta U_\eta^i + x_\eta^i)$ and $y \in \text{supp } u_\eta^j = (\eta U_\eta^j + x_\eta^j)$, $i \neq j$. Hence,

$$\int_{\mathbb{T}^3} \int_{\mathbb{T}^3} \frac{u_\eta^i u_\eta^j}{4\pi|x-y|} dx dy = \frac{m_\eta^i m_\eta^j}{4\pi|x_\eta^i - x_\eta^j|} + O(\eta|x_\eta^i - x_\eta^j|^{-2}) = \frac{m^i m^j}{4\pi|x_\eta^i - x_\eta^j|} (1 - o(1)). \quad (4.5)$$

To estimate the confinement term, choose k for which $|x_\eta^k| = \beta_\eta$. For $x \in (\eta U_\eta^k + x_\eta^k)$ we have $|x| \geq \frac{1}{2}\beta_\eta$ and hence using (H3) on the structure of $\rho(x)$ near zero, we may conclude the rough estimate:

$$\rho(x) - \rho_{\max} \leq -q(x) + o(|x|^2) \leq -c_2\beta_\eta^2,$$

for constant $c_2 > 0$ independent of η . This implies that

$$\int_{\mathbb{T}^3} u_\eta^k \rho(x) dx \leq m_\eta^k (\rho_{\max} - c_2\beta_\eta). \quad (4.6)$$

We thus have a rough lower bound on the energy,

$$\begin{aligned} E_\eta(v_\eta) &\geq \sum_{i=1}^n (e_0(m_\eta^i) - m_\eta^i \rho_{\max}) + \sum_{\substack{i,j=1 \\ i \neq j}}^n \eta \frac{m^i m^j}{4\pi|x_\eta^i - x_\eta^j|} (1 - o(1)) + c_1 m_\eta^k \beta_\eta^2 \\ &\geq e_0(M) - M\rho_{\max} + \sum_{\substack{i,j=1 \\ i \neq j}}^n \eta \frac{m^i m^j}{4\pi|x_\eta^i - x_\eta^j|} (1 - o(1)) + c_2 \beta_\eta^2, \end{aligned} \quad (4.7)$$

since $M = \sum_{i=1}^n m_\eta^i$, with $c_2 > 0$ independent of η .

Step 2. The scale of concentration, $n \geq 2$. Let $k \neq \ell \in \{1, \dots, n\}$ be such that (along some subsequence $\eta \rightarrow 0$) $|x_\eta^k| = \beta_\eta$, and $|x_\eta^k - x_\eta^\ell| = \lambda_\eta$. Matching the upper bound (2.2) to the lower bound (4.7),

$$\begin{aligned} e_0(M) - M\rho_{\max} + \eta^{2/3} \mu_0 &\geq \tilde{E}_\eta(v_\eta) \\ &\geq e_0(M) - M\rho_{\max} + \eta \frac{m_k m_\ell}{4\pi|x_\eta^k - x_\eta^\ell|} (1 - o(1)) + c_2 \beta_\eta^2 \\ &\geq e_0(M) - M\rho_{\max} + \eta \frac{c_1}{\lambda_\eta} + c_2 \beta_\eta^2, \end{aligned}$$

We may then conclude that

$$\lambda_\eta \geq C_1 \eta^{1/3} \quad \text{and} \quad \beta_\eta \leq C_2 \eta^{1/3}$$

for some constants $C_1, C_2 > 0$, as desired when $n \geq 2$.

Step 3. When $n = 1$, we have the simpler lower bound, since there are no interaction terms:

$$E_\eta(v_\eta) \geq e_0(M) - M\rho_{\max} + |y_\eta|^2 - o(\eta) = e_0(M) - M\rho_{\max} + \beta_\eta^2 - o(\eta) \quad (4.8)$$

Comparing the upper bound (2.3) from Lemma 2.4 with the lower bound (4.8), we have

$$e_0(M) - M\rho_{\max} + \beta_\eta^2 - \mathcal{O}(\eta) \leq \mathbb{E}_\eta(v_\eta) \leq e_0(M) - M\rho_{\max} + \mathcal{O}(\eta)$$

and hence $|y_\eta| = \beta_\eta \leq \mathcal{O}(\eta^{1/2})$ as demanded. \square

We are now ready to prove the main result stated in the Introduction.

Proof of Theorem 1.2. Let $v_\eta \in BV(\mathbb{T}^3; \{0, 1/\eta^3\})$ with $\int_{\mathbb{T}^3} v_\eta dx = M$ be a minimizer of \mathbb{E}_η for each $\eta > 0$. Applying Lemma 2.5, there exist $n \in \mathbb{N}$, $\{m^i\}_{i=1}^n \in \mathcal{M}$, and $\{x_\eta^i\}_{i=1}^n \subset \mathbb{T}^3$ satisfying the conditions given there. If $n = 1$, then conclusion (ii) in the statement of the theorem holds, by the second part of Lemma 4.1 and (2.8) from Lemma 2.5.

If instead $n \geq 2$, by the first part of Lemma 4.1 we have that $x_\eta^i = \mathcal{O}(\eta^{1/3})$ for every $i = 1, \dots, n$. That is, the sequences $\{\eta^{-1/3}x_\eta^i\}_{\eta>0} \subset \mathbb{T}^3$ are bounded for each $i = 1, \dots, n$. Passing to a subsequence let us define

$$x_i := \lim_{\eta \rightarrow 0} \eta^{-1/3}x_\eta^i \in \mathbb{T}^3$$

for each $i = 1, \dots, n$. Moreover, by Lemma 3.5, using a change of variables as in (4.2), we have the precise lower bound,

$$\begin{aligned} \mathbb{E}_\eta(v_\eta) &\geq \sum_{i=1}^n \mathbb{E}_\eta(u_\eta^i) + \sum_{\substack{i,j=1 \\ i \neq j}}^n \eta \int_{\mathbb{T}^3} \int_{\mathbb{T}^3} \frac{u_\eta^i u_\eta^j}{4\pi|x-y|} dx dy + \mathcal{O}(\eta) \\ &\geq \sum_{i=1}^n \left(\mathcal{G}(U_\eta^i) - \int_{U_\eta^i} \rho(x_\eta^i + \eta\xi) d\xi \right) \\ &\quad + \sum_{\substack{i,j=1 \\ i \neq j}}^n \eta \int_{U_\eta^i} \int_{U_\eta^j} \frac{1}{4\pi|(x_\eta^i - x_\eta^j) + \eta(\xi - \zeta)|} d\xi d\eta + \mathcal{O}(\eta) \\ &\geq \sum_{i=1}^n e_0(m_\eta^i) - M\rho_{\max} + \sum_{\substack{i,j=1 \\ i \neq j}}^n \eta \int_{U_\eta^i} \int_{U_\eta^j} \frac{1}{4\pi|(x_\eta^i - x_\eta^j) + \eta(\xi - \zeta)|} d\xi d\eta \\ &\quad + \sum_{i=1}^n \int_{U_\eta^i} [\rho_{\max} - \rho(x_\eta^i + \eta\xi)] d\xi + \mathcal{O}(\eta) \\ &\geq e_0(M) - M\rho_{\max} + \sum_{\substack{i,j=1 \\ i \neq j}}^n \eta^{2/3} \int_{U_\eta^i} \int_{U_\eta^j} \frac{1}{4\pi|\eta^{-1/3}(x_\eta^i - x_\eta^j) + \eta^{2/3}(\xi - \zeta)|} d\xi d\eta \\ &\quad + \eta^{2/3} \sum_{i=1}^n \int_{U_\eta^i} q(\eta^{-1/3}x_\eta^i + \eta^{2/3}\xi) d\xi + o(\eta^{2/3}) \end{aligned}$$

Now, since each $\eta^{-1/3}x_\eta^i \rightarrow x_i$, and the $U_\eta^i \rightarrow \Omega_i$ globally, and all sets are compact in \mathbb{R}^3 , by Dominated Convergence we may conclude that

$$\lim_{\eta \rightarrow 0} \int_{U_\eta^i} \int_{U_\eta^j} \frac{1}{4\pi|\eta^{-1/3}(x_\eta^i - x_\eta^j) + \eta^{2/3}(\xi - \zeta)|} d\xi d\eta = \frac{m^i m^j}{4\pi|x_i - x_j|},$$

and

$$\lim_{\eta \rightarrow 0} \int_{U_\eta^i} q(\eta^{-1/3}x_\eta^i + \eta^{2/3}\xi) d\xi = m^i q(x_i),$$

and arrive at the desired lower bound,

$$E_\eta(v_\eta) \geq e_0(M) - \rho_{\max}M + \eta^{2/3} \left\{ \sum_{\substack{i,j=1 \\ i \neq j}}^n \frac{m^i m^j}{4\pi|x_i - x_j|} + \sum_{i=1}^n m^i q(x_i) \right\} + o(\eta^{2/3}).$$

Matching this lower bound with the upper bound from Lemma 2.4, the two match up to order $o(\eta^{2/3})$, with the $\{m^i\}$ and $\{x_j\}$ as selected by the concentration lemma. Since the upper bound holds for any choice of masses and rescaled centers, it holds for those which minimize the quantity F_{m^1, \dots, m^n} , and so those must be the values which appear in a minimizing sequence for E_η . Thus, the conclusion (iii) holds whenever $n \geq 2$ in Lemma 2.5.

It remains to connect the number n of components with the question of whether $M \in \mathcal{M}_0$ or not. First, if $M \notin \mathcal{M}_0$, then $n \geq 2$ by the last assertion in the statement of Lemma 2.5. To complete the argument, assume $M \in \mathcal{M}_0$ and $n \geq 2$. But then, Lemma 2.4 provides a stronger upper bound which doesn't match with the lower bound for $n \geq 2$,

$$e_0(M) - M\rho_{\max} + \mathcal{O}(\eta) \geq E_\eta(v_\eta) \geq e_0(M) - M\rho_{\max} + \eta^{2/3}F_{m^1, \dots, m^n} + o(\eta^{2/3}).$$

This can only be true if $F_{m^1, \dots, m^n} = 0$, which is impossible. Thus we must have (i) holding for $M \in \mathcal{M}_0$, by (2.8) in Lemma 2.5. \square

5. Penalization measures with degenerate maxima. Finally, in this section we consider the situation where the penalization measure defined via the density function ρ is degenerate at its maximum value, in the sense that the Hessian degenerates at a single maximum point. Note that by Remark 2.2 the first-order approximation still holds in such cases. It is at the second order that the behavior will differ, as the interaction energy will depend on the structure of ρ at its maximum.

The case where ρ attains its maximum at a unique point (say at $x = 0$), but with vanishing Hessian, does not entail much variation from the arguments used in the nondegenerate case above. Indeed, the concentration structure given in Lemmas 2.5 and 3.5 does not rely on the nondegeneracy hypothesis (H3), which only makes itself felt in the scaling arguments which determine the interdroplet distance in the upper bound construction of Lemma 2.4 and in the corresponding lower bound estimates of Lemma 4.1. In particular, we may replace the nondegeneracy condition (H3) by a homogeneity assumption of the form,

(H3') There exist $q > 2$ and $r, \rho_1 > 0$ so that $\rho \in C^1(B_r(0))$ and

$$\rho(x) := \rho(0) - \rho_1|x|^q + o(|x|^q), \quad \text{as } x \rightarrow 0,$$

without changing the concentration structure of minimizers. However, this choice will alter the geometry of minimizing droplet configurations: the appropriate secondary scale for pattern formation occurs at distance of $\mathcal{O}(\eta^{1/(q+1)})$, and the energy scale for interaction between droplets will be of order $\mathcal{O}(\eta^{q/(q+1)})$. That is, minimizers v_η will have an energy expansion of the form

$$E_\eta(v_\eta) = e_0(M) - M\rho_{\max} + \eta^{\frac{q}{q+1}}F_0(v) + o\left(\eta^{\frac{q}{q+1}}\right),$$

with

$$F_{m^1, \dots, m^n}(v) := \begin{cases} \rho_1 \sum_{i=1}^n m^i |x_i|^q + \frac{1}{4\pi} \sum_{\substack{i, j=1 \\ i \neq j}}^n \frac{m^i m^j}{|x_i - x_j|} \\ \quad \text{if } v = \sum_{i=1}^n m^i \delta_{x_i} \text{ with } \{x_i\} \text{ distinct, } \{m^i\} \in \mathcal{M}, \\ +\infty \quad \text{otherwise} \end{cases}$$

and $\{m^i\} \in \mathcal{M}$, $\{x_i\}$ chosen to minimize $F_{m^1, \dots, m^n}(v)$. The minimizing droplet configurations for examples with $q = 2, 4, 10$ are shown in Figure 2.

6. Appendix. Here we briefly discuss the physics behind diblock copolymers, the role of nanoparticle attraction in confinement, and the droplet scaling regime.

Diblock copolymers and confinement. Diblock copolymers are macromolecules composed of two chemically distinct homogeneous polymer chains (of monomer species A and B respectively) linked together by a covalent bond (cf. [3, 13]). The thermodynamical incompatibility between the different sub-chains drives the system to phase separate; however, the covalent bonds between the different sub-chains prevent phase separation at a macroscopic length scale. As a result of these two competing trends, block copolymers undergo phase separation at a nanometer length scale, leading to an amazingly rich array of nanostructures. Perhaps the simplest model for self-assembly of diblock copolymers is via the *Ohta–Kawasaki functional* [26], a phase-field model (diffuse interface) with nonlocal interactions. Recently, there has been considerable interest in the physics and engineering communities on self-assembly of diblock copolymers under confinement (see for example [28] and the references therein). Of particular interest is the situation where the length of the confinement region is comparable with the length of the diblock copolymer macromolecules.

Recently three of the authors (see [2]) introduced and analyzed a simple model for a polymer/nanoparticle blend, wherein the nanoparticles were attracted to one of the two molecular species in the polymers. Copolymer/nanoparticle interaction is modeled in the Ohta–Kawasaki functional by the addition of a penalization term which prefers the $u = 1$ phase in the neighborhood of each nanoparticle [14]. Assuming a fixed limiting distribution of a large number of nanoparticles occupying a vanishingly small volume of the sample, and passing to the sharp interface limit, [2] obtain a model for diblock copolymers under confinement of the form,

$$\mathcal{E}_{\gamma, \sigma}(u) = \int_{\mathbb{T}^3} |\nabla u| + \gamma \|u - m\|_{H^{-1}(\mathbb{T}^3)}^2 + \sigma \int_{\mathbb{T}^3} (1 - u(x)) \rho(x) dx, \quad \int_{\mathbb{T}^3} u dx = m.$$

For relatively large $\sigma > 0$, the region of high nanoparticle density effectively confines the $\{u = 1\}$ phase to this region, and we may simply view the penalization term in the energy as a confinement term subject to the density ρ . Indeed, an example in [2] illustrates how confinement modifies minimizing patterns in the isoperimetric problem ($\sigma = 0$), from lamellar to a sphere, with increasing σ .

The Droplet scaling regime. We arrive at the desired singularly perturbed energy functional (1.1) by extending the droplet scaling of [8] to encompass the confinement term in (6.1). First, we define an auxiliary shifted energy by

$$\begin{aligned} E_{\gamma,\sigma}(u) &:= \mathcal{E}_{\gamma,\sigma}(u) - \sigma \int_{\mathbb{T}^3} \rho(x) dx \\ &= \int_{\mathbb{T}^3} |\nabla u| + \gamma \|u - m\|_{H^{-1}(\mathbb{T}^3)}^2 - \sigma \int_{\mathbb{T}^3} u(x) \rho(x) dx. \end{aligned}$$

We now introduce the master parameter $0 < \eta \ll 1$, intended to represent the approximate diameter of the droplets of the phase $\{u = 1\}$. The droplet volume will tend to zero, so we need to adjust the order parameter to produce particles of fixed mass, $M > 0$. Thus, we introduce a concentrating order parameter $v_\eta : \mathbb{T}^3 \rightarrow \{0, 1/\eta^3\}$ via

$$v_\eta(x) := \eta^{-3} u(x), \quad \text{so} \quad \int_{\mathbb{T}^3} v_\eta dx = \eta^{-3} m := M.$$

We note that the support of v_η is the same as that of u . We then have

$$E_{\gamma,\sigma}(u) = \eta^2 \left[\eta \int_{\mathbb{T}^3} |\nabla v_\eta| + \gamma \eta^4 \|v_\eta - M\|_{H^{-1}(\mathbb{T}^3)}^2 - \sigma \eta \int_{\mathbb{T}^3} v_\eta \rho(x) dx \right]. \quad (6.1)$$

Given the anticipated scaling for v_η as suggested by the ansatz (1.3), we expect each term in the brackets to be of order one, and thus we define the droplet regime by the following choice of the parameters,

$$\gamma = \frac{1}{\eta^3} \quad \text{and} \quad \sigma = \frac{1}{\eta}.$$

Thus, we define the rescaled energy as follows:

$$\begin{aligned} E_\eta(v) &:= \frac{1}{\eta^2} E_{\gamma,\sigma}(u) = \frac{1}{\eta^2} \left(\mathcal{E}_{\gamma,\sigma}(u) - \sigma \int_{\mathbb{T}^3} \rho \right) \\ &= \eta \int_{\mathbb{T}^3} |\nabla v| + \eta \|v - M\|_{H^{-1}(\mathbb{T}^3)}^2 - \int_{\mathbb{T}^3} v(x) \rho(x) dx. \end{aligned} \quad (6.2)$$

Acknowledgements. The authors were supported by NSERC (Canada) Discovery Grants. IT was also supported by a Field–Ontario Postdoctoral Fellowship. We are also grateful to An-Chang Shi for pointing out that the functional used in [2] can also be used as a model problem for diblock copolymers under confinement.

REFERENCES

- [1] Emilio Acerbi, Nicola Fusco and Massimiliano Morini, [Minimality via second variation for a nonlocal isoperimetric problem](#), *Comm. Math. Phys.*, **322** (2013), 515–557.
- [2] Stan Alama, Lia Bronsard and Ihsan Topaloglu, [Sharp interface limit of an energy modelling nanoparticle-polymer blends](#), *Interfaces Free Bound.*, **18** (2016), 263–290.
- [3] Frank S. Bates and Glenn H. Fredrickson, Block copolymers—designer soft materials, *Physics Today*, **52** (1999), 32–38.
- [4] Marco Bonacini and Riccardo Cristoferi, [Local and global minimality results for a nonlocal isoperimetric problem on \$\mathbb{R}^N\$](#) , *SIAM J. Math. Anal.*, **46** (2014), 2310–2349.
- [5] Almut Burchard, Rustum Choksi and Ihsan Topaloglu, [Nonlocal shape optimization via interactions of attractive and repulsive potentials](#), *Indiana Univ. Math. J.*, **67** (2018), 375–395.
- [6] Djalil Chafaï, Nathael Gozlan and Pierre-André Zitt, [First-order global asymptotics for confined particles with singular pair repulsion](#), *Ann. Appl. Probab.*, **24** (2014), 2371–2413.
- [7] Rustum Choksi, Cyril B. Muratov and Ihsan Topaloglu, An old problem resurfaces nonlocally: Gamow’s liquid drops inspire today’s research and applications, *Notices Amer. Math. Soc.*, **64** (2017), 1275–1283.

- [8] Rustum Choksi and Mark A. Peletier, [Small volume fraction limit of the diblock copolymer problem: I. Sharp-interface functional](#), *SIAM J. Math. Anal.*, **42** (2010), 1334–1370.
- [9] Yao-Li Chuang, Maria R. D’Orsogna, Daniel Marthaler, Andrea L. Bertozzi, and Lincoln S. Chayes, [State transitions and the continuum limit for a 2D interacting, self-propelled particle system](#), *Phys. D*, **232** (2007), 33–47.
- [10] Rupert Frank, Rowan Killip and Phan Thành Nam, [Nonexistence of large nuclei in the liquid drop model](#), *Lett. Math. Phys.*, **106** (2016), 1033–1036.
- [11] Rupert L. Frank and Elliot H. Lieb, [A “liquid-solid” phase transition in a simple model for swarming, based on the “no flat-spots” theorem for subharmonic functions](#), *Indiana Univ. Math. J.*, **67** (2018), 1547–1569.
- [12] Rupert L. Frank and Elliott H. Lieb, [A compactness lemma and its application to the existence of minimizers for the liquid drop model](#), *SIAM J. Math. Anal.*, **47** (2015), 4436–4450.
- [13] Glenn Fredrickson, *Equilibrium Theory of Inhomogeneous Polymers*, Oxford Science Publications, 2005.
- [14] Valeriy V. Ginzburg, Feng Qiu, Marco Paniconi, Gongwen Peng, David Jasnow and Anna C Balazs, [Simulation of hard particles in a phase-separating binary mixture](#), *Phys. Rev. Lett.*, **82** (1999), 4026–4029.
- [15] Shay Gueron and Itai Shafir, [On a discrete variational problem involving interacting particles](#), *SIAM J. Appl. Math.*, **60** (2000), 1–17 (electronic).
- [16] Vesa Julin, [Isoperimetric problem with a Coulomb repulsive term](#), *Indiana Univ. Math. J.*, **63** (2014), 77–89.
- [17] Hans Knüpfer and Cyrill B. Muratov, [On an isoperimetric problem with a competing nonlocal term I: The planar case](#), *Comm. Pure Appl. Math.*, **66** (2013), 1129–1162.
- [18] Hans Knüpfer and Cyrill B. Muratov, [On an isoperimetric problem with a competing nonlocal term II: The general case](#), *Comm. Pure Appl. Math.*, **67** (2014), 1974–1994.
- [19] Hans Knüpfer, Cyrill B. Muratov and Matteo Novaga, [Low density phases in a uniformly charged liquid](#), *Comm. Math. Phys.*, **345** (2016), 141–183.
- [20] Pierre-Louis Lions, [The concentration-compactness principle in the calculus of variations. The locally compact case. I](#), *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **1** (1984), 109–145.
- [21] Jiangfeng Lu and Felix Otto, [Nonexistence of a minimizer for Thomas-Fermi-Dirac-von Weizsäcker model](#), *Comm. Pure Appl. Math.*, **67** (2014), 1605–1617.
- [22] Jiangfeng Lu and Felix Otto, [An isoperimetric problem with Coulomb repulsion and attraction to a background nucleus](#), [arXiv:1508.07172](#), 2015.
- [23] Francesco Maggi, *Sets of Finite Perimeter and Geometric Variational Problems*, volume 135 of Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, first edition, 2012.
- [24] Daniela Morale, Vincenzo Capasso and Karl Oelschläger, [An interacting particle system modeling aggregation behavior: from individuals to populations](#), *J. Math. Biol.*, **50** (2005), 49–66.
- [25] Phan Thành Nam and Hanne van den Bosch, [Nonexistence in Thomas-Fermi-Dirac-von Weizsäcker theory with small nuclear charges](#), *Math. Phys. Anal. Geom.*, **20** (2017), Art. 6, 1–32.
- [26] Takao Ohta and Kyozi Kawasaki, [Equilibrium morphology of block copolymer melts](#), *Macromolecules*, **19** (1986), 2621–2632.
- [27] Etienne Sandier and Sylvia Serfaty, *Vortices in The Magnetic Ginzburg-Landau Model*, Progress in Nonlinear Differential Equations and their Applications, 70. Birkhäuser Boston, Inc., Boston, MA, 2007.
- [28] An-Chang Shi and Baohui Li, [Self-assembly of diblock copolymers under confinement](#), *Soft Matter*, **9** (2013), 1398–1413.
- [29] James H. von Brecht, David Uminsky, Theodore Kolokolnikov and Andrea L. Bertozzi, [Predicting pattern formation in particle interactions](#), *Math. Models Methods Appl. Sci.*, **22**(suppl. 1) (2012), 1140002, 1–31.

Received August 2018; revised March 2019.

E-mail address: alama@mcmaster.ca

E-mail address: iatopaloglu@vcu.edu

E-mail address: bronsard@mcmaster.ca

E-mail address: rustum.choksi@mcgill.ca