

Problem Solving Seminar. Fall 2022. Problem Set 9. Miscellaneous.

1. **Putnam 2018. A1.** Find all ordered pairs  $(a, b)$  of positive integers for which

$$\frac{1}{a} + \frac{1}{b} = \frac{3}{2018}.$$

2. **Putnam 2018. A2.** Let  $S_1, S_2, \dots, S_{2^n-1}$  be the nonempty subsets of  $\{1, 2, \dots, n\}$  in some order, and let  $M$  be the  $(2^n - 1) \times (2^n - 1)$  matrix whose  $(i, j)$  entry is

$$m_{ij} = \begin{cases} 0 & \text{if } S_i \cap S_j = \emptyset; \\ 1 & \text{otherwise.} \end{cases}$$

Calculate the determinant of  $M$ .

3. **Putnam 2018. A3.** Determine the greatest possible value of  $\sum_{i=1}^{10} \cos(3x_i)$  for real numbers  $x_1, x_2, \dots, x_{10}$  satisfying  $\sum_{i=1}^{10} \cos(x_i) = 0$ .
4. **Putnam 2013. A1.** Recall that a regular icosahedron is a convex polyhedron having 12 vertices and 20 faces; the faces are congruent equilateral triangles. On each face of a regular icosahedron is written a nonnegative integer such that the sum of all 20 integers is 39. Show that there are two faces that share a vertex and have the same integer written on them.
5. **Putnam 2013. A2.** Let  $S$  be the set of all positive integers that are *not* perfect squares. For  $n$  in  $S$ , consider choices of integers  $a_1, a_2, \dots, a_r$  such that  $n < a_1 < a_2 < \dots < a_r$  and  $n \cdot a_1 \cdot a_2 \cdots a_r$  is a perfect square, and let  $f(n)$  be the minimum of  $a_r$  over all such choices. For example,  $2 \cdot 3 \cdot 6$  is a perfect square, while  $2 \cdot 3, 2 \cdot 4, 2 \cdot 5, 2 \cdot 3 \cdot 4, 2 \cdot 3 \cdot 5, 2 \cdot 4 \cdot 5,$  and  $2 \cdot 3 \cdot 4 \cdot 5$  are not, and so  $f(2) = 6$ . Show that the function  $f$  from  $S$  to the integers is one-to-one.
6. **Putnam 2017. B1.** Let  $L_1$  and  $L_2$  be distinct lines in the plane. Prove that  $L_1$  and  $L_2$  intersect if and only if, for every real number  $\lambda \neq 0$  and every point  $P$  not on  $L_1$  or  $L_2$ , there exist points  $A_1$  on  $L_1$  and  $A_2$  on  $L_2$  such that  $\overrightarrow{PA_2} = \lambda \overrightarrow{PA_1}$ .
7. **Putnam 2017. B2.** Suppose that a positive integer  $N$  can be expressed as the sum of  $k$  consecutive positive integers

$$N = a + (a + 1) + (a + 2) + \cdots + (a + k - 1)$$

for  $k = 2017$  but for no other values of  $k > 1$ . Considering all positive integers  $N$  with this property, what is the smallest positive integer  $a$  that occurs in any of these expressions?

8. **Putnam 2017. B3.** Suppose that  $f(x) = \sum_{i=0}^{\infty} c_i x^i$  is a power series for which each coefficient  $c_i$  is 0 or 1. Show that if  $f(2/3) = 3/2$ , then  $f(1/2)$  must be irrational.