

NON-PARAMETRIC STATISTICS

THE ROLE OF RANDOMIZATION/PERMUTATION TESTS

Randomization or **Permutation** procedures are useful for computing **exact** null distributions for non-parametric test statistics when sample sizes are small.

We focus first on two sample comparisons; suppose that two data samples $x_1 \dots, x_{n_1}$ and $y_1 \dots, y_{n_2}$ (where $n_1 \geq n_2$) have been obtained, and we wish to carry out a comparison of the two populations from which the samples are drawn. The Wilcoxon test statistic, W , is the sum of the ranks for the second sample. The permutation test proceeds as follows:

1. Let $n = n_1 + n_2$. Assuming that there are no ties, the pooled and ranked samples will have ranks

$$1 \quad 2 \quad 3 \quad \dots \quad n$$

2. The test statistic is $W = R_2$, the rank sum for sample two items. For the observed data, W will be the sum of n_2 of the ranks given in the list above.
3. If the null hypothesis

$$H_0 : \text{No difference between population 1 and population 2}$$

were **true**, then we would expect **no pattern** in the arrangements of the group labels when sorted into ascending order. That is, the sorted data would give rise a **random** assortment of group 1 and group 2 labels.

4. To obtain the exact distribution of W under H_0 (which is what we require for the assessment of statistical significance), we could compute W for all possible permutations of the group labels, and then form the probability distribution of the values of W . We call this the **permutation null distribution**.
5. But W is a rank sum, so we can compute the permutation null distribution simply by tabulating **all possible subsets** of size n_2 of the set of ranks $\{1, 2, 3, \dots, n\}$.

6. There are

$$\binom{n}{n_2} = \frac{n!}{n_1! n_2!} = N$$

say possible subsets of size n_2 . For example, for $n = 6$ and $n_2 = 2$, the number of subsets of size n_2 is

$$\binom{6}{2} = \frac{6!}{4! 2!} = 15$$

However, the number of subsets increases dramatically as n increases; for $n_1 = n_2 = 10$, so that $n = 20$, the number of subsets of size n_2 is

$$\binom{20}{10} = \frac{20!}{10! 10!} = 184756$$

7. The exact rejection region and p -value are computed from the permutation null distribution. Let $W_i, i = 1, \dots, N$ denote the value of the Wilcoxon statistic for the N possible subsets of the ranks of size n_2 . The probability that the test statistic, W , is less than or equal to w is

$$\Pr[W \leq w] = \frac{\text{Number of } W_i \leq w}{N}$$

We seek the values of w that give the appropriate rejection region, \mathcal{R} , so that

$$\Pr[W \in \mathcal{R}] = \frac{\text{Number of } W_i \in \mathcal{R}}{N} = \alpha$$

It may not be possible to find critical values, and define \mathcal{R} , so that this probability is **exactly** α as the distribution of W is **discrete**.

EXAMPLE : Simple Example

Suppose $n_1 = 7$ and $n_2 = 3$. There are

$$\binom{10}{3} = \frac{10!}{7!3!} = 120$$

subsets of the ranks $\{1, 2, 3, \dots, 10\}$ of size 3. The subsets are listed below, together with the rank sums.

| Ranks | W | Ranks | W | Ranks | W | Ranks | W |
|--------|----|--------|----|--------|----|--------|----|
| 1 2 3 | 6 | 1 7 8 | 16 | 2 7 10 | 19 | 4 6 7 | 17 |
| 1 2 4 | 7 | 1 7 9 | 17 | 2 8 9 | 19 | 4 6 8 | 18 |
| 1 2 5 | 8 | 1 7 10 | 18 | 2 8 10 | 20 | 4 6 9 | 19 |
| 1 2 6 | 9 | 1 8 9 | 18 | 2 9 10 | 21 | 4 6 10 | 20 |
| 1 2 7 | 10 | 1 8 10 | 19 | 3 4 5 | 12 | 4 7 8 | 19 |
| 1 2 8 | 11 | 1 9 10 | 20 | 3 4 6 | 13 | 4 7 9 | 20 |
| 1 2 9 | 12 | 2 3 4 | 9 | 3 4 7 | 14 | 4 7 10 | 21 |
| 1 2 10 | 13 | 2 3 5 | 10 | 3 4 8 | 15 | 4 8 9 | 21 |
| 1 3 4 | 8 | 2 3 6 | 11 | 3 4 9 | 16 | 4 8 10 | 22 |
| 1 3 5 | 9 | 2 3 7 | 12 | 3 4 10 | 17 | 4 9 10 | 23 |
| 1 3 6 | 10 | 2 3 8 | 13 | 3 5 6 | 14 | 5 6 7 | 18 |
| 1 3 7 | 11 | 2 3 9 | 14 | 3 5 7 | 15 | 5 6 8 | 19 |
| 1 3 8 | 12 | 2 3 10 | 15 | 3 5 8 | 16 | 5 6 9 | 20 |
| 1 3 9 | 13 | 2 4 5 | 11 | 3 5 9 | 17 | 5 6 10 | 21 |
| 1 3 10 | 14 | 2 4 6 | 12 | 3 5 10 | 18 | 5 7 8 | 20 |
| 1 4 5 | 10 | 2 4 7 | 13 | 3 6 7 | 16 | 5 7 9 | 21 |
| 1 4 6 | 11 | 2 4 8 | 14 | 3 6 8 | 17 | 5 7 10 | 22 |
| 1 4 7 | 12 | 2 4 9 | 15 | 3 6 9 | 18 | 5 8 9 | 22 |
| 1 4 8 | 13 | 2 4 10 | 16 | 3 6 10 | 19 | 5 8 10 | 23 |
| 1 4 9 | 14 | 2 5 6 | 13 | 3 7 8 | 18 | 5 9 10 | 24 |
| 1 4 10 | 15 | 2 5 7 | 14 | 3 7 9 | 19 | 6 7 8 | 21 |
| 1 5 6 | 12 | 2 5 8 | 15 | 3 7 10 | 20 | 6 7 9 | 22 |
| 1 5 7 | 13 | 2 5 9 | 16 | 3 8 9 | 20 | 6 7 10 | 23 |
| 1 5 8 | 14 | 2 5 10 | 17 | 3 8 10 | 21 | 6 8 9 | 23 |
| 1 5 9 | 15 | 2 6 7 | 15 | 3 9 10 | 22 | 6 8 10 | 24 |
| 1 5 10 | 16 | 2 6 8 | 16 | 4 5 6 | 15 | 6 9 10 | 25 |
| 1 6 7 | 14 | 2 6 9 | 17 | 4 5 7 | 16 | 7 8 9 | 24 |
| 1 6 8 | 15 | 2 6 10 | 18 | 4 5 8 | 17 | 7 8 10 | 25 |
| 1 6 9 | 16 | 2 7 8 | 17 | 4 5 9 | 18 | 7 9 10 | 26 |
| 1 6 10 | 17 | 2 7 9 | 18 | 4 5 10 | 19 | 8 9 10 | 27 |

There are 22 possible rank sums, $\{6, 7, 8, \dots, 25, 26, 27\}$; the number of times each is observed is displayed in the table below, with the corresponding probabilities and cumulative probabilities.

| | | | | | | | | | | | |
|------------------|-------|-------|--------------|-------|-------|-------|-------|-------|-------|-------|-------|
| W | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 |
| Frequency | 1 | 1 | 2 | 3 | 4 | 5 | 7 | 8 | 9 | 10 | 10 |
| Prob. | 0.008 | 0.008 | 0.017 | 0.025 | 0.033 | 0.042 | 0.058 | 0.067 | 0.075 | 0.083 | 0.083 |
| Cumulative Prob. | 0.008 | 0.017 | 0.033 | 0.058 | 0.092 | 0.133 | 0.192 | 0.258 | 0.333 | 0.417 | 0.500 |
| W | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 |
| Frequency | 10 | 10 | 9 | 8 | 7 | 5 | 4 | 3 | 2 | 1 | 1 |
| Prob. | 0.083 | 0.083 | 0.075 | 0.067 | 0.058 | 0.042 | 0.033 | 0.025 | 0.017 | 0.008 | 0.008 |
| Cumulative Prob. | 0.583 | 0.667 | 0.742 | 0.808 | 0.867 | 0.908 | 0.942 | 0.967 | 0.983 | 0.992 | 1.000 |

Thus, for example, the probability that $W = 19$ is 0.075, with a frequency of 9 out of 120. From this table, we deduce that

$$\Pr[8 \leq W \leq 25] = 0.983 - 0.017 = 0.966$$

implying that the two-sided rejection region for $\alpha = 0.05$ is the set $\mathcal{R} = \{6, 7, 26, 27\}$.

EXAMPLE : Placenta Permeability Data

Using the placenta permeability data from Assignment 3, the data and ranks for are displayed below:

| | | | | | | | | | | | | | | | |
|-------|------|------|------|------|------|------|------|------|------|------|------|------|-----|------|------|
| Group | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 2 | 2 | 2 | 2 | 2 |
| Obs. | 0.73 | 0.80 | 0.83 | 1.04 | 1.38 | 1.45 | 1.46 | 1.64 | 1.89 | 1.91 | 0.74 | 0.88 | 0.9 | 1.15 | 1.21 |
| Rank | 1 | 3 | 4 | 7 | 10 | 11 | 12 | 13 | 14 | 15 | 2 | 5 | 6 | 8 | 9 |

Thus the Wilcoxon statistic is

$$W = R_2 = 2 + 5 + 6 + 8 + 9 = 30$$

Now, here $n_1 = 10$ and $n_2 = 5$. There are

$$\binom{15}{5} = \frac{15!}{10! 5!} = 3003$$

subsets of the ranks $\{1, 2, 3, \dots, 15\}$ of size 5.

In the permutation null distribution, the possible values of W are $\{15, 16, \dots, 64, 65\}$; the probabilities are given below.

| | | | | | | | | | | | | | |
|------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| W | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 |
| Frequency | 1 | 1 | 2 | 3 | 5 | 7 | 10 | 13 | 18 | 23 | 30 | 36 | 45 |
| Prob. | 0.000 | 0.000 | 0.001 | 0.001 | 0.002 | 0.002 | 0.003 | 0.004 | 0.006 | 0.008 | 0.010 | 0.012 | 0.015 |
| Cumulative Prob. | 0.000 | 0.001 | 0.001 | 0.002 | 0.004 | 0.006 | 0.010 | 0.014 | 0.020 | 0.028 | 0.038 | 0.050 | 0.065 |
| W | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 40 |
| Frequency | 53 | 63 | 72 | 83 | 92 | 103 | 111 | 121 | 127 | 134 | 137 | 141 | 141 |
| Prob. | 0.018 | 0.021 | 0.024 | 0.028 | 0.031 | 0.034 | 0.037 | 0.040 | 0.042 | 0.045 | 0.046 | 0.047 | 0.047 |
| Cumulative Prob. | 0.082 | 0.103 | 0.127 | 0.155 | 0.185 | 0.220 | 0.257 | 0.297 | 0.339 | 0.384 | 0.430 | 0.477 | 0.523 |
| W | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 |
| Frequency | 141 | 137 | 134 | 127 | 121 | 111 | 103 | 92 | 83 | 72 | 63 | 53 | 45 |
| Prob. | 0.047 | 0.046 | 0.045 | 0.042 | 0.040 | 0.037 | 0.034 | 0.031 | 0.028 | 0.024 | 0.021 | 0.018 | 0.015 |
| Cumulative Prob. | 0.570 | 0.616 | 0.661 | 0.703 | 0.743 | 0.780 | 0.815 | 0.845 | 0.873 | 0.897 | 0.918 | 0.935 | 0.950 |
| W | 54 | 55 | 56 | 57 | 58 | 59 | 60 | 61 | 62 | 63 | 64 | 65 | |
| Frequency | 36 | 30 | 23 | 18 | 13 | 10 | 7 | 5 | 3 | 2 | 1 | 1 | |
| Prob. | 0.012 | 0.010 | 0.008 | 0.006 | 0.004 | 0.003 | 0.002 | 0.002 | 0.001 | 0.001 | 0.000 | 0.000 | |
| Cumulative Prob. | 0.962 | 0.972 | 0.980 | 0.986 | 0.990 | 0.994 | 0.996 | 0.998 | 0.999 | 0.999 | 1.000 | 1.000 | |

By inspection of the table, we see that

$$\Pr[25 \leq W \leq 55] = 0.972 - 0.028 = 0.944$$

and

$$\Pr[24 \leq W \leq 56] = 0.980 - 0.020 = 0.960$$

Thus for a symmetric two-sided interval which contains at most probability 0.95, we take the interval

$$\{25, 26, \dots, 54, 55\}$$


and hence define the rejection region

$$\mathcal{R} = \{15, 16, 17, \dots, 23, 24, 56, 57, \dots, 64, 65\}$$

Note that this choice of rejection region ensures that there is at least probability 0.025 in each tail.

The permutation null distribution of W is displayed below.

Permutation Null Distribution with Normal Approximation



The normal approximation is given by

$$W \approx \text{Normal} \left(\frac{n_2(n_1 + n_2 + 1)}{2}, \frac{n_1 n_2 (n_1 + n_2 + 1)}{12} \right)$$